

## PROTEINS AND NUCLEIC ACIDS ENCODING SAME

### RELATED APPLICATIONS

This application claims priority from U.S.S.N. 60/241,040, filed October 17, 2000; U.S.S.N. 60/241,058, filed October 17, 2000; U.S.S.N. 60/241,063, filed October 17, 2000; U.S.S.N. 60/241,243 filed October 17, 2000; U.S.S.N. 60/242,152, filed October 20, 2000; U.S.S.N. 60/242,482, filed October 23, 2000; U.S.S.N. 60/242,611, filed October 23, 2000; U.S.S.N. 60/242,612, filed October 23, 2000; U.S.S.N. 60/242,880, filed October 24, 2000; U.S.S.N. 60/242,881, filed October 24, 2000; U.S.S.N. 60/259,028, filed December 29, 2000; U.S.S.N. 60/269,813, filed February 20, 2001; U.S.S.N. 60/286,324, filed April 25, 2001; U.S.S.N. 60/294,108, filed May 29, 2001; and U.S.S.N. 60/303,968, filed July 9, 2001 each of which is incorporated by reference in its entirety.

### FIELD OF THE INVENTION

The invention generally relates to nucleic acids and polypeptides encoded thereby.

### BACKGROUND OF THE INVENTION

The invention generally relates to nucleic acids and polypeptides encoded therefrom. More specifically, the invention relates to nucleic acids encoding cytoplasmic, nuclear, membrane bound, and secreted polypeptides, as well as vectors, host cells, antibodies, and recombinant methods for producing these nucleic acids and polypeptides.

### SUMMARY OF THE INVENTION

The invention is based in part upon the discovery of nucleic acid sequences encoding novel polypeptides. The novel nucleic acids and polypeptides are referred to herein as NOVX, or NOV1, NOV2, NOV3, NOV4, NOV5, NOV6, NOV7, NOV8, NOV9, and NOV10 nucleic acids and polypeptides. These nucleic acids and polypeptides, as well as derivatives, homologs, analogs and fragments thereof, will hereinafter be collectively designated as "NOVX" nucleic acid or polypeptide sequences.

In one aspect, the invention provides an isolated NOVX nucleic acid molecule encoding a NOVX polypeptide that includes a nucleic acid sequence that has identity to the

nucleic acids disclosed in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28. In some embodiments, the NOVX nucleic acid molecule will hybridize under stringent conditions to a nucleic acid sequence complementary to a nucleic acid molecule that includes a protein-coding sequence of a NOVX nucleic acid sequence. The invention also includes an isolated nucleic acid that encodes a NOVX polypeptide, or a fragment, homolog, analog or derivative thereof. For example, the nucleic acid can encode a polypeptide at least 80% identical to a polypeptide comprising the amino acid sequences of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, and 29. The nucleic acid can be, for example, a genomic DNA fragment or a cDNA molecule that includes the nucleic acid sequence of any of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28.

Also included in the invention is an oligonucleotide, *e.g.*, an oligonucleotide which includes at least 6 contiguous nucleotides of a NOVX nucleic acid (*e.g.*, SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28) or a complement of said oligonucleotide.

Also included in the invention are substantially purified NOVX polypeptides (SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, and 29). In certain embodiments, the NOVX polypeptides include an amino acid sequence that is substantially identical to the amino acid sequence of a human NOVX polypeptide.

The invention also features antibodies that immunoselectively bind to NOVX polypeptides, or fragments, homologs, analogs or derivatives thereof.

In another aspect, the invention includes pharmaceutical compositions that include therapeutically- or prophylactically-effective amounts of a therapeutic and a pharmaceutically-acceptable carrier. The therapeutic can be, *e.g.*, a NOVX nucleic acid, a NOVX polypeptide, or an antibody specific for a NOVX polypeptide. In a further aspect, the invention includes, in one or more containers, a therapeutically- or prophylactically-effective amount of this pharmaceutical composition.

In a further aspect, the invention includes a method of producing a polypeptide by culturing a cell that includes a NOVX nucleic acid, under conditions allowing for expression of the NOVX polypeptide encoded by the DNA. If desired, the NOVX polypeptide can then be recovered.

In another aspect, the invention includes a method of detecting the presence of a NOVX polypeptide in a sample. In the method, a sample is contacted with a compound that selectively binds to the polypeptide under conditions allowing for formation of a complex between the polypeptide and the compound. The complex is detected, if present, thereby identifying the NOVX polypeptide within the sample.

The invention also includes methods to identify specific cell or tissue types based on their expression of a NOVX.

Also included in the invention is a method of detecting the presence of a NOVX nucleic acid molecule in a sample by contacting the sample with a NOVX nucleic acid probe or primer, and detecting whether the nucleic acid probe or primer bound to a NOVX nucleic acid molecule in the sample.

In a further aspect, the invention provides a method for modulating the activity of a NOVX polypeptide by contacting a cell sample that includes the NOVX polypeptide with a compound that binds to the NOVX polypeptide in an amount sufficient to modulate the activity of said polypeptide. The compound can be, *e.g.*, a small molecule, such as a nucleic acid, peptide, polypeptide, peptidomimetic, carbohydrate, lipid or other organic (carbon containing) or inorganic molecule, as further described herein.

Also within the scope of the invention is the use of a therapeutic in the manufacture of a medicament for treating or preventing disorders or syndromes including, *e.g.*, adrenoleukodystrophy, congenital adrenal hyperplasia, leukodystrophies, breast cancer, Small-cell cancer of lung, squamous cell carcinomas, Colorectal cancer, Malignant melanoma, cutaneous, Neuroblastoma, Prostate cancer-brain cancer susceptibility, Alzheimer's disease, epilepsy, Huntington's disease, anxiety, ataxia-telangiectasia, behavioral disorders, multiple sclerosis, muscular dystrophy, myasthenia gravis, neurodegeneration, neuroprotection, Parkinson's disease, pain, stroke, Stroke, Aneurysm, Embolism, autoimmune disease, allergies, addiction, asthma, ARDS, allergy, endometriosis, endocrine dysfunctions, graft versus host disease, graft versus host disease (GVHD), idiopathic thrombocytopenic purpura, immunodeficiencies, IgA nephropathy, lymphoedema, systemic lupus erythematosus, scleroderma, transplantation, psoriasis, crohn's disease, HIV infection, Muscle-eye-brain disease, Neuropathy, paraneoplastic sensory, Charcot-Marie-Tooth neuropathy-2A, SCID due to LCK deficiency, bone marrow transplantation, Kostmann neutropenia, immunodeficiency, thrombocytopenia, eczema, lymphoid malignancy, impaired monocyte motility, Lymphoedema, atherosclerosis, aortic stenosis, atrial septal defect (ASD), atrioventricular (A-V) canal defect, ductus arteriosus, cerebral palsy, cirrhosis, cardiomyopathy, congenital heart defects, hypertension, hemophilia, hypercoagulation, pulmonary stenosis, subaortic stenosis, ventricular septal defect (VSD), valve diseases, bacterial and viral infections, cerebral vascular disease, osteoarthritis, rheumatoid arthritis, Heart block, nonprogressive, Heart block, progressive, 2, Ventricular fibrillation, idiopathic, entricular tachycardia, idiopathic, Thrombocytopenia, congenital amegakaryocytic, Bypass surgery, Bleeding disorders,

Adrenoleukodystrophy, Congenital Adrenal Hyperplasia, hypercoagulation, diabetes, obesity, metabolic disorders such as familial amyloidotic polyneuropathy, hyperkinetic diseases, Galactose epimerase deficiency, Glucose transport defect, blood-brain barrier, diverticular disease, emphysema, glomerulonephritis, hypercalcaemia, interstitial nephritis, inflammatory bowel disease, Lesch-Nyhan syndrome, polycystic kidney disease, pancreatitis, renal artery stenosis, renal tubular acidosis, tuberous sclerosis, Von Hippel-Lindau (VHL) syndrome, ulcers, neonatal apnea, eagle's syndrome, renal fibrogenesis, Meckel syndrome, skin disorders, connective tissue disorders such as type VIIC Ehlers-Danros syndrome, Primary bile acid malabsorption (PBAM, an idiopathic intestinal disorder), congenital diarrhea, steatorrhea, refractory infantile diarrhea, interruption of the enterohepatic circulation of bile acids, reduced plasma cholesterol levels, Hirschsprung's disease, Cirrhosis, growth failure, Aicardi-Goutieres syndrome 1, Brugada syndrome, Deafness, autosomal recessive 6, Ichthyosiforme erythroderma, congenital, nonbullous, Long QT syndrome-3, Night blindness, congenital stationary, Pituitary ACTH-secreting adenoma, Elliptocytosis-1, Fucosidosis, Hypophosphatasia (adult, childhood, infantile), Porphyria cutanea tarda, Porphyria, hepatoerythropoietic, Schwartz-Jampel syndrome, Myopathy due to succinate dehydrogenase deficiency, Bartter syndrome, type 3, Corneal dystrophy, crystalline, Schnyder, Hyperprolinemia, type II, erythrokeratoderma variabilis, palmoplantar keratoderma, diseases and disorders involving intercellular metabolic and electrical communication, diseases and disorders involving coordination, proliferation and differentiation, diseases and disorders involving maintenance of tissue homeostasis, growth control, development, and synchronized response of cells to stimuli, Wiskott-Aldrich syndrome, cytoskeletal abnormalities, trauma, tissue regeneration (in vitro and in vivo), respiratory disease, gastro-intestinal diseases, muscle, bone, joint and skeletal disorders, hematopoietic disorders, urinary system disorders, Tissue and organ transplantation, Fibromuscular dysplasia, Hyperparathyroidism, Hypoparathyroidism, Hyperthyroidism and Hypothyroidism, SIDS, Xerostomia, Tonsillitis, Osteoporosis, Ankylosing spondylitis, Scoliosis, Tendinitis, Dental disease and infection, growth and reproductive disorders, hypogonadism, fertility, and/or other pathologies and disorders of the like.

30           The therapeutic can be, *e.g.*, a NOVX nucleic acid, a NOVX polypeptide, or a NOVX-specific antibody, or biologically-active derivatives or fragments thereof.

For example, the compositions of the present invention will have efficacy for treatment of patients suffering from the diseases and disorders disclosed above and/or other pathologies and disorders of the like. The polypeptides can be used as immunogens to produce antibodies



specific for the invention, and as vaccines. They can also be used to screen for potential agonist and antagonist compounds. For example, a cDNA encoding NOVX may be useful in gene therapy, and NOVX may be useful when administered to a subject in need thereof. By way of non-limiting example, the compositions of the present invention will have efficacy for treatment of patients suffering from the diseases and disorders disclosed above and/or other pathologies and disorders of the like.

The invention further includes a method for screening for a modulator of disorders or syndromes including, *e.g.*, the diseases and disorders disclosed above and/or other pathologies and disorders of the like. The method includes contacting a test compound with a NOVX polypeptide and determining if the test compound binds to said NOVX polypeptide. Binding of the test compound to the NOVX polypeptide indicates the test compound is a modulator of activity, or of latency or predisposition to the aforementioned disorders or syndromes.

Also within the scope of the invention is a method for screening for a modulator of activity, or of latency or predisposition to disorders or syndromes including, *e.g.*, the diseases and disorders disclosed above and/or other pathologies and disorders of the like by administering a test compound to a test animal at increased risk for the aforementioned disorders or syndromes. The test animal expresses a recombinant polypeptide encoded by a NOVX nucleic acid. Expression or activity of NOVX polypeptide is then measured in the test animal, as is expression or activity of the protein in a control animal which recombinantly-expresses NOVX polypeptide and is not at increased risk for the disorder or syndrome. Next, the expression of NOVX polypeptide in both the test animal and the control animal is compared. A change in the activity of NOVX polypeptide in the test animal relative to the control animal indicates the test compound is a modulator of latency of the disorder or syndrome.

In yet another aspect, the invention includes a method for determining the presence of or predisposition to a disease associated with altered levels of a NOVX polypeptide, a NOVX nucleic acid, or both, in a subject (*e.g.*, a human subject). The method includes measuring the amount of the NOVX polypeptide in a test sample from the subject and comparing the amount of the polypeptide in the test sample to the amount of the NOVX polypeptide present in a control sample. An alteration in the level of the NOVX polypeptide in the test sample as compared to the control sample indicates the presence of or predisposition to a disease in the subject. Preferably, the predisposition includes, *e.g.*, the diseases and disorders disclosed above and/or other pathologies and disorders of the like. Also, the expression levels of the new

polypeptides of the invention can be used in a method to screen for various cancers as well as to determine the stage of cancers.

In a further aspect, the invention includes a method of treating or preventing a pathological condition associated with a disorder in a mammal by administering to the subject  
5 a NOVX polypeptide, a NOVX nucleic acid, or a NOVX-specific antibody to a subject (*e.g.*, a human subject), in an amount sufficient to alleviate or prevent the pathological condition. In preferred embodiments, the disorder, includes, *e.g.*, the diseases and disorders disclosed above and/or other pathologies and disorders of the like.

In yet another aspect, the invention can be used in a method to identify the cellular  
10 receptors and downstream effectors of the invention by any one of a number of techniques commonly employed in the art. These include but are not limited to the two-hybrid system, affinity purification, co-precipitation with antibodies or other specific-interacting molecules.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention  
15 belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods,  
20 and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

## DETAILED DESCRIPTION OF THE INVENTION

25 The present invention provides novel nucleotides and polypeptides encoded thereby. Included in the invention are the novel nucleic acid sequences and their encoded polypeptides. The sequences are collectively referred to herein as "NOVX nucleic acids" or "NOVX polynucleotides" and the corresponding encoded polypeptides are referred to as "NOVX polypeptides" or "NOVX proteins." Unless indicated otherwise, "NOVX" is meant to refer to  
30 any of the novel sequences disclosed herein. Table A provides a summary of the NOVX nucleic acids and their encoded polypeptides.

**TABLE A. Sequences and Corresponding SEQ ID Numbers**

NOVX Assignment	Internal Identification	SEQ ID NO (nucleic acid)	SEQ ID NO (polypeptide)	Homology
1a	SC138213196_A	1	2	Zinc Metalloprotease-like
1b	137043926_EXT_1	3	4	zinc metalloprotease-1-like
1c	CG52952-03	5	6	ADAM-TS 7-like
1d	CG52952-04	7	8	ADAM-TS 7-like
2	SC_78316254_A	9	10	Alpha-2-macroglobulin precursor-like
3	GMAC079237_A	11	12	Ileal Sodium/Bile Acid Cotransporter-like
4	AL161453_A	13	14	Prohibitin-like
5	dj1182a14_da1	15	16	Macrophage Stimulating Protein Precursor-like
6	GM382a20_A	17	18	Fatty Acid-Binding Protein-like
7	sggc_draft_dj895c5_2 0000819	19	20	Gap junction beta-5 protein-like
8	56072181_da1	21	22	Metallothionein-like
9	2855519_0_19_da1	23	24	CIP4-like
10a	129297354_EXT	25	27	hepsin/plasma transmembrane serine protease-like
10b	CG106783-02	28	29	Spinesin-like

NOVX nucleic acids and their encoded polypeptides are useful in a variety of applications and contexts. The various NOVX nucleic acids and polypeptides according to the invention are useful as novel members of the protein families according to the presence of domains and sequence relatedness to previously described proteins. Additionally, NOVX nucleic acids and polypeptides can also be used to identify proteins that are members of the family to which the NOVX polypeptides belong.

NOV1 is homologous to a Zinc Metalloprotease/ADAM-TS 7-like family of proteins. Thus, the NOV1 nucleic acids, polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example; adrenoleukodystrophy, Alzheimer's disease, autoimmune disease, allergies, addiction, anxiety, ataxia-telangiectasia, asthma, ARDS, atherosclerosis, behavioral disorders, aortic stenosis, atrial septal defect (ASD), atrioventricular (A-V) canal defect, ductus arteriosus, allergy, cerebral palsy, congenital adrenal hyperplasia, cirrhosis, cardiomyopathy, congenital heart defects, diabetes, diverticular disease, epilepsy, emphysema, endometriosis, endocrine dysfunctions, graft versus host disease, glomerulonephritis, graft versus host disease (GVHD), growth and reproductive disorders, hemophilia, hypercoagulation, hypercalcaemia, Huntington's disease, hypertension, hypogonadism, fertility, idiopathic thrombocytopenic purpura, immunodeficiencies, interstitial nephritis, IgA nephropathy, lymphoedema, inflammatory bowel disease, Lesch-Nyhan syndrome, leukodystrophies, multiple sclerosis, muscular dystrophy, myasthenia gravis, neurodegeneration, neuroprotection, obesity,

Parkinson's disease, pain, polycystic kidney disease, pulmonary stenosis, pancreatitis, renal artery stenosis, renal tubular acidosis, stroke, systemic lupus erythematosus, scleroderma, subaortic stenosis, transplantation, tuberous sclerosis, Von Hippel-Lindau (VHL) syndrome, ventricular septal defect (VSD), valve diseases, Von Hippel-Lindau (VHL) syndrome, ulcers,  
5 bacterial and viral infections, neonatal apnea, eagle's syndrome, atherosclerosis, metabolic disorders such as familial amyloidotic polyneuropathy, hyperkinetic diseases, muscular dystrophy, cerebral vascular disease, hypertension, cardiovascular diseases, renal fibrogenesis, inflammatory bowel disease, Meckel syndrome, colorectal cancer, papillomavirus infection and cervical carcinoma, liver malignancies, skin disorders,  
10 connective tissue disorders such as type VIIC Ehlers-Danros syndrome, osteoarthritis, rheumatoid arthritis, or other pathologie or conditions.

NOV2 is homologous to the Alpha-2-macroglobulin precursor-like family of proteins. Thus NOV2 nucleic acids, polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example;  
15 asthma, allergy and psoriasis, Alzheimer disease, Emphysema, pulmonary disease, immune disorders and Cancer and/or other pathologies and disorders.

NOV3 is homologous to a family of Ileal Sodium/Bile Acid Cotransporter-like proteins. Thus, the NOV3 nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated  
20 in, for example: Primary bile acid malabsorption (PBAM, an idiopathic intestinal disorder), congenital diarrhea, steatorrhea, refractory infantile diarrhea, interruption of the enterohepatic circulation of bile acids, reduced plasma cholesterol levels, crohn's disease, Inflammatory bowel disease, Diverticular disease, Hirschsprung's disease, Cirrhosis, Transplantation, Hypercalceimia, Ulcers, growth failure and/or other pathologies.

25 NOV4 is homologous to the Prohibitin-like family of proteins. Thus, NOV4 nucleic acids, polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example: breast cancer (In a sporadic breast cancer, Sato et al. (1992) found a missense mutation from valine (GTC) to alanine (GCC) at codon 88 of the PHB gene), and/or other pathologies.

30 NOV5 is homologous to the Macrophage Stimulating Protein Precursor-like family of proteins. Thus NOV5 nucleic acids, polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example: Aicardi-Goutieres syndrome 1, Brugada syndrome, Deafness, autosomal recessive 6, Heart block, nonprogressive, Heart block, progressive, 2, Ichthyosiforme

erythroderma, congenital, nonbullous, Long QT syndrome-3, Night blindness, congenital stationary, Pituitary ACTH-secreting adenoma, Small-cell cancer of lung, Ventricular fibrillation, idiopathic, entricular tachycardia, idiopathic, HIV infection, susceptibility/resistance to, Von Hippel-Lindau (VHL) syndrome, Cirrhosis, Transplantation, and/or other pathologies/disorders.

NOV6 is homologous to the Fatty Acid-Binding Protein-like family of proteins. Thus NOV6 nucleic acids, polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example: Cardiomyopathy, Atherosclerosis, Hypertension, Congenital heart defects, Aortic stenosis, Atrial septal defect (ASD), Atrioventricular (A-V) canal defect, Ductus arteriosus, Pulmonary stenosis, Subaortic stenosis, Ventricular septal defect (VSD), valve diseases, Tuberous sclerosis, Scleroderma, Transplantation, Endometriosis, Inflammatory bowel disease, Diverticular disease, Hirschsprung's disease, Crohn's Disease, Hemophilia, hypercoagulation, Idiopathic thrombocytopenic purpura, immunodeficiencies, Osteoporosis, Hypercalceimia, Arthritis, Von Hippel-Lindau (VHL) syndrome, Alzheimer's disease, Stroke, Tuberous sclerosis, hypercalceimia, Parkinson's disease, Huntington's disease, Cerebral palsy, Epilepsy, Asthma, allergy, ARDS, Lesch-Nyhan syndrome, Multiple sclerosis, Leukodystrophies, Behavioral disorders, Addiction, Anxiety, Pain, Neuroprotection Fertility, psoriasis, cancer including but not limited to basal and squamous cell carcinomas, obesity, diabetes, and/or other pathologies and disorders involving fatty acid transport of skin, oral mucosa as well as other organs, and/or other pathologies/disorders.

NOV7 is homologous to members of the Gap junction beta-5 protein-like family of proteins. Thus, the NOV7 nucleic acids, polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example; Deafness, autosomal dominant 2, Elliptocytosis-1, Fucosidosis, Hypophosphatasia (adult, childhood, infantile), Muscle-eye-brain disease, Neuropathy, paraneoplastic sensory, Porphyria cutanea tarda, Porphyria, hepatoerythropoietic, Schwartz-Jampel syndrome, Thrombocytopenia, congenital amegakaryocytic, Charcot-Marie-Tooth neuropathy-2A, Galactose epimerase deficiency, Glucose transport defect, blood-brain barrier, Kostmann neutropenia, Muscular dystrophy, congenital, with early spine rigidity, Myopathy due to succinate dehydrogenase deficiency, SCID due to LCK deficiency, Colorectal cancer, resistance to, Bartter syndrome, type 3, Breast cancer, ductal, Corneal dystrophy, crystalline, Schnyder, Hyperprolinemia, type II, Inflammatory bowel disease 7, Malignant melanoma, cutaneous, Neuroblastoma, Prostate cancer-brain cancer susceptibility, erythrokeratodermia

variabilis, palmoplantar keratoderma, diseases and disorders involving intercellular metabolic and electrical communication, diseases and disorders involving coordination, proliferation and differentiation, diseases and disorders involving maintenance of tissue homeostasis, growth control, development, and synchronized response of cells to stimuli, diseases and disorders involving the immune system, diseases and disorders involving regulation of bone cell differentiation, and/or other pathologies/disorders.

NOV8 is homologous to the Metallothionein-like family of proteins. Thus, NOV8 nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example; Gitelman syndrome, Menkes disease, Wilson's disease, acrodermatitis enteropathica, myelomonocytic leukemia, eosinophil disorders, hepatic disorders such as hepatic copper toxicity, and/or other pathologies/disorders.

NOV9 is homologous to the CDC-42 interacting protein 4-like family of proteins. Thus, NOV9 nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example; Wiskott-Aldrich syndrome, immunodeficiency, thrombocytopenia, eczema, lymphoid malignancy cytoskeletal abnormalities, impaired monocyte motility, Muscular dystrophy, Lesch-Nyhan syndrome, Myasthenia gravis, Von Hippel-Lindau (VHL) syndrome, Alzheimer's disease, Stroke, Tuberous sclerosis, hypercalcaemia, Parkinson's disease, Huntington's disease, Cerebral palsy, Epilepsy, Lesch-Nyhan syndrome, Ataxia-telangiectasia, Leukodystrophies, Behavioral disorders, Addiction, Anxiety, Pain, Neuroprotection, Fertility, Cardiomyopathy, Atherosclerosis, Hypertension, Congenital heart defects, Aortic stenosis, Atrial septal defect (ASD), Atrioventricular (A-V) canal defect, Ductus arteriosus, Pulmonary stenosis, Subaortic stenosis, Ventricular septal defect (VSD), valve diseases, Tuberous sclerosis, Scleroderma, Obesity, Transplantation, Diabetes, Pancreatitis, Obesity, Systemic lupus erythematosus, Autoimmune disease, Asthma, Emphysema, Scleroderma, allergy, ARDS, Cirrhosis, Transplantation, Diabetes, Autoimmune disease, Renal artery stenosis, Interstitial nephritis, Glomerulonephritis, Polycystic kidney disease, Renal tubular acidosis, IgA nephropathy, Hypercalcaemia, and/or other pathologies/disorders.

NOV10 is homologous to the hepsin/plasma transmembrane serine protease/spinesin-like family of proteins. Thus, NOV10 nucleic acids and polypeptides, antibodies and related compounds according to the invention will be useful in therapeutic and diagnostic applications implicated in, for example; cancer, trauma, tissue regeneration (in vitro and in vivo), viral/bacterial/parasitic infections, immunological disease, respiratory disease, gastro-intestinal

diseases, reproductive health, neurological and neurodegenerative diseases, bone marrow transplantation, metabolic and endocrine diseases, allergy and inflammation, nephrological disorders, cardiovascular diseases, muscle, bone, joint and skeletal disorders, hematopoietic disorders, urinary system disorders, Tissue and organ transplantation, Cardiomyopathy,

5 Atherosclerosis, Hypertension, Congenital heart defects, Aortic stenosis, Atrial septal defect (ASD), Atrioventricular (A-V) canal defect, Ductus arteriosus, Pulmonary stenosis, Subaortic stenosis, Ventricular septal defect (VSD), valve diseases, Scleroderma, Obesity, Hypertension, Fibromuscular dysplasia, Stroke, Aneurysm, Myocardial infarction, Embolism, Bypass surgery, Anemia, Bleeding disorders, Adrenoleukodystrophy, Congenital Adrenal

10 Hyperplasia, Diabetes, Von Hippel-Lindau (VHL) syndrome, Pancreatitis, Hyperparathyroidism, Hypoparathyroidism, Hyperthyroidism and Hypothyroidism, SIDS, Endometriosis, infertility, Xerostomia, Hypercalcaemia, Ulcers, Cirrhosis, Inflammatory bowel disease, Diverticular disease, Hirschsprung's disease, Crohn's Disease, Appendicitis, Hemophilia, hypercoagulation, autoimmune disease, allergies, immunodeficiencies,

15 transplantation, Graft versus host disease (GVHD), Ataxia-telangiectasia, Autoimmune disease, Hemophilia, Hypercoagulation, Idiopathic thrombocytopenic purpura, Immunodeficiencies, Lymphedema, Allergies, Hemophilia, hypercoagulation, idiopathic thrombocytopenic purpura, Lymphoedema, Tonsillitis, Osteoporosis, Hypercalcaemia, Arthritis, Ankylosing spondylitis, Scoliosis, Tendinitis, Muscular dystrophy, Lesch-Nyhan

20 syndrome, Myasthenia gravis, Dental disease and infection, Alzheimer's disease, Stroke, Tuberous sclerosis, hypercalcaemia, Parkinson's disease, Huntington's disease, Cerebral palsy, Epilepsy, Lesch-Nyhan syndrome, Multiple sclerosis, Ataxia-telangiectasia, Behavioral disorders, Addiction, Anxiety, Pain, Neuroprotection, Endocrine dysfunctions, Growth and reproductive disorders, Myasthenia gravis, Leukodystrophies, Pain, Neuroprotection, Systemic

25 lupus erythematosus, Autoimmune disease, Emphysema, Scleroderma, ARDS, Pharyngitis, Laryngitis, Asthma, Hearing loss, Tinnitus, Psoriasis, Actinic keratosis, Tuberous sclerosis, Acne, Hair growth, alopecia, pigmentation disorders, endocrine disorders, cystitis, incontinence, Autoimmune disease, Renal artery stenosis, Interstitial nephritis, Glomerulonephritis, Polycystic kidney disease, Systemic lupus erythematosus, Renal tubular

30 acidosis, IgA nephropathy, Hypercalcaemia, Lesch-Nyhan syndrome, Vesicoureteral reflux, and/or other pathologies/disorders.

The NOVX nucleic acids and polypeptides can also be used to screen for molecules, which inhibit or enhance NOVX activity or function. Specifically, the nucleic acids and polypeptides according to the invention may be used as targets for the identification of small

molecules that modulate or inhibit, *e.g.*, neurogenesis, cell differentiation, cell proliferation, hematopoiesis, wound healing and angiogenesis.

Additional utilities for the NOVX nucleic acids and polypeptides according to the invention are disclosed herein.

## 5 NOV1

NOV1 includes three novel zinc metalloprotease/ADAM-TS 7-like proteins disclosed below. The disclosed sequences have been named NOV1a, NOV1b, NOV1c, and NOV1d.

### NOV1a

A disclosed NOV1a nucleic acid of 2997 nucleotides (also referred to as SC138213196  
10 \_A) encoding a novel Zinc Metalloprotease-like protein is shown in Table 1A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 10-12 and ending with a TAA codon at nucleotides 2968-2970. A putative untranslated region upstream from the initiation codon and downstream from the termination codon is underlined in Table 1A. The start and stop codons are in bold letters.

15

Table 1A. NOV1a nucleotide sequence (SEQ ID NO:1).

```
CGCTCCTGGATGAAGCCCCGCGCGCGGATGGCGGGGCTTGGCGGCGCTGTGGATGCTGTTGGCGCAGGTG
GCCGAGCAGGTGAGTCCCGGGCGCTCCACAGCGCGGAAACCGGGTCCGGACAGCTGGAGGCGAGTCCC
CCGCGGCTCCTCTCCCGGGACCCCGCGTCTCACCGCGATGTCGCGCTGTTTTCCGCGAGCACCTGCGTG
CGCCATGGGACCCGAGCGGAGCGCTGGGAGCCGAGCGTCCCGCGTCTCCTCCACCCGCGGAGCGGCC
GGGCTGGATGGAAAGGGCGGGACATGGATGAAGCTGGAAACCATCGTTCTCAGCAAACCTAACACAGGAACA
GAAACCAAACTGCATGTTCTCACTCAATATGACCTGGTCTCTGCCTACGAGGTTGACCACAGGGGCGAT
TACGTGTCCCATGAAATCATGCACCATCAGCGGCGGAGAAGAGCAGTGCCCGTGTCCGAGGTTGAGTCTCTT
CACCTTCGGCTGAAAGGCCCGGACGACTTCCACATGGATCTGAGGACTTCCAGCAGCCTAGTGGCTCCT
GGCTTTATGTGCAGACGTTGGGAAAGACAGGCACTAAGTCTGTGCAGACTTACCGCCAGAGGACTTCTGT
TTCTATCAAGGCTCTTTGCGATCACACAGAACTCGCCATCGCATGGAGGGAAGTTCTGTGAGGGCTCCACT
CGCACTCTGAAGCTCTGCAACAGTCAGAAATGTCCCGGGACAGTGTGACTTCCGTGCTGCTCGACTGTGCC
GAGCACACAGCAGACGATTGAGAGGCGGCACTACAAGTGAAGCCTTACACTCAAGTAGAAGNNGACTTA
TGCAAACTCTACTGTATCGCAGAAGGATTTGATTCTTCTTTCTTTGTCAAATAAAGTCAAAGATGGGACT
CCATGCTCGGAGGATAGCCGTAATGTTGTATAGATGGGATATGTGAGCTCAGTGTGGTGTCCACATCTGCG
CACATGCCCCAGCCTCCCAAGGAAGACCTCTTCATCTTGCCAGATGAGTATAAGTCTTGCTTACGGCATAAG
CGCTCTCTTCTGAGGTCCCATAGAAATGAAGAACTGAACGTGGAGACCTTGGTGGTGTGCGACAAAAGATG
ATGCAAAACCATGGCCATGAAAATATCACCACTACGTGCTCACGATACTCAACATGGTATCTGCTTTATTC
AAAGATGGAACAATAGGAGGAAACATCAACATTGCAATTGTAGGTCTGATTCTCTAGAAGATGAACAGCCA
GGACTGGTGATAAGTCAACACGCAGACCACACCTTAAGTAGCTTCTGCCAGTGGCAGTCTGGATTGATGGGG
AAAGATGGGACTCGTCATGACCACGCCATCTTACTGACTGGTCTGGATATATGTTCTTGAAGAATGAGCCC
TGTGACACTTTGGGATTTGCACCCATAAGTGGAATGTGTAGTAAATATCGCAGCTGCACGATTAATGAAGAT
ACAGGTCTTGGACTGGCCTTCAACATTGCCCATGAGTCTGGACACAACCTTTGGCATGATTGATGAGAGAA
GGGAACATGTGTAAAAAGTCCGAGGGCAACATCATGTCCCTACATTGGCAGGACGCAATGGAGTCTTCTCC
TGGTCACCCTGCAGCCGCCAGTATCTACAAAATTTCTAAGCACCGCTCAAGCTATCTGCCTTGCTGATCAG
CCAAAGCCTGTGAAGGAATACAAGATATCTGAGAAATTGCCAGGAGAATTATATGATGCAAAACACAGATGC
AAGTGGCAGTTTCGGAGAGAAAGCCAAGCTCTGCATGCTGGACTTTAAAAAGGACATCTGTAAAGCCCTGTGG
TGCCATCGTATTGGAAGGAAATGTGAGACTAAATTTATGCCAGCAGCAGAAGGCACAATTTGTGGGCATGAC
ATGTGGTGCCGGGGAGGACAGTGTGTGAAATATGGTGATGAAGGCCCAAGCCCAACCATGGCCACTGGTCCG
GACTGGTCTTCTTGGTCCCCATGCTCCAGGACCTGCGGAGGGGGAGTATCTCATAGGAGTCGCCCTCTGCACC
AACCCCAATCCATCGCATGGAGGGAAGTTCTGTGAGGGCTCCACTCGCACTCTGAAGCTCTGCAACAGTCAG
AAATGTCCCCGGGACAGTGTGACTTCCGTGCTGCTCAGTGTGCCGAGCACAAACAGCAGACGATTGAGAGGG
CGGCACTACAAGTGAAGCCTCAGGACTTATGCAAACTCTACTGTATCGCAGAAGGATTTGATTCTTCTTT
TCTTTGTCAAATAAAGTCAAAGATGGGACTCCATGCTCGGAGGATAGCCGTAATGTTTGTATAGATGGGATA
TGTGAGNTTGGATGTGACAAATGTCCTTGGATCTGATGCTGTTGAAGACGTCTGTGGGGTGTGTAACGGGAAT
AACTCAGCCTGCACGATTACAGGGGTCTCTACCAAGCACCACACCAACCAATTATCATATGGTCACC
```



```

ATTCTTCTGGAGCCCGGAGTATCCGCATCTATGAAATGAACGTCTCTACCTCCTACATTTCTGTGCGCAAT
GCCCTCAGAAGGTACTACCTGAATGGGCACTGGACCGTGGACTGGCCCGGCCGGTACAAATTTTCGGGCACT
ACTTTCGACTACAGACGGTCCTATAATGAGCCCGAGAACTTAATCGCTACTGGACCAACCAACGAGACACTG
ATTGTGGAGCTGCTGTTTCAGGGAAGGAACCCGGGTGTTGCCTGGGAATACTCCATGCCTCGCTTGGGGACC
GAGAAGCAGCCCCCTGCCAGCCAGCTACACTTGGGCCATCGTGCGCTCTGAGTGCTCCGTGCTCGCGGA
GGGGGTAGGTGCCTTCCAGTGCTGCTCCTGGAGGCAGCATGTGAGCCTTCAGCCACTGCGTACATTGCACTG
GCCTTTCTGAATCCTAATGAGCAGCCCGNGCTTCTCCTGCCA

```

In a search of public sequence databases, the NOV1a nucleic acid sequence, located on chromosome 5 has 250 of 375 bases (66%) identical to a zinc metalloprotease ADAMTS6 (ADAMTS6) mRNA from *Homo sapiens* (GENBANK-ID: AF140674). Public nucleotide databases include all GenBank databases and the GeneSeq patent database.

In all BLAST alignments herein, the "E-value" or "Expect" value is a numeric indication of the probability that the aligned sequences could have achieved their similarity to the BLAST query sequence by chance alone, within the database that was searched. For example, the probability that the subject ("Sbjct") retrieved from the NOV1 BLAST analysis, e.g., thioredoxin mRNA from *Ovis aries*, matched the Query NOV1 sequence purely by chance is  $9.4e^{-25}$ . The Expect value (E) is a parameter that describes the number of hits one can "expect" to see just by chance when searching a database of a particular size. It decreases exponentially with the Score (S) that is assigned to a match between two sequences. Essentially, the E value describes the random background noise that exists for matches between sequences.

The Expect value is used as a convenient way to create a significance threshold for reporting results. The default value used for blasting is typically set to 0.0001. In BLAST 2.0, the Expect value is also used instead of the P value (probability) to report the significance of matches. For example, an E value of one assigned to a hit can be interpreted as meaning that in a database of the current size one might expect to see one match with a similar score simply by chance. An E value of zero means that one would not expect to see any matches with a similar score simply by chance. See, e.g., <http://www.ncbi.nlm.nih.gov/Education/BLASTinfo/>. Occasionally, a string of X's or N's will result from a BLAST search. This is a result of automatic filtering of the query for low-complexity sequence that is performed to prevent artifactual hits. The filter substitutes any low-complexity sequence that it finds with the letter "N" in nucleotide sequence (e.g., "NNNNNNNNNNNNNN") or the letter "X" in protein sequences (e.g., "XXXXXXXXXX"). Low-complexity regions can result in high scores that reflect compositional bias rather than significant position-by-position alignment. (Wootton and Federhen, *Methods Enzymol* 266:554-571, 1996).

The disclosed NOV1a polypeptide (SEQ ID NO:2) encoded by SEQ ID NO:1 has 986 amino acid residues and is presented in Table 1B using the one-letter amino acid code. Signal P, Psort and/or Hydropathy results predict that NOV1a has a signal peptide and is likely to be localized in extracellularly with a certainty of 0.5469. In other embodiments, NOV1a may also be localized to the lysosome (lumen) with a certainty of 0.1900, the microbody (peroxisome) with a certainty of 0.1297 or in the endoplasmic reticulum (membrane) with a certainty of 0.1000. The most likely cleavage site for a NOV1a peptide is between amino acids 26 and 27, at: VAE-QV.

**Table 1B. Encoded NOV1a protein sequence (SEQ ID NO:2).**

```

MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQRCNRRSGQLEASPPRLLSRGPRRLTAMSPLFSACTCVRHG
TRSGSAWEPPERPASSSTRGAAGLDGKGRDMDEAGNHRSSQQTNTGTENQTLHVLTQYDLVSAEVDHRGDYVS
HEIMHHQRRRRRAVAVSEVESLHLRLKGRHDFHMDLRTSSSLVAPGFIVQTLGKTGTSKVQTLPPEDFCFYQ
GSLRSHRNSPSHGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKWKPYTQVEXDLCKL
YCIAEGFDFFFLSNKVKDGTPCSEDSRNVICIDGICELSVVSTSAHMPQPPKEDLFILPDEYKSLRHKRSL
LRSHRNEELNVETLVVVDDKMMQNHGHEINITTYVLTILNMVSALFKDGTIGGNINIAIVGLILLEDEQPGLV
ISHHADHTLSSFCQWQSGLMGKDGTRHDHAILLTGLDICSWKNEPCDTLGFAPISGMCSKYRSTINEDTGL
GLAFTIAHESGHNFMIHDGEGNMCKKSEGNIMSPTLAGRNGVFSWSPCSRQYLHKFLSTAQAICLADQPKP
VKEYKYPEKLPGLYDANTQCKWQFGEKAKLCMLDFKKDICKALWCHRIGRKCETKFMPPAAEGTICGHDMWC
RGGQCVKYGDEGPKPTHGHWSWSSWSPCSRTCGGVSHRSRLCTNPNSHGGKFCEGSTRTLKLCNSQKCP
RDSVDFRAAQCAEHNSRRFRGRHYKWKPDLCCLYICIAEGFDFFFLSNKVKDGTPCSEDSRNVICIDGICEX
GCDNVLGSDAEDVCGVCNGNNSACTIHRGLYTKHHHTNHYHMTIPSGARSIRIYEMNVSTSYISVRNALR
RYYLNGHWTVDWPGRYKFSGTTFDYRRSYNEPENLIATGPTNETLIVELLFQGRNPGVAWEYSMPRLGTEKQ
PPAQPSYTWAIVRSECSVSCGGGRCLPVLLLEAACQPSATAYIALAFLES

```

A search of sequence databases reveals that the NOV1a amino acid sequence has 257 of 579 amino acid residues (44%) identical to, and 356 of 579 amino acid residues (61%) similar to, the 997 amino acid residue Zinc Metalloprotease Adamts7 protein from *Homo sapiens* (Human) (Q9UKP4) ( $E = 6.1 \times 10^{-149}$ ). Public amino acid databases include the GenBank databases, SwissProt, PDB and PIR.

NOV1a is expressed in the lung.

#### NOV1b

A disclosed NOV1b nucleic acid of 2433 nucleotides (also referred to as 137043926\_EXT\_1) encoding a novel zinc metalloprotease-1-like protein is shown in Table 1C. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 31-33 and ending with a TAA codon at nucleotides 2404-2406. A putative untranslated regions upstream from the initiation codon and downstream of the termination codon are underlined in Table 1C. The start and stop codons are in bold letters.

**Table 1C. NOV1b nucleotide sequence (SEQ ID NO:3).**

GTGGCCCCCTAGCCCCCTCGGAGCGCTCCTGGATGAAGCCCCGCGCGCGGATGGCGGGGCTTGGCGGCGCTG  
TGGATGCTGCTGGCGCAGGTGGCCGAGCAGGTGAGTCCCGGGCGCTCCACAGCGCGGAAACCGCGGGTCC  
GGACAGCTGGAGCGAGTCCCCCGCGCTCCTCTCCCGCGGACCCCGCGCTCTACCCGCGATGTCGCGCGTG  
TTTTCCGAGGCACCTGCGTGCGCCATGGGACCCGAGCGGCAGCGCCTGGGAGCCCGAGCGTCCCGCGTCC  
TCCTCCACCCCGGAGCGCGCGGGCTGGATGGAAAAGGCGGGACATGGATGAAGCTGGAAACCATCGTTCT  
CAGCAAATAACACAGGAACAGAAAACCAAACACTGCATGTTCTCACTCGTGAATATGACCTGGTCTCTGCC  
TACGAGTTGACCACAGGGGCGATTACGTGTCCCATGAAATCATGCACCATCAGCGGCGGAGAAGAGCAGTG  
GCCGTGTCCGAGGTTGAGTCTCTTACCTTCGGCTGAAAGGCCCGAGGCACGACTTCCACATGGATCTGAGG  
ACTTCCAGCAGCCTAGTGGCTCCTGGCTTTATTGTGCAGACGTTGGGAAAGACAGGCATAAGTCTGTGCAG  
ACTTTACCGCCAGAGGACTTCTGTTTCTATCAAGGCTCTTTCGATCACACAGAACTCCCCATCGCATGGA  
GGGAAGTTCTGTAGGGCTCCACTCGCACTCTGAAGCTCTGCAACAGTCAGAAATGTCCCGGGACAGTGTT  
GACTTCCGTGCTGCTCAGTGTGCCGAGCACAAACAGCAGACGATTAGAGGGCGGCATACAAGTGGAAAGCCT  
TACACTCAAGTAGAACAGGACTTATGCAAACCTACTGTATCGCAGAAGGATTTGATTTCTTCTTTCTTTG  
TCAAATAAAGTCAAAGATGGGACTCCATGCTCGGAGGATAGCCGTAATGTTTGTATAGTGGATATGTGAG  
ATGCCCCAGCCTCCCAAGGAAGACCTTTCATCTTGCCAGATGAGTATAAGTCTTGCTTACGGCATAAGCGC  
TCTCTTCTGAGGTCCCATAGAAATGAAGAACTGAACGTGGAGACCTTGGTGGTGGTCGACAAAAAGATGATG  
CAAAACCATGGCCATGAAAATATCACACCTACGTGCTCAGGATCTCAACATGGTATCTGCTTTATTCAA  
GATGGAACAATAGGAGGAACATCAACATTGCAATTGTAGGTCTGATTCTTCTAGAAGATGAACAGGACATC  
GTAAAGCCCTGTGGTGCCATCGTATTGGAAGGAAATGTGAGACTAAATTTATGCCAGCAGAGAAGGCACA  
ATTTGTGGCATGACATGTGGTGCCGGGGAGGACAGTGTGTGAAATATGGTGATGAAGCCCCAAGCCCACC  
CATGGCCACTGGTCGGACTGGTCTTCTTGGTCCCATGTCCAGGACCTGCGGAGGGGAGTATCTCATAGG  
AGTCGCCTCTGCACCAACCCAGGCCATCGCATGGAGGGAAGTTCTGTGAGGGCTCCACTCGCACTCTGAAG  
CTCTGCAACAGTCAGAAATGTCCCGGGACAGTGTGACTTCCGTGCTGCTCAGTGTGCCGAGCACAAACAGC  
AGACGATTTCAGAGGGCGGCCTACAAGTGGAAAGCCTCAGGACTTATGCAAACCTCTACTGTATCGCAGAAGGA  
TTTGATTTCTTCTTTCTTTGTCAAATAAAGTCAAAGATGGGACTCCATGCTCGGAGGATAGCCGTAATGTT  
TGTATAGATGGGATATGTGAGGGATGTGACAAATGCTTGGATCTGATGCTGTTGAAGACGTCGTGGGGTG  
TGTAACGGGAATAACTCAGCCTGCACGATTACAGGGGTCTCTACACCAAGCACCACCACTATTAT  
CACATGGTCACCATTCCTTCTGGAGCCCGAGTATCCGATCTATGAAATGAACGTCCTTACCTCTACATT  
TCTGTGCGCAATGCCCTCAGAAGGTACTACCTGAATGGGCACTGGACCGTGGACTGGCCCGCGCGGTACAAA  
TTTTCGGGCACTACTTTCGACTACAGACGGTCTTATAATGAGCCCGAGAACTTAATCGCTACTGGACCAACC  
AACGAGACACTGATTGTGGAGCTGCTGTTTTCAGGGAAGGAACCCGGGTGTTGCCTGGGAATACTCCATGCCT  
CGCTTGGGGACCGAGAAGCAGCCCCCTGCCCAGCCCAGCTACACTTGGGCCATCGTGCGCTCTGAGTGCTCC  
GTGTCCTGCGGAGGGGTAGGTGCCTTCCAGTGCTGCTCCTGGAGGCAGCATGTGAGCCTTACGCACTGCG  
TACATTGCACTGGCCTTCTTGAATCCTAATGAGCAGCCCGGGCTTCTCCCTGCCA

In a search of public sequence databases, the NOV1b nucleic acid sequence, located on chromosome 5 has 101 of 126 bases (80%) identical to agb:GENBANK-

ID:HSA400877|acc:AJ400877.1 mRNA from *Homo sapiens* (*Homo sapiens* ASCL3 gene,

5 CEGP1 gene, C11orf14 gene, C11orf15 gene, C11orf16 gene and C11orf17 gene) ( $E = 2.3e^{-7}$ ).

Public nucleotide databases include all GenBank databases and the GeneSeq patent database.

The disclosed NOV1b polypeptide (SEQ ID NO:4) encoded by SEQ ID NO:3 has 791 amino acid residues and is presented in Table 1D using the one-letter amino acid code. Signal P, Psort and/or Hydropathy results predict that NOV1b has a signal peptide and is likely to be  
10 localized in extracellularlye with a certainty of 0.5469. In other embodiments, NOV1b may also be localized to the lysosome (lumen) with acertainty of 0.1900, the microbody (peroxisome) with a certainty of 0.1144 or in the endoplasmic reticulum (membrane) with a certainty of 0.1000. The most likely cleavage site for a NOV1b peptide is between amino acids 23 and 24, at: VAE-QV.

**Table 1D. Encoded NOV1b protein sequence (SEQ ID NO:4).**

MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQRGNRSGGLEASPPRLLSRGPRRLTAMSPLFSAGTCVRHG  
 TRSGSAWEFERPASPSTRGAAGLDGKGRDMDEAGNHRSQQTNTGTENQTLHVLTREYDLVSAYEVDHRGDYV  
 SHEIMHHQRRRRRAVAVSEVESLHLRLKGRHDFHMDLRTSSSLVAPGFIVQTLGKTGTSVQTLPPEDFCFY  
 QGSLRSHRNSPSHGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKWKPYTQVEQDLCK  
 LYCIAEGFDFFFSLSNKVKDGTPCSEDSRNVCIDGICEMPQPPKEDLFILPDEYKSLRHKRSLRSHRNEE  
 LNVETLVVVDKKMMQNHGHENITTYVLTILNMVSALFKDGTIGGNINIAIVGLILLEDEQDICKALWCHRIG  
 RKCETKFMPAEAGTICGHDMMCRGGQCVKYGDEGPKPTHGHWSWSSWSPCSRTCGGGVSHRSRLCTNPRPS  
 HGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKWKPDLCCKLYCIAEGFDFFFSLSNK  
 VKDGTPCSEDSRNVCIDGICGCDNVLGSDAVEDVCGVCNGNNSACTIHRGLYTKHHHTNYHYMVTIPSGAR  
 SIRIYEMNVSTSYISVRNALRRYYLNGHWTVDWPGRYKFSGTTFDYRRSYNEPENLIATGPTNETLIVELLF  
 QGRNPGVAWEYSMPRLGTEKQPPAQPSYTWAIVRSECSVSCGGGRCLPVLLLEAACQPSATAYIALAFLES

A search of sequence databases reveals that the NOV1b amino acid sequence has 152 of 357 amino acid residues (42%) identical to, and 216 of 357 amino acid residues (60%) similar to, the 860 amino acid residue ptnr:SWISSNEW-ACC:Q9UKP5 protein from *Homo sapiens* (Human) (ADAM-TS 6 precursor (EC 3.4.24.-) (A Disintegrin And Metalloproteinase With Thrombospondin Motifs 6) (ADAMTS-6) (ADAM-TS6) ( $E = 4.8e^{-101}$ ). Public amino acid databases include the GenBank databases, SwissProt, PDB and PIR.

NOV1b is expressed in at least the following tissues: brain, liver, spleen, uterus, colon, tonsil, lung, germ cells. This information was derived by determining the tissue sources of the sequences that were included in the invention including but not limited to SeqCalling sources, Public EST sources, Genomic Clone sources, Literature sources, and/or RACE sources.

#### NOV1c

In the present invention, the target sequence identified previously, NOV1a, was subjected to the exon linking process to confirm the sequence. PCR primers were designed by starting at the most upstream sequence available, for the forward primer, and at the most downstream sequence available for the reverse primer. In each case, the sequence was examined, walking inward from the respective termini toward the coding sequence, until a suitable sequence that is either unique or highly selective was encountered, or, in the case of the reverse primer, until the stop codon was reached. Such primers were designed based on in silico predictions for the full length cDNA, part (one or more exons) of the DNA or protein sequence of the target sequence, or by translated homology of the predicted exons to closely related human sequences sequences from other species. These primers were then employed in PCR amplification based on the following pool of human cDNAs: adrenal gland, bone marrow, brain - amygdala, brain - cerebellum, brain - hippocampus, brain - substantia nigra, brain - thalamus, brain - whole, fetal brain, fetal kidney, fetal liver, fetal lung, heart, kidney, lymphoma - Raji, mammary gland, pancreas, pituitary gland, placenta, prostate, salivary

gland, skeletal muscle, small intestine, spinal cord, spleen, stomach, testis, thyroid, trachea, uterus. Usually the resulting amplicons were gel purified, cloned and sequenced to high redundancy. The resulting sequences from all clones were assembled with themselves, with other fragments in CuraGen Corporation's database and with public ESTs. Fragments and ESTs were included as components for an assembly when the extent of their identity with another component of the assembly was at least 95% over 50 bp. In addition, sequence traces were evaluated manually and edited for corrections if appropriate. These procedures provide the sequence reported below, which is designated NOV1c. This differs from the previously identified sequence NOV1a in having a different N-terminus

A disclosed NOV1c nucleic acid of 2902 nucleotides (also referred to as CG52952-03) encoding a novel ADAM-TS 7-like protein is shown in Table 1E. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 182-184 and ending with a TAA codon at nucleotides 2750-2752. A putative untranslated region upstream from the initiation codon and downstream from the termination codon is underlined in Table 1E. The start and stop codons are in bold letters.

**Table 1E. NOV1c nucleotide sequence (SEQ ID NO:5).**

TTGGCGGCGCTGTGGATGCTGCTGGCGCAGGTGGCCGAGCAGGTGAGTCCCGGGCGCTCCCACCAGCGCGGA  
AACC CGGGTCCGGACAGCTGGAGGCGAGTCCCCCGGGCTCCTCTCCCGCGGACCCGCGCTCACC CGG  
ATGTCGCGCGCTGTTTCCG CAGGCACCTGCGTGCGCCATGGGACCCG CAGCGGCAGCGCTGGGAGCCCGAG  
CGTCCCGCGTCTCTCCAGACCTGGTCTCTGCTACGAGGTTGACCACAGGGGCGATTACGTGTCCCATGA  
AATCATGCACCATCAGCGGCGGAGAAGAGCAGTGGCCGTGTCCGAGGTTGAGCCAGCCTTTCTCCAGGTATG  
CAGAGCCAGAGAGCTCAGACTGTGTGTGGAGGCTTTCCCATTTGCTAATTCTCAGCCGGGTTTTTGAACCT  
TTCCAATGTTTCGCTCTCAGTGGAGGAACAGCATGCTTCCAAGAGAATAATAACAAATGCAATGCTTGGAGA  
ATCGGCCCTGGCTTCAACCAGAAAGTCTAATTGTGTTTTCTTTCTTCTTTATTTTTTCCAGTCAGGCAT  
GATACGAACAGAAGAGGCAGATTACTTCTTAAGGCCACTTCTTCACACCTCTCATGGAACTCGGCAGAGC  
TGCCCAAGGCAGCTCGCCATCCACGTA CTGTACAAGAGAGAGGTCCTGGTGACCTCAAGGACATGGGAGCT  
GGCACATCAACCCCTGCACAGCAGCGACCTTCGCTGGGACTGCCACAAAAGCAGCATTCTGTGGAAGACG  
CAAGAAATACATGCCCCAGCCTCCCAAGGAAGACCTCTTCATCTTGCCAGATGAGTATAAGTCTTGCTTACG  
GCATAAGCGCTCTCTTCTGAGGTCCCATAGAAATGAAGAACTGAACGTGGAGACCTTGGTGGTGGTGCACAA  
AAAGATGATGCAAAACCATGGCCATGAAAATATCACCACCTACGTGCTCAGGATACTCAACATGGTATCTGC  
TTTATTCAAAGATGGAACAATAGGAGGAAACATCAACATTGCAATTGTAGGTCTGATTCTTCTAGAAGATGA  
ACAGCCAGGACTGGTGATAAGTCACACGCAGACCACACCTTAAGTAGTTTTCTGCCAGTGGCAGTCTGGATT  
GATGGGGAAAGATGGGACTCGTCATGACCACGCCATCTTACTGACTGGTCTGGATATATGTTCTCTGGAAGAA  
TGAGCCCTGTGACACTTTGGGATTTCACCCATAAGTGAATGTGTAGTAAATATCGCAGCTGCACGATTAA  
TGAAGATACAGGCTCTGGACTGGCCTTCACCATTTGCCCATGAGTCTGGACACAACCTTTGGCATGATTCTGA  
TGGAGAAGGGAACATGTGCAAAAAGTCCGAGGGCAACATCATGTCCCTACATTGGCAGGACGCAATGGAGT  
CTTCTCCTGGTCACCTGCAGCCGCCAGTATCTACACAAATTTCTAAGCACCCTCAAGCTATCTGCCTTGC  
TGATCAGCCAAAGCCTGTGAAGGAATACAAGTATCTTGAGAAATTGCCAGGAGAATTATATGGTGCAACAC  
ACAGTGCAAGTGGCAGTTCGGAGAGAAAGCCAAGCTCTGCATGCTGGACTTTAAAAGGACATCTGTAAAGC  
CCTGTGGTGCCATCGTATTGGAAGGAATGTGAGACTAAATTTATGCCAGCAGCAGAAGGCACAATTTGTGG  
GCATGAACATGGTGCCGGAGGACAGTGTGTGAAATATGGTGATGAAGGCCCAAGCCCAACCATGGCCACTG  
GTCGGACTGGTCTTCTTGGTCCCCATGCTCCAGGACCTGCGGAGGGGAGTATCTCATAGGAGTCCGCTCTCA  
AAATACACATTCCAGGCCATCGCATGGAGGAAGTCTGTGAGGGCTCCACTCGCACTCTGAAGCTCTGCAA  
CAGTCAGAAATGTCCCGGGACAGTGTGACTTCCGTGTGCTCAGTGTGCCGAGCACAACAGCAGACGATT  
CAGAGGGCGGCACTACAAGTGAAGCCTGATCAGGACTTATGCAAACTCTACTGTATCGCAGAAGGATTTGA  
TTTCTTCTTTCTTTGTCAAATAAAGTCAAAGATGGGACTCCATGCTCGGAGGATAGCCGTAATGTTGTAT  
AGATGGGATATGTGAGAGAGTTGGATGTGACAAATGTCCTTGGATCTGATGCTGTTGAAGACGCTCTGTGGGGT  
GTGTAACGGGAATAACTCAGCCTGCACGATTACAGGGGTCTCTACCTAGAGTATTATCATATGGTCACCAT  
TCCTTCTGGAGCCCGAGTATCCGCATCTATGAAATGAACGTCTCTACCTCTACATTCTGTGCGCAATGC  
CCTCAGAAGGTACTACCTGAATGGGCACTGGACCGTGGACTGGCCCGGCGGTACAAATTTTCGGGCACTAC

TTTCGACTACAGACGGTCTCTATAATGAGCCCCGAGAACTTAATCGCTACTGGACCAACCAACGAGACACTGAT  
TGTGGAGCTGCTGTTTCAGGGAAGGAACCCGGGTGTTGCCTGGGAATACTCCATGCCTCGCTTGGGGACCGA  
GAAGCAGCCCCCTGCCAGCCAGCTACACTTGGGCCATCGTGCGCTCTGAGTGCTCCGTGTCTGCGGAGG  
GGGTAGGTGCCCTCCAGTGCTGCTCCTGGAGGCAGCATGCAGCCTTTAGCCACTGCGTACATTGCACTGGC  
CTTTCCTGAATCCTAATGAGCAGCCCCGGGCTTCTCCCTGCCAGTAGCAGTGACATTCCCCAAGGTGGGGAGT  
GGTGGTCTGAGTGCTACTTGTCGGCCCGAGCTGCCTTCTCCAGTCTATCTGCTTCAGTGTGTACTCTGAG  
GAAGTCAGTAGATGCATTGCTT

In a search of public sequence databases, the NOV1c nucleic acid sequence, located on chromosome 5 has 646 of 1089 bases (59%) identical to a gb:GENBANK-

- 5 ID:AF140675|acc:AF140675.1 mRNA from *Homo sapiens* (*Homo sapiens* zinc metalloprotease ADAMTS7 (ADAMTS7) mRNA, complete cds) ( $E = 1.0e^{-23}$ ). Public nucleotide databases include all GenBank databases and the GeneSeq patent database.

The disclosed NOV1c polypeptide (SEQ ID NO:6) encoded by SEQ ID NO:5 has 856 amino acid residues and is presented in Table 1F using the one-letter amino acid code. Signal  
10 P, Psort and/or Hydropathy results predict that NOV1c has no signal peptide and is likely to be localized in the nucleus with a certainty of 0.7000. In other embodiments, NOV1c may also be localized to the microbody (peroxisome) with a certainty of 0.3813, the mitochondrial matrix space with a certainty of 0.1000, or in the lysosome (lumen) with a certainty of 0.1000.

**Table 1F. Encoded NOV1c protein sequence (SEQ ID NO:6).**

MGPAAPGSPSPVPRPPDLVSAYEVDHRGDYVSHEIMHHQRRRRRAVAVSEVEPAFLQVCRARELRLCVEAF  
PIANSQPGFLNLSNVRSHWREQHASKRIITNAMLGESALASTRKSNCVFFLSFYFFQSGMIRTEADYFLRP  
LPSHLSWKLGRAAQSSPSHVLYKREVLVTSRTWELAHQPLHSSDLRLGLPQKQHFGRKKYMPQPPKEDL  
FILPDEYKSLRHKRSLRLSHRNEELNVETLVVVDKMMQNHGHENITTYVLTILNMVSALFKDGTIGGNIN  
IAIVGLILLEDEQPLVISHHADHTLSSFCQWQSGLMGKDGRHDHAILLTGLDICSWKNEPCDTLGFAPIS  
GMCSKYRSCTINEDTGLGLAFTIAHESGHNFMIHDGEGNMCKKSEGNIMSPTLAGRNGVFSWSPCSRQYLH  
KFLSTAQAICLADQPKPVKEYKYPEKLPGLYANTQCKWQFGEKAKLCMLDFKKDICKALWCHIRIGRKCE  
KFMPAAEGTICGHEHGAGGQCVKYGDEGPKPTHGHSWSSWSPCSRTCGGGVSHRSRQNTSHRPSHGGKF  
CEGSTRTLKLCNSQKCPRDSVDFRAAQAEHNSRRFRGRHYKWKPDQDLCKLYCIAEGDFFFSLSNKVKDG  
TPCSEDSRNVCIDGICERVGCDNVLGSDAVEDVCGVCNGNNSACTIHRGLYLEYYHMTIPSGARSIRIYEM  
NVSTSYISVRNALRRYYLNGHWTVDWPGRYKFSGTTFDYRRSYNEPENLIATGPTNETLIVELLFQGRNPGV  
AWEYSMPRLGTEKQPPAQPSYTWAIVRSECSVSCGGRCPLVLLLEAACQPLATAYIALAFLES

15 A search of public sequence databases reveals that the NOV1c amino acid sequence has 256 of 579 amino acid residues (44%) identical to, and 352 of 579 amino acid residues (60%) similar to, the 997 amino acid residue ptrn:SWISSNEW-ACC:Q9UKP4 protein from *Homo sapiens* (Human) (ADAM-TS 7 precursor (EC 3.4.24.-) (A Disintegrin And  
20 Metalloproteinase With Thrombospondin Motifs 7) (ADAMTS-7) (ADAM-TS7) ( $E = 1.9e^{-144}$ ). Public amino acid databases include the GenBank databases, SwissProt, PDB and PIR.

NOV1c is expressed in at least the following tissues: adrenal gland, bone marrow, brain - amygdala, brain - cerebellum, brain - hippocampus, brain - substantia nigra, brain - thalamus, brain -whole, fetal brain, fetal kidney, fetal liver, fetal lung, heart, kidney,

lymphoma - Raji, mammary gland, pancreas, pituitary gland, placenta, prostate, salivary gland, skeletal muscle, small intestine, spinal cord, spleen, stomach, testis, thyroid, trachea and uterus. Expression information was derived from the tissue sources of the sequences that were included in the derivation of the sequence of NOV1c.

5

# NOV1d

A disclosed NOV1d nucleic acid of 2895 nucleotides (also referred to as CG52952-04) encoding a novel ADAM-TS 7-like protein is shown in Table 1G. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 10-12 and ending with a TAA codon at nucleotides 2866-2868. A putative untranslated region upstream from the initiation codon and downstream from the termination codon is underlined in Table 1G. The start and stop codons are in bold letters.

10

**Table 1G. NOV1d nucleotide sequence (SEQ ID NO:7).**

CGTCCTGGATGAAGCCCCGCGCGCGGATGGCGGGCTTGGCGGCGCTGTGGATGCTGTTGGCGCAGGTG  
GCGGAGCAGGTGAGTCCCGGGCGCTCCACCAGCGCGGAAACCGCGGGTCCGGACAGCTGGAGGCGAGTCCC  
CCGCGGCTCCTCTCCCGCGGACCCCGCGTCTCACCGCGATGTCGCGCTGTTTTCCGCAGGCACCTGCGTG  
CGCCATGGGACCCGAGCGGCGAGCGCTGGGAGCCCGAGCGTCCCGCGTCCCTCCACCCGCGGAGCGGGC  
GGGCTGGATGGAAAAGGGCGGGACATGGATGAAGCTGGAACCATCGTTCTCAGCAAATAACACAGGAACA  
GAAACCAAACTGCATGTTCTCACTCAATATGACCTGGTCTCTGCCACGAGGTTGACACAGGGGCGAT  
TACGTGTCCCATGAAATCATGCACCATCAGCGGCGGAGAAGAGCAGTGGCCGTGTCCGAGGTTGAGTCTCTT  
CACCTTCGGCTGAAAGGCCCGAGGCAGCACTTCCACATGGATCTGAGGACTTCCAGCAGCCTAGTGGCTCCT  
GGCTTTATTGTGCAGACGTTGGGAAAGACAGGCACTAAGTCTGTGCAGACTTACCGCCAGAGGACTTCTGT  
TTCTATCAAGGCTTTTGCATCACACAGAACTGCCATCGCATGGAGGGAAGTTCTGTGAGGCTCCACT  
CGCACTCTGAAGCTCTGCAACAGTCAGAAATGTCGCCGAGCAGTGTGACTTCCGTGCTGCTCAGTGTGCC  
GAGCACAACAGCAGACGATTACAGAGGCGGCACTACAAGTGAAGCCTTACACTCAAGTAGAAGCCGACTTA  
TGCAAACTCTACTGTATCGCAGAAGGATTTGATTTCTTCTTTCTTGTCAAATAAAGTCAAAGATGGGACT  
CCATGCTCGGAGGATAGCCGTAATGTTGTATAGATGGGATATGTGAGCTCAGTGTGGTGTCCACATCTGCG  
CACATGCCCCAGCCTCCCAAGGAAGACCTCTTCATCTTGCCAGATGAGTATAAGTCTTGCTTACGGCATAAG  
CGCTCTCTTCTGAGGTCCCATAGAAATGAAGAACTGAACGTGGAGACCTTGGTGGTGGTGCACAAAAGATG  
ATGCAAAACCATGGCCATGAAATATACCCACCTACGTGCTCACGATACTCAACATGGTATCTGCTTTATTC  
AAAGATGGATTGATGGGGAAGATGGGACTCGTCATGACCACGCCATCTTACTGACTGGTCTGGATATGT  
TCCTGGAAGAATGAGCCCTGTGACACTTTGGGATTTGCACCCATAAGTGGAATGTGTAGTAAATCTCGAGC  
TGCACGATTAATGAAGATACAGGTCTTGGACTGGCCTTACCATTGCCCATGAGTCTGGACACAACTTTGGC  
ATGATTATGATGGAGAAGGAACATGTGTAAGTCCGAGGGCAACATCATGTCCCTACATTGGCAGGA  
CGCAATGGAGTCTTCTCCTGGTCACCTGCAGCCGCCAGTATCTACACAAATTTCTAAGCACCCTCAAGCT  
ATCTGCCTTGCTGATCAGCCAAAGCCTGTGAAGGAATACAAGTATCCTGAGAAATTGCCAGGAGAATTATAT  
GATGCAAAACACAGTGCAAGTGGCAGTTCGGAGAGAAAGCCAAGCTCTGCATGCTGGACTTTAAAAAGGAC  
ATCTGTAAGCCCTGTGGTGCCATCGTATTGGAAGGAAATGTGAGACTAAATTTATGCCAGCAGCAGAAGGC  
ACAATTTGTGGGCATGACATGTGGTGCCGGGGAGGACAGTGTGTGAAATATGGTGATGAAGGCCCAAGCCC  
ACCCATGGCCACTGGTCCGACTGGTCTTCTTGGTCCCCATGCTCCAGGACCTGCGGAGGGGGAGTATCTCAT  
AGGAGTCGCCTCTGCACCAACCCCAAGCCATCGCATGGAGGGAAGTTCTGTGAGGGTCCACTCGCACTCTG  
AAGCTCTGCAACAGTCAGAAATGTCCCGGGACAGTGTGACTTCCGTGCTGCTCAGTGTGCCGAGCACAAC  
AGCAGACGATTACAGAGGCGGCACTACAAGTGAAGCCTTACACTCAAGTAGAAGATCAGGACTTATGCAAA  
CTCTACTGTATCGCAGAAGGATTTGATTTCTTCTTTCTTGTCAAATAAAGTCAAAGATGGGACTCCATGC  
TCGGAGGATAGCCGTAATGTTGTATAGATGGGATATGTGAGAGAGTTGGATGTGACAATGTCCTTGGATCT  
GATGCTGTGAAGACGTCTGTGGGTGTGTAAACGGGAATAACTCAGCCTGCAGATTACAGGGGTCTCTAC  
ACCAAGCACCAACCAACCAAGTATATCAGATGGTCACTTCCCTTCTGGAGCCCGGATCTCCGATC  
TATGAAATGAACGTCTCTACCTCTACATTTCTGTGCGCAATGCCCTCAGAAGGTAACCTGAATGGGCAC  
TGGACCGTGGACTGGCCCGGCGGTACAAATTTTCGGGCACTACTTTCGACTACAGACGGTCTTATAATGAG  
CCCGAGAACTTAATCGCTACTGGACCAACCAACGAGACACTGATTGTGGAGCTGCTGTTTCAGGGAAGGAAC  
CCGGGTGTTGCTGGGAATACTCCATGCCTCGTTGGGACCGAGAAGCAGCCCCCTGACGCCAGCTAC  
ACTTGGGCCATCGTGCTCTGAGTGTCTCGTGTCTGCGGAGGGGTAGGTGCCCTTCCAGTGTGCTCCTG  
GAGGCAGCATGTCAGCCTTACGCCACTGCGTACATTGCACTGGCCTTCTTGAATCCT**TAA**TGAGCAGCCCGG  
GGCTTCTCCCTGCCA

In a search of public sequence databases, the NOV1d nucleic acid sequence, located on chromosome 5 has 380 of 614 bases (61%) identical to a gb:GENBANK-ID:AF140675|acc:AF140675.1 mRNA from *Homo sapiens* (*Homo sapiens* zinc metalloprotease ADAMTS7 (ADAMTS7) mRNA, complete cds) ( $E = 7.0e^{-16}$ ). Public nucleotide databases include all GenBank databases and the GeneSeq patent database.

The disclosed NOV1d polypeptide (SEQ ID NO:8) encoded by SEQ ID NO:7 has 952 amino acid residues and is presented in Table 1H using the one-letter amino acid code. Signal P, Psort and/or Hydropathy results predict that NOV1d has a signal peptide and is likely to be localized extracellularly with a certainty of 0.3700. In other embodiments, NOV1d may also be localized to the lysosome (lumen) with a certainty of 0.1900, the microbody (peroxisome) with a certainty of 0.1270, or in the endoplasmic reticulum (membrane) with a certainty of 0.1000. The most likely cleavage site for a NOV1a peptide is between amino acids 23 and 24, at: VAE-QV.

**Table 1H. Encoded NOV1d protein sequence (SEQ ID NO:8).**

<p>MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQGRNGSGGLEASPPRLLSRGPRRLTAMSPLFSAGTCVRHG  TRSGSAWEPPERPASSSTRGAAGLDGKGRMDDEAGNHRSSQQTNTGTENQTLHVLTQYDLVSAYEVDHRGDYVS  HEIMHHQRRRAVAVSEVESLHLRLKGRPHDFHMDLRTSSSLVAPGFIVQTLGKTGTSKVQTLPPEDFCFYQ  GSLRSHRNSPSHGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKWKPYTQVEADLCCL  YCIAEGFDFFFSLSNKVKDGTPCSEDSRNVCIDGICELSVVSTSAHMPQPPKEDLFILPDEYKSLRHKRSL  LRSHRNEELNVETLVVVDKMMQNHGHEINITTYVLTILNMVSALFKDGLMGKDGTGRHDHAILLTGLDICSWK  NEPCDTLGFAPISGMCSKYRSCTINEDTGLGLAFTIAHESGHNFNGMIHDGEGNMCKKSEGNIMSPTLAGRNG  VFSWSPCSRQYLHKFLSTAQAICLADQPKPVKEYKYPEKLPGEYDANTQCKWQFGEKAKLCMLDFKKDICK  ALWCHRIGRKCEKFMFAAEGTICGHDMWCRGGQCVKYGDEGPKPTHGHWSWSSWSPCSRTCGGGVSHRSR  LCTNPKPSHGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKWKPYTQVEDQLCKLYC  IAEGFDFFFSLSNKVKDGTPCSEDSRNVCIDGICERVGCNVLGSDAVEDVCGVCNGNSACTIHRGLYTKH  HHTNQYYHMTIPSGARSIRIYEMNVSTSYISVRNALRRYYLNGHWTVDPGRYKFSGTTFDYRRSYNEPEN  LIATGPTNETLIVELLFQGRNPGVAWEYSMPRLGTEKQPPAQPSYTWIVRSECSVSCGGGRCLPVLLLEAA  CQPSATAYIALAFLES</p>
---

A search of public sequence databases reveals that the NOV1d amino acid sequence has 207 of 483 amino acid residues (42%) identical to, and 287 of 483 amino acid residues (59%) similar to, the 997 amino acid residue ptnr:SWISSNEW-ACC:Q9UKP4 protein from *Homo sapiens* (Human) (ADAM-TS 7 precursor (EC 3.4.24.-) (A Disintegrin And Metalloproteinase With Thrombospondin Motifs 7) (ADAMTS-7) (ADAM-TS7)) ( $E = 7.0e^{-16}$ ). Public amino acid databases include the GenBank databases, SwissProt, PDB and PIR.

NOV1d is expressed in at least the following tissues: adrenal gland, bone marrow, brain - amygdala, brain - cerebellum, brain - hippocampus, brain - substantia nigra, brain - thalamus, brain -whole, fetal brain, fetal kidney, fetal liver, fetal lung, heart, kidney, lymphoma - Raji, mammary gland, pancreas, pituitary gland, placenta, prostate, salivary gland, skeletal muscle, small intestine, spinal cord, spleen, stomach, testis, thyroid, trachea and



uterus. Expression information was derived from the tissue sources of the sequences that were included in the derivation of the sequence of NOV1d and the expression pattern of (GENBANK-ID: gb:GENBANK-ID:AF140675|acc:AF140675.1) a closely related Homo sapiens zinc metalloprotease ADAMTS7 (ADAMTS7) mRNA, complete cds homolog in species Homo sapiens.

The proteins encoded by the NOV1a, 1b, 1c, and 1d nucleotides are very closely homologous as is shown in the alignment in Table II. TaqMan data for NOV1 can be found below in Example 2. SNP data for NOV1a can be found below in Example 3.

**Table II Alignment of NOV1a, 1b, 1c, and 1d.**

10		10	20	30	40	50
	NOV1a	MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQRGNGRSGQLEASPPRLLS				
	NOV1b	MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQRGNGRSGQLEASPPRLLS				
	NOV1d	MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQRGNGRSGQLEASPPRLLS				
15	NOV1c	-----MGPAAAI-----				
		60	70	80	90	100
	NOV1a	RGPRRLTAMSPLFSAGTCVRHGTRSGSAWEPPERPASSSTRGAAGLDGKGF				
20	NOV1b	RGPRRLTAMSPLFSAGTCVRHGTRSGSAWEPPERPASSSTRGAAGLDGKGF				
	NOV1d	RGPRRLTAMSPLFSAGTCVRHGTRSGSAWEPPERPASSSTRGAAGLDGKGF				
	NOV1c	-----GSPSVRPP-----				
		110	120	130	140	150
25	NOV1a	DMDEAGNHRSQQTNTGTENQTLHVLT-CYDLVSAYEVDHRGDYVSHEIMH				
	NOV1b	DMDEAGNHRSQQTNTGTENQTLHVLTREYDLVSAYEVDHRGDYVSHEIMH				
	NOV1d	DMDEAGNHRSQQTNTGTENQTLHVLT-CYDLVSAYEVDHRGDYVSHEIMH				
	NOV1c	-----DLVSAYEVDHRGDYVSHEIMH				
30		160	170	180	190	200
	NOV1a	HQRRRAVAVSEVESLHLRLKGRHDFHMDLRTSSSLVAPGFIVQTLGKT				
	NOV1b	HQRRRAVAVSEVESLHLRLKGRHDFHMDLRTSSSLVAPGFIVQTLGKT				
35	NOV1d	HQRRRAVAVSEVESLHLRLKGRHDFHMDLRTSSSLVAPGFIVQTLGKT				
	NOV1c	HQRRRAVAVSEVEPAFLQYCRAPELRLCVEAFPIANSQPGHINLENVRS				
		210	220	230	240	250
40	NOV1a	GTKSVQTLI--PEDFCFYQGSLSHRNSPSHGKFCGSTRTLKLCNSQF				
	NOV1b	GTKSVQTLI--PEDFCFYQGSLSHRNSPSHGKFCGSTRTLKLCNSQF				
	NOV1d	GTKSVQTLI--PEDFCFYQGSLSHRNSPSHGKFCGSTRTLKLCNSQF				
	NOV1c	HWREQHASKRIITNAMLGSSASTKES---NCVFLS----FYFFCSGM				
45		260	270	280	290	300
	NOV1a	CPRDSVDFRAAQCAEHNSRRFRGRHYKWKPYTQVEQDLCKLYCIAEGFDF				
	NOV1b	CPRDSVDFRAAQCAEHNSRRFRGRHYKWKPYTQVEQDLCKLYCIAEGFDF				
	NOV1d	CPRDSVDFRAAQCAEHNSRRFRGRHYKWKPYTQVEADLCKLYCIAEGFDF				
50	NOV1c	IRTEAEALFLRPLPSHLSWKLGRAAQGSSESHVLYKREVLVTSRTWELAH				
		310	320	330	340	350
	NOV1a	FFSLSNKVVVDGTPCSEDSRNVCIDGICELSVVSTSAHMFQPPKEDLFILE				
55	NOV1b	FFSLSNKVVVDGTPCSEDSRNVCIDGICE-----MFQPPKEDLFILE				
	NOV1d	FFSLSNKVVVDGTPCSEDSRNVCIDGICELSVVSTSAHMFQPPKEDLFILE				
	NOV1c	QPLHSDDLRLQLKQHFCCRKK-----MFQPPKEDLFILE				
60		360	370	380	390	400

NOV1a DEYFSCLEHHFSLRSHRNEELNVETLVVVDFKMMQNHGHENITTYVLT  
 NOV1b DEYFSCLEHHFSLRSHRNEELNVETLVVVDFKMMQNHGHENITTYVLT  
 NOV1d DEYFSCLEHHFSLRSHRNEELNVETLVVVDFKMMQNHGHENITTYVLT  
 NOV1c DEYFSCLEHHFSLRSHRNEELNVETLVVVDFKMMQNHGHENITTYVLT

5

410 420 430 440 450  
 NOV1a LNMVSALFEDGTICGNINIAIVGLILLEDEQPLVISHHADHTLSSFCQW  
 NOV1b LNMVSALFEDGTIC-----  
 10 NOV1d LNMVSALFEDGLEYK-----  
 NOV1c LNMVSALFEDGTICGNINIAIVGLILLEDEQPLVISHHADHTLSSFCQW

460 470 480 490 500  
 15 NOV1a QSGLMGKDGTRHDHAILLTGLDICSWFNEPCDTLGFAPISGMCSKYRSC  
 NOV1b -----  
 NOV1d -----DGTRHDHAILLTGLDICSWFNEPCDTLGFAPISGMCSKYRSC  
 NOV1c QSGLMGKDGTRHDHAILLTGLDICSWFNEPCDTLGFAPISGMCSKYRSC

20

510 520 530 540 550  
 NOV1a INEDTGLGLAFTIAHESGHNFPMIHDGEGNMCKKSEGNIMSPTLAGRNGV  
 NOV1b -----GNIN-----  
 NOV1d INEDTGLGLAFTIAHESGHNFPMIHDGEGNMCKKSEGNIMSPTLAGRNGV  
 25 NOV1c INEDTGLGLAFTIAHESGHNFPMIHDGEGNMCKKSEGNIMSPTLAGRNGV

560 570 580 590 600  
 30 NOV1a FSWSPCSRQYLHKFLSTAQAICLADQPKPKVKEYKYPEKLPGLYDANTQC  
 NOV1b -----ALVG-----  
 NOV1d FSWSPCSRQYLHKFLSTAQAICLADQPKPKVKEYKYPEKLPGLYDANTQC  
 NOV1c FSWSPCSRQYLHKFLSTAQAICLADQPKPKVKEYKYPEKLPGLYDANTQC

610 620 630 640 650  
 35 NOV1a KWQFGEKAKLCMLDFKKDICKALWCHRIGRKCEKFMPPAAEGTICGHDMW  
 NOV1b -----LLEKEDICKALWCHRIGRKCEKFMPPAAEGTICGHDMW  
 NOV1d KWQFGEKAKLCMLDFKKDICKALWCHRIGRKCEKFMPPAAEGTICGHDMW  
 NOV1c KWQFGEKAKLCMLDFKKDICKALWCHRIGRKCEKFMPPAAEGTICGHDMW

40

660 670 680 690 700  
 45 NOV1a CRGGQCVKYGDEGPKPTHGHWSWSSWSPCSRTCGGGVSHRSF-LCTNEN  
 NOV1b CRGGQCVKYGDEGPKPTHGHWSWSSWSPCSRTCGGGVSHRSF-LCTNEN  
 NOV1d CRGGQCVKYGDEGPKPTHGHWSWSSWSPCSRTCGGGVSHRSF-LCTNEN  
 NOV1c A-GGQCVKYGDEGPKPTHGHWSWSSWSPCSRTCGGGVSHRSFSQNTLSF

710 720 730 740 750  
 50 NOV1a PSHGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKW  
 NOV1b PSHGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKW  
 NOV1d PSHGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKW  
 NOV1c PSHGGKFCEGSTRTLKLCNSQKCPRDSVDFRAAQCAEHNSRRFRGRHYKW

55

760 770 780 790 800  
 NOV1a KF-----QDLCKLYCIAEGFDFFFSLSNKVVDGTPCEDSRNVCIDGIC  
 NOV1b KF-----QDLCKLYCIAEGFDFFFSLSNKVVDGTPCEDSRNVCIDGIC  
 NOV1d KFYTQVEDQDLCKLYCIAEGFDFFFSLSNKVVDGTPCEDSRNVCIDGIC  
 60 NOV1c KF-----DQDLCKLYCIAEGFDFFFSLSNKVVDGTPCEDSRNVCIDGIC

810 820 830 840 850  
 65 NOV1a EX-GCDNVLGSDAVEDVCGVCNGNNSACTIHRGLYTKHHHTN-VYHMTI  
 NOV1b E--GCDNVLGSDAVEDVCGVCNGNNSACTIHRGLYTKHHHTN-VYHMTI  
 NOV1d ERVGCNDVLGSDAVEDVCGVCNGNNSACTIHRGLYTKHHHTN-VYHMTI  
 NOV1c ERVGCNDVLGSDAVEDVCGVCNGNNSACTIHRGLYTKHHHTN-VYHMTI

70

860 870 880 890 900

NOV1a PSQARSIRIYEENYSTSYISVFNALRRYYLNGHWTVDWPGRYKFSGTTFI  
 NOV1b PSQARSIRIYEENYSTSYISVFNALRRYYLNGHWTVDWPGRYKFSGTTFI  
 NOV1d PSQARSIRIYEENYSTSYISVFNALRRYYLNGHWTVDWPGRYKFSGTTFI  
 NOV1c PSQARSIRIYEENYSTSYISVFNALRRYYLNGHWTVDWPGRYKFSGTTFI

5

910 920 930 940 950  
 NOV1a YRRSYNEPENLIATGPTNETLIVELLFQGRNPGVAWEYSMPRLGTEKQPI  
 NOV1b YRRSYNEPENLIATGPTNETLIVELLFQGRNPGVAWEYSMPRLGTEKQPI  
 NOV1d YRRSYNEPENLIATGPTNETLIVELLFQGRNPGVAWEYSMPRLGTEKQPI  
 NOV1c YRRSYNEPENLIATGPTNETLIVELLFQGRNPGVAWEYSMPRLGTEKQPI

10

960 970 980 990  
 NOV1a AQPSTWAIVRSECSVSCGGGRCLPVLLEAACQPSATAYIALAFLES (SEQ ID NO:2)  
 NOV1b AQPSTWAIVRSECSVSCGGGRCLPVLLEAACQPSATAYIALAFLES (SEQ ID NO:4)  
 NOV1d AQPSTWAIVRSECSVSCGGGRCLPVLLEAACQPSATAYIALAFLES (SEQ ID NO:6)  
 NOV1c AQPSTWAIVRSECSVSCGGGRCLPVLLEAACQPSATAYIALAFLES (SEQ ID NO:8)

15

20

Homologies to any of the above NOV1 proteins will be shared by the other two NOV1 proteins insofar as they are homologous to each other as shown above. Any reference to NOV1 is assumed to refer to all three of the NOV1 proteins in general, unless otherwise noted.

The disclosed NOV1a polypeptide has homology to the amino acid sequences shown in the BLASTP data listed in Table 1J.

25

Table 1J. BLAST results for NOV1a					
Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 13569928 ref NP_112217.1	a disintegrin-like and metalloprotease with thrombospondin type 1 motif, 12 [Homo sapiens]	1593	269/616 (43%)	371/616 (59%)	e-135
gi 11493589 gb AAG35563.1 AF163762_1 (AF163762)	zinc metalloendopeptidase [Homo sapiens]	1077	253/624 (40%)	359/624 (56%)	e-128
gi 10645199 ref NP_055087.1	a disintegrin and metalloprotease with thrombospondin motifs-7 preproprotein; a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 7 [Homo sapiens]	997	247/571 (43%)	344/571 (59%)	e-127

gi 15309931 ref XP_054419.1	a disintegrin-like and metalloprotease domain with thrombospondin type I repeats 10 [Homo sapiens]	854	245/604 (40%)	348/604 (57%)	e-124
gi 7656869 ref NP_055088.1	a disintegrin and metalloprotease with thrombospondin motifs-6 preproprotein; a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 6 [Homo sapiens]	860	244/630 (38%)	345/630 (54%)	e-115

The homology between these and other sequences is shown graphically in the ClustalW analysis shown in Table 1K. In the ClustalW alignment of the NOV1 proteins, as well as all other ClustalW analyses herein, the black outlined amino acid residues indicate regions of conserved sequence (*i.e.*, regions that may be required to preserve structural or functional properties), whereas non-highlighted amino acid residues are less conserved and can potentially be altered to a much broader extent without altering protein structure or function.

**Table 1K. ClustalW Analysis of NOV1**

1)	Novel NOV1a	(SEQ ID NO:2)
2)	Novel NOV1b	(SEQ ID NO:4)
3)	Novel NOV1c	(SEQ ID NO:6)
4)	Novel NOV1d	(SEQ ID NO:8)
4)	gi 13569928 ref NP_112217.1	a disintegrin-like and metalloprotease with thrombospondin type 1 motif, 12 [Homo sapiens] (SEQ ID NO:30)
5)	gi 11493589 gb AAG35563.1	AF163762_1 (AF163762) zinc metalloendopeptidase [Homo sapiens] (SEQ ID NO:31)
6)	gi 10645199 ref NP_055087.1	a disintegrin and metalloprotease with thrombospondin motifs-7 preproprotein; a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 7 [Homo sapiens] (SEQ ID NO:32)
7)	gi 15309931 ref XP_054419.1	a disintegrin-like and metalloprotease domain with thrombospondin type I repeats 10 [Homo sapiens] (SEQ ID NO:33)
8)	gi 7656869 ref NP_055088.1	a disintegrin and metalloprotease with thrombospondin motifs-6 preproprotein; a disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 6 [Homo sapiens] (SEQ ID NO:34)
<div> <div>10</div> <div>20</div> <div>30</div> <div>40</div> </div>		
NOV1a	MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQRGNRGSGQ	
NOV1b	MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQRGNRGSGQ	
NOV1c	-----	
NOV1d	MKPRARGWRGLAALWMLLAQVAEQVSPGRSHQRGNRGSGQ	
gi 13569928	-----	
gi 11493589	-----	
gi 10645199	-----	

```

gi|15309931| -----
gi|7656869| -----

          50      60      70      80
5  NOV1a    LEASPPRLLSRGPRRLTAMSPLFSAAGTCVRHGTRSGSAWE
   NOV1b    LEASPPRLLSRGPRRLTAMSPLFSAAGTCVRHGTRSGSAWE
   NOV1c    -----MGP-----AAAAP-----GSPSV
   NOV1d    LEASPPRLLSRGPRRLTAMSPLFSAAGTCVRHGTRSGSAWE
10 gi|13569928| ---MPCAQRSWLANLSVVAQLLNFGALCYGRQPQGPVRF
   gi|11493589| -----
   gi|10645199| ---MPGGPSRSPAPLLRPLLLLCALAPG-APGPAPGRA
   gi|15309931| -----
   gi|7656869| -----MHILWKTLTWILSLIMASSEFHS DHRL

          90      100     110     120
20 NOV1a    PERPASSSTRGAAGLDGKGRDMDEAGNHRSSQQTNTGTENQ
   NOV1b    PERPASSSTRGAAGLDGKGRDMDEAGNHRSSQQTNTGTENQ
   NOV1c    PRPPP-----
   NOV1d    PERPASSSTRGAAGLDGKGRDMDEAGNHRSSQQTNTGTENQ
   gi|13569928| PDRRQEHFIKGL-----
   gi|11493589| --RSQDEFLSSL-----
   gi|10645199| TEGR-----
25 gi|15309931| -----
   gi|7656869| SYSSQEEFLTYL-----

          130     140     150     160
30 NOV1a    TLHVLT-QYOLVSAYEVDHRCQYVSHEIMHHQR-RRRAVA
   NOV1b    TLHVLTREYOLVSAYEVDHRCQYVSHEIMHHQR-RRRAVA
   NOV1c    -----DIYSAYEVDHRCQYVSHEIMHHQR-RRRAVA
   NOV1d    TLHVLT-QYOLVSAYEVDHRCQYVSHEIMHHQR-RRRAVA
   gi|13569928| -----PEMHVVGPRVVDASCHTSLSYGLHYPTSSRRKRD
35 gi|11493589| -----ESMIAFPTRVDHNCALLAFSPPPP-RRQRRTG
   gi|10645199| -----AALDIIVHPVRVDAGCSFLSYELWP--R-ALRKRD
   gi|15309931| -----
   gi|7656869| -----EHYQLTIPIRVDQNGAFLSPTVKN-D-HSRRRRS

          170     180     190     200
40 NOV1a    VS-----EVES-LHLRLKGRPHDFHMDTRTSSSLVAPGFI
   NOV1b    VS-----EVES-LHLRLKGRPHDFHMDTRTSSSLVAPGFI
   NOV1c    VS-----EVES-AFLVCRARELRCLVEAFPIANSQPGFI
45 NOV1d    VS-----EVES-LHLRLKGRPHDFHMDTRTSSSLVAPGFI
   gi|13569928| LD-----GSELDW-VYVYISHEEKDLFFITVNOGFLSNSTI
   gi|11493589| AT-----AERRLPYVVASPSTHFLLRTRSSRLLAGHVS
   gi|10645199| VS-----VRRDAPAFYELQYRGRELRFMTANQHLLAPGFV
   gi|15309931| -----
50 gi|7656869| MDPIDPQQAVSKLFFKLSAYGKHFLHNTLTMTDFVSKHET

          210     220     230     240
55 NOV1a    VQTLGKTCTKSVQTLF--P-EDFCFYQCSLRSHRNSPSHG
   NOV1b    VQTLGKTCTKSVQTLF--P-EDFCFYQCSLRSHRNSPSHG
   NOV1c    NLSNVRSWREQHASKRII-TNAMLGESALASTRKS---N
   NOV1d    VQTLGKTCTKSVQTLF--P-EDFCFYQCSLRSHRNSPSHG
   gi|13569928| MEKR-YGNLSHVMMAS--SAPLCHLSCTVLQQGTRVGTA
   gi|11493589| VYWTREELAWQRAARP-----HCLYACHLQGGASS-HV
60 gi|10645199| SETRRRGCLGRAHIRAH---TPACHLLCEVDPELEGGLA
   gi|15309931| -----
   gi|7656869| VYTWGKDEPQWKHDFLD-----NCHYTCYLDQQRSTT-KV

          250     260     270     280
65 NOV1a    GKFCGSTRTLKLCNSQKCPRDSVDTRAAQCAEHNSRRFR
   NOV1b    GKFCGSTRTLKLCNSQKCPRDSVDTRAAQCAEHNSRRFR
   NOV1c    CVFFL-S---FYFFQSGMIRTHEADYFLRPLPSHLSWKLG
   NOV1d    GKFCGSTRTLKLCNSQKCPRDSVDTRAAQCAEHNSRRFR
70 gi|13569928| ALSACHG-----LTGFFQLPHGDFIEPVKKHPLVEG-

```

```

gi|11493589| AISTCGG-----LHGLIVADEEYLLIEPLHGGPKGS--
gi|10645199| AISACDG-----LKGVFQLSNEDYFIEPLDSAPARPG-
gi|15309931| -----
gi|7656869| ALSNCVG-----LHGVIATEDERYFIEPLKNTTDSKH

5
290 300 310 320
NOV1a GRHYKWKPTQVLDLCKLYCIAEGDFFFSLENKVKDGT
NOV1b GRHYKWKPTQVLDLCKLYCIAEGDFFFSLENKVKDGT
10 NOV1c RAAQGSSPSHVLYKREVLVTSRTWELAHQPLHSDLELGL
NOV1d GRHYKWKPTQVLDLCKLYCIAEGDFFFSLENKVKDGT
gi|13569928| ----GYHPEIVYRRQKVP-----ETKEP-TCGLKDSV
gi|11493589| RSPESGPEVYVYRSSLRHPHLDTACGV-RDEKPKWGRPW
gi|10645199| ----HAQPEVYVYRQAPER--LAQRGDSSAPSTCGVQVYP
15 gi|15309931| -----
gi|7656869| FSYENGHEVYVYKKSALQQRHLYDHSCHGVSDPTRSGKPW

330 340 350 360
20 NOV1a PCSEDSRNVCIDGICELSVVSTSAHMPQPPKEDLFILPDE
NOV1b PCSEDSRNVCIDGICE-----MPQPPKEDLFILPDE
NOV1c PQKQHFCGRKK-----YMPQPPKEDLFILPDE
NOV1d PCSEDSRNVCIDGICELSVVSTSAHMPQPPKEDLFILPDE
25 gi|13569928| NISQ-----KQELWREK
gi|11493589| WLRT-----LKPPPAPR
gi|10645199| ELES-----RRERWEQR
gi|15309931| -----
gi|7656869| WLN-----DTSTVSYS

370 380 390 400
30 NOV1a YKSLRHKRSLLSHRMELNVETLVVVDKMMQNHGHEN
NOV1b YKSLRHKRSLLSHRMELNVETLVVVDKMMQNHGHEN
NOV1c YKSLRHKRSLLSHRMELNVETLVVVDKMMQNHGHEN
35 NOV1d YKSLRHKRSLLSHRMELNVETLVVVDKMMQNHGHEN
gi|13569928| WERHNLPSRSLSRSSISERWVETLVVVDKMMQNHGHEN
gi|11493589| LGNETERGQPLRSVSRERYVETLVVVDKMMQNHGHEN
gi|10645199| QQWRRPRLRLPHQRSVSEKQVETLVVVDKMMQNHGHEN
gi|15309931| -----MMVAYHGRRD
40 gi|7656869| LPINNTTHIHRQKRSVSIERFVETLVVVDKMMQNHGHEN

410 420 430 440
45 NOV1a ITTYVLTINMVSALFDGTCGG-----NINIAIVGLLIL
NOV1b ITTYVLTINMVSALFDGTCGG-----NINIAIVGLLIL
NOV1c ITTYVLTINMVSALFDGTCGG-----NINIAIVGLLIL
NOV1d ITTYVLTINMVSALFDGLCKDGTTRHDHAILLTGLDTC
50 gi|13569928| VESYVLTINMVTGLPHPHICN-----ATIEIVVRLIL
gi|11493589| VEQYVLAI NNVAKLFQDSSIGS-----TVNIIIVRLIL
gi|10645199| VESYVLTINMVSALFDGTCGG-----NINIAIVGLLIL
gi|15309931| VEQYVLAI NNVAKLFQDSSIGS-----TVNIIIVRLIL
gi|7656869| IEHYILAV NNVAKLFQDSSIGS-----VNNIIIVARLIL

450 460 470 480
55 NOV1a EDLQPGVISHHADHTLSSFCWQSGLMGKDG-----
NOV1b EDLQ-----
NOV1c EDLQPGVISHHADHTLSSFCWQSGLMGKDG-----
NOV1d SWLN-----
60 gi|13569928| EELQGLKIVHHAETLSSFCWQKSINPKS-----DL
gi|11493589| TEQPTLEITHHAGKLDSSFCWQKSIVNHSGHGNAIPEN
gi|10645199| EDLQEDLITHHADNTLKSFCWQKSINMKG-----DA
gi|15309931| TEQPTLEITHHAGKLDSSFCWQKSIVNHSGHGNAIPEN
gi|7656869| TEQPNLEINHHADKLDSSFCWQKSILSHQSDGNTIPEN

490 500 510 520
70 NOV1a --TRHDHAILLTGLDICSWN-PCDTLGFAPISGMCSKYR
NOV1b --TRHDHAILLTGLDICSWN-PCDTLGFAPISGMCSKYR
NOV1c --TRHDHAILLTGLDICSWN-PCDTLGFAPISGMCSKYR

```

NOV1d  
gi|13569928| NPVHHDVAILLIRKIDICAGFNPCETLGLSHLFGMCQPHR  
gi|11493589| GVANHDVAILLIRYDICIYKNPCGTLGLAPVGMCCERER  
5 gi|10645199| HPLHHDVAILLIRKIDICAGFNPCETLGLSHVFGMCQPHR  
gi|15309931| GVANHDVAILLIRYDICIYKNPCGTLGLAPVGMCCERER  
gi|7656869| GIAHHDVAILLIRYDICIYKNPCGTLGLASVAGMCEPER

530 540 550 560  
NOV1a SCINEDTGLGLAFTIAHESGHFGMIHDGEGNCKKSEG  
NOV1b SCINEDTGLGLAFTIAHESGHFGMIHDGEGNCKKSEG  
NOV1c SCINEDTGLGLAFTIAHESGHFGMIHDGEGNCKKSEG  
NOV1d SCINEDTGLGLAFTIAHESGHFGMIHDGEGNCKKSEG  
10 gi|13569928| SCINEDSGIPLAFTIAHEIGHFGIHDGKENDCEPVGR  
gi|11493589| SCINEDTGLPLAFTIAHEIGHFGIHDGSGNCEPVGR  
15 gi|10645199| SCINEDTGLPLAFTIAHEIGHFGIHDGSGNCEPVGR  
gi|15309931| SCINEDTGLATAFTIAHEIGHFGMIHDGSGNCEPVGR  
gi|7656869| SCINEDTGLSAFTIAHEIVHFGMIHDGSGNCEPVGR

570 580 590 600  
NOV1a N---TMSPTLAGRNGVTSWSPCSRQYGHKFLSTAQAICLA  
NOV1b N---TMSPTLAGRNGVTSWSPCSRQYGHKFLSTAQAICLA  
NOV1c N---TMSPTLAGRNGVTSWSPCSRQYGHKFLSTAQAICLA  
25 NOV1d N---TMSPTLAGRNGVTSWSPCSRQYGHKFLSTAQAICLA  
gi|13569928| HP-YIMRQLQYDPTPLTWSKCSERYTRFLDRGWGFCLD  
gi|11493589| DPAKIMAAHTMTKTNPVWSSCSRQYTSFLDSGLGLCLN  
gi|10645199| RP-FIMSPQLLYDAAPLTWSRCSRQYTRFLDRGWGFCLD  
gi|15309931| DPAKIMAAHTMTKTNPVWSSCSRQYTSFLDSGLGLCLN  
30 gi|7656869| VMLQONTGSSHYCEYQSFELVCLQ

610 620 630 640  
NOV1a DQPKPVKEYKPEKLPGEYDANTOCWOFGEKAKLCMLD  
NOV1b DQPKPVKEYKPEKLPGEYDANTOCWOFGEKAKLCMLD  
NOV1c DQPKPVKEYKPEKLPGEYDANTOCWOFGEKAKLCMLD  
35 NOV1d DQPKPVKEYKPEKLPGEYDANTOCWOFGEKAKLCMLD  
gi|13569928| DIPKK-KGLKSKVIAPGVYVDVHQCQLOYG--PNATFCQ  
gi|11493589| MRPPR-QDFVYPTVAPQAYDADEQCQFCH--GVKSRQC  
40 gi|10645199| DPAK-DIIDPSPVPGVLYDVSHQCQLOYG--AYSAPCE  
gi|15309931| MRPPR-QDFVYPTVAPQAYDADEQCQFCH--GVKSRQC  
gi|7656869| SRX-----HHQ

650 660 670 680  
NOV1a FKKDLCALWCHRIGRCETKFPAAEGTTCG----HDMW  
NOV1b ---DKALWCHRIGRCETKFPAAEGTTCG----HDMW  
NOV1c FKKDLCALWCHRIGRCETKFPAAEGTTCG----HEHG  
45 NOV1d FKKDLCALWCHRIGRCETKFPAAEGTTCG----HDMW  
gi|13569928| EVENVCOTLWCSVKG-FCRSLDAAAQTCQCG----EKKW  
50 gi|11493589| KYGLVCSLWCLSKSNFCITNSIPAAEGTTCQTHTIDKGW  
gi|10645199| DMDNVCHTLWCSVGT-TCHSLDAAVIGTRCG----ENKW  
gi|15309931| KYGLVCSLWCLSKSNFCITNSIPAAEGTTCQTHTIDKGW  
gi|7656869| LFRVVCRLWCLSKSNFCVTNSIPAAEGTTCQTNIEKGW

690 700 710 720  
NOV1a CRGGCVKICDEGPPTHGHSWSSWSPCSRTCGGGVSH  
NOV1b CRGGCVKICDEGPPTHGHSWSSWSPCSRTCGGGVSH  
NOV1c A-CGCVKICDEGPPTHGHSWSSWSPCSRTCGGGVSH  
60 NOV1d CRGGCVKICDEGPPTHGHSWSSWSPCSRTCGGGVSH  
gi|13569928| CMAGCTIVCKK-P-SIPCGWRWSPWHSRSCGAGVQS  
gi|11493589| CYKRVCPVCSR-P-GVDGAWGPWPWGDCSRTCGGGVSS  
gi|10645199| CLSGECVPVCFR-P-PAVDGGSWGSWHSRSCGAGVQS  
65 gi|15309931| CYKRVCPVCSR-P-GVDGAWGPWPWGDCSRTCGGGVSS  
gi|7656869| CVOGDCVPVCTW-P-SIDGGWGPWSLWGECSRTCGGGVSS

730 740 750 760  
NOV1a RSR-LCTHNPESHGGKCEGSTRTLLKCNQKCPRDSVDF

	NOV1b	RSR-LCTPRPFGGKCEGSTRTLCLCNQKCPRDSVDF
	NOV1c	RSRSQNTHSRPFHGKCEGSTRTLCLCNQKCPRDSVDF
	NOV1d	RSR-LCTPRPFGGKCEGSTRTLCLCNQKCPRDSVDF
5	gi 13569928	AER-LCNLEPKFEGGKCTGERIRYRLCNVHPCRSEPTF
	gi 11493589	SSR-HCDPRPTIGGKCLGERRRHSCNTDDCPGGSODF
	gi 10645199	AER-QCTQPTPKYKGLCVGERIRFRCLNLQACBAGRPSF
	gi 15309931	SSR-HCDPRPTIGGKCLGERRRHSCNTDDCPGGSODF
	gi 7656869	SIR-HCDPRAPGGGKCLGERIRYRSCNTDPCPLGSRDF
10		770 780 790 800
	NOV1a	RAAQC EHI SRRFRG HYKWKP-----QDLC LMC AEG
	NOV1b	RAAQC EHI SRRFRG HYKWKP-----QDLC LMC AEG
	NOV1c	RAAQC EHI SRRFRG HYKWKP-----QDLC LMC AEG
15	NOV1d	RAAQC EHI SRRFRG HYKWKPYTQVEDQDLC LMC AEG
	gi 13569928	ROMQC EFT VPM NELYHWEIFN--PAHPC LMC CRPID
	gi 11493589	REVQC EFL SIPFRG FYKWKTYRGG-GVKACSLTSAEG
	gi 10645199	RHVQC HFD MLMLGLTWVEVVN--DVNPC LMC CRPAN
	gi 15309931	REVQC EFL SIPFRG FYKWKTYRGG-GVKACSLTSAEG
20	gi 7656869	REKQC AFD MPFRG IYNNWKPYTGG-GVKPCALNCAEG
		810 820 830 840
25	NOV1a	FDF FSLSKVKDGTGPS--EDSRVCI GIC X-GCDNV
	NOV1b	FDF FSLSKVKDGTGPS--EDSRVCI GIC --GCDNV
	NOV1c	FDF FSLSKVKDGTGPS--EDSRVCI GIC RVGCDNV
	NOV1d	FDF FSLSKVKDGTGPS--EDSRVCI GIC RVGCDNV
	gi 13569928	GQSEKMLDAVIDGTGCFEGGSRVCI GIC MVGCDYE
	gi 11493589	FDFYTERAAAVVDGTPCR--PDVVDICVSGEC HVGCDRV
30	gi 10645199	EYSAKKLRLAAVVDGTPCYQVRASRQCI GIC NVGCDYE
	gi 15309931	FDFYTERAAAVVDGTPCR--PDVVDICVSGEC HVGCDRV
	gi 7656869	YDFYTERAAVVDGTQCN--ADSLDICI GEC HVGCDNV
		850 860 870 880
35	NOV1a	LGSDAVEDVCGVCNGINSACTIHRGLTKHHHTN-HYHLV
	NOV1b	LGSDAVEDVCGVCNGINSACTIHRGLTKHHHTN-HYHLV
	NOV1c	LGSDAVEDVCGVCNGINSACTIHRGL--LE--HYHLV
	NOV1d	LGSDAVEDVCGVCNGINSACTIHRGLTKHHHTNQHYHLV
40	gi 13569928	IDSATEDRCGVCLGSGSCQTVRKIKQKE-GSGYVDYG
	gi 11493589	LGSDLREDRCRVCGGGSACETIEGVSPASPGAGYEDVV
	gi 10645199	IDSAMEDRCGVCHGSGTCHTVSGTTEAE-GLGYVDYG
	gi 15309931	LGSDLREDRCRVCGGGSACETIEGVSPASPGAGYEDVV
	gi 7656869	LGSDAREDRRCRVCGGSGTCDALIEGFNDSLPRGYMEVV
45		890 900 910 920
	NOV1a	TIPSGARSIRIYE N YSTS L YVRN-ALRRYYLNGHWTVD
	NOV1b	TIPSGARSIRIYE N YSTS L YVRN-ALRRYYLNGHWTVD
50	NOV1c	TIPSGARSIRIYE N YSTS L YVRN-ALRRYYLNGHWTVD
	NOV1d	TIPSGARSIRIYE N YSTS L YVRN-ALRRYYLNGHWTVD
	gi 13569928	LIPKGARDIRIMELEGAGHIALRSEDPEYYLNGGWTIQ
	gi 11493589	WIPKGVHIEFDLNLNLSLHIALGDO-ESLLLECLPGTP
	gi 10645199	LIPAGAREIRIOEVAELANTALRSEDPEYYLNGGWTIQ
55	gi 15309931	WIPKGVHIEFDLNLNLSLHIALGDO-ESLLLECLPGTP
	gi 7656869	QIPRGVHIEVREVALSKNYIALKSEG-DDYYDNGAWTLD
		930 940 950 960
60	NOV1a	WPGRYFEGTTFDYRRSYNPELIATGPTNE L L V L L F
	NOV1b	WPGRYFEGTTFDYRRSYNPELIATGPTNE L L V L L F
	NOV1c	WPGRYFEGTTFDYRRSYNPELIATGPTNE L L V L L F
	NOV1d	WPGRYFEGTTFDYRRSYNPELIATGPTNE L L V L L F
	gi 13569928	WQNYLALGVFOYDRKGD-LEKLMTAGPTNE L L V L L F
65	gi 11493589	QPHRLPLAGTTFQLRQGPQVQSLALGPINASLIVMYLA
	gi 10645199	WQDYQVAGTTFTHARRGN-WEALTPGPTKEPTWID---
	gi 15309931	QPHRLPLAGTTFQLRQGPQVQSLALGPINASLIVMYLA
	gi 7656869	WPRKFDVAGTAFHYRPTDPELEALGPTSENLIVMYLL
70		970 980 990 1000





		1210	1220	1230	1240
5	NOV1a	-----	-----	SEC	VSCGGGRC
	NOV1b	-----	-----	SEC	VSCGGGRC
	NOV1c	-----	-----	SEC	VSCGGGRC
	NOV1d	-----	-----	SEC	VSCGGGRC
	gi 13569928	SMSTSTPAISSPSPTTASKEGDLGGKQWQDS	--	STQPELS	
10	gi 11493589	---PRPPVLEACHGPTCPPEW	--AALDWSEC	TPSCGPCLR	
	gi 10645199	-T-SVSPSLKWPNLVAAVHRG	---GWGQA	--PLGLGGW	
	gi 15309931	---PRPPVLEACHGPTCPPEW	--AALDWSEC	TPSCGPCLR	
	gi 7656869	-----	-----	FAS	SCN--LA
15	NOV1a	-----	-----	-----	-----
	NOV1b	-----	-----	-----	-----
	NOV1c	-----	-----	-----	-----
	NOV1d	-----	-----	-----	-----
20	gi 13569928	SRYLSTGTSQPIILTSQS	SIQPSEENVSSSDTGPTSEG		
	gi 11493589	HRVVLCKSA	DHRATLPPAHCS	PAAKP	
	gi 10645199	RRHLVLMG	---PRLPTQL	PQE	---SNP
	gi 15309931	HRVVLCKSA	DHRATLPPAHCS	PAAKP	
	gi 7656869	KETLL			
25		1290	1300	1310	1320
	NOV1a	-----	-----	-----	-----
30	NOV1b	-----	-----	-----	-----
	NOV1c	-----	-----	-----	-----
	NOV1d	-----	-----	-----	-----
	gi 13569928	GLVATTTSGSGLSSSRNPITWPVTPFYNTLT	KGPEMEIHS		
	gi 11493589	-----	-----	-----	PAT
	gi 10645199	G	-----	VHYEYT	-----IHR
35	gi 15309931	-----	-----	-----	PAT
	gi 7656869	-----	-----	-----	-----
40	NOV1a	-----	-----	-----	-----
	NOV1b	-----	-----	-----	-----
	NOV1c	-----	-----	-----	-----
	NOV1d	-----	-----	-----	-----
45	gi 13569928	GSGEEREQPEDKDES	NPVIWTKIRVPGNDAPVESTEMPLA		
	gi 11493589	MRCNLRRCPP	-----	-----	-----
	gi 10645199	EAGGHDEVPP	-----	-----	-----
	gi 15309931	MRCNLRRCPP	-----	-----	-----
	gi 7656869	-----	-----	-----	-----
50		1370	1380	1390	1400
	NOV1a	-----	-----	-----	-----
	NOV1b	-----	-----	-----	-----
	NOV1c	-----	-----	-----	-----
	NOV1d	-----	-----	-----	-----
55	gi 13569928	PPLTPDLSRESWWPPF	STVMEGLLP	SQRPTTSETGTPRVE	
	gi 11493589	-----	-----	-----	-----
	gi 10645199	-----	-----	-----	-----
	gi 15309931	-----	-----	-----	-----
	gi 7656869	-----	-----	-----	-----
60		1410	1420	1430	1440
	NOV1a	-----	-----	-----	-----
65	NOV1b	-----	-----	-----	-----
	NOV1c	-----	-----	-----	-----
	NOV1d	-----	-----	-----	-----
	gi 13569928	GMVTEKPANTLLPLGGDHQPE	PSGKTANRNHLKLPNNMNQ		
	gi 11493589	-----	-----	-----	-----
70	gi 10645199	-----	-----	-----	-----

```

gi|15309931| -----
gi|7656869| -----

                    1450      1460      1470      1480
5      ....|....|....|....|....|....|....|....|
      NOV1a -----
      NOV1b -----
      NOV1c -----
      NOV1d -----
10     gi|13569928| TKSEPVLTEEDATSLITEGFLLNASNYKQLTNGHGSAAHW
      gi|11493589| -----ARW
      gi|10645199| ----PVFS-----W
      gi|15309931| -----ARW
      gi|7656869| -----

                    1490      1500      1510      1520
20     ....|....|....|....|....|....|....|....|
      NOV1a -----
      NOV1b -----
      NOV1c -----
      NOV1d -----
      gi|13569928| IVGNWSECSTTCGLGAYWKRVECTTQMDSDCAAIQRPDPA
      gi|11493589| VAGEWGECSAQCGVGQRQSRVCTS-----
      gi|10645199| HYGPWTKCTVTTCGRGE-----
25     gi|15309931| VAGEWGECSAQCGVGQRQSRVCTS-----
      gi|7656869| -----

                    1530      1540      1550      1560
30     ....|....|....|....|....|....|....|....|
      NOV1a -----
      NOV1b -----
      NOV1c -----
      NOV1d -----
      gi|13569928| KRCHLRPCAGWKVGNWSKCSRNCSTGGFKIREIQCVDSRDH
35     gi|11493589| -----HTGQASHECTEALR-----
      gi|10645199| -----KWGRHSPTCRG-----LVSGQGH
      gi|15309931| -----HTGQASHECTEALR-----
      gi|7656869| -----

                    1570      1580      1590      1600
40     ....|....|....|....|....|....|....|....|
      NOV1a -----
      NOV1b -----
      NOV1c -----
      NOV1d -----
45     gi|13569928| RNLRPFHCCQFLAGIPPLSMSCNPEPCAWQVEPWSQCSR
      gi|11493589| -----PPTTQQCEA
      gi|10645199| -----WLQLPA
      gi|15309931| -----PPTTQQCEA
50     gi|7656869| -----

                    1610      1620      1630      1640
55     ....|....|....|....|....|....|....|....|
      NOV1a -----
      NOV1b -----
      NOV1c -----
      NOV1d -----
      gi|13569928| SCGGGVQERGVFCPGLCDWTKRPTSTMSCNEHLCCHWAT
      gi|11493589| KCDS-----PTPG
60     gi|10645199| HC-----WAT
      gi|15309931| KCDS-----PTPG
      gi|7656869| -----

                    1650      1660      1670      1680
65     ....|....|....|....|....|....|....|....|
      NOV1a -----
      NOV1b -----
      NOV1c -----
      NOV1d -----
70     gi|13569928| GNWDLCSSTSCGGGFQKRIVQCVPSSEGNKTEDQDQCLCDHK

```

```

5  gi|11493589| DGPEECKDVN-----
   gi|10645199| TGLEVCFS-----E-----
   gi|15309931| DGPEECKDVN-----
   gi|7656869|  -----
                                     1690 1700 1710 1720
   NOV1a      .....|.....|.....|.....|.....|.....|.....|.....|
   NOV1b      -----
   NOV1c      -----
   NOV1d      -----
   gi|13569928| PRPPEFKKCNQQACKKSADLLCTKDKLSASFQTLKAMKK
   gi|11493589| -----KVAYCPLVLKRFQF
   gi|10645199| ---P-----QF-----S-ICEMRLAIAL
   gi|15309931| -----KVAYCPLVLKRFQF
   gi|7656869|  -----
                                     1730 1740 1750
   NOV1a      .....|.....|.....|.....|.....|.....|.....|..
   NOV1b      -----
   NOV1c      -----
   NOV1d      -----
   gi|13569928| CSVPTVRAECCFSCPQTHITHTQRQRRLQKSKEL
   gi|11493589| CSRAYFRQMCCCKTCQGH-----
   gi|10645199| CPRPAGRVHG-----
   gi|15309931| CSRAYFRQMCCCKTCHGH-----
   gi|7656869|  -----

```

30

The presence of identifiable domains in NOV1, as well as all other NOVX proteins, was determined by searches using software algorithms such as PROSITE, DOMAIN, Blocks, Pfam, ProDomain, and Prints, and then determining the Interpro number by crossing the domain match (or numbers) using the Interpro website (<http://www.ebi.ac.uk/interpro>).

35 DOMAIN results for NOV1 as disclosed in Tables 1L-IO, were collected from the Conserved Domain Database (CDD) with Reverse Position Specific BLAST analyses. This BLAST analysis software samples domains found in the Smart and Pfam collections. For Table 1E and all successive DOMAIN sequence alignments, fully conserved single residues are indicated by black shading or by the sign (!) and "strong" semi-conserved residues are

40 indicated by grey shading or by the sign (+). The "strong" group of conserved amino acid residues may be any one of the following groups of amino acids: STA, NEQK, NHQK, NDEQ, QHRK, MILV, MILF, HY, FYW.

Tables 1L-IO lists the domain description from DOMAIN analysis results against NOV1. This indicates that the NOV1 sequence has properties similar to those of other

45 proteins known to contain this domain.

**Table 1L. Domain Analysis of NOV1**

gnl|Pfam|pfam01421, Reprolysin, Reprolysin (M12B) family zinc metalloprotease. The members of this family are enzymes that cleave peptides. These proteases require zinc for catalysis. Members of this family are also known as adamalysins. Most members of this family are snake venom endopeptidases, but there are also some mammalian proteins, and fertilin. Fertilin and closely related proteins appear to not have some active site residues and may not be active enzymes. (SEQ ID NO:76)  
 CD-Length = 199 residues, 95.0% aligned  
 Score = 122 bits (307), Expect = 7e-29

5 Query: 378 DKKMMQNHG--HENITTYVLTLNMVSALFKDGTIGGNINIAIVGLILLEDEQPGLVISH 435  
 | | + | | | +|+|+ +++ | | + +| | | + | + +  
 Sbjct: 10 DHGMFTKYGSDLNKIRQRVHQIVNLVNEIYRPL---NIRVVLVGLLEIWSGDGK-ITVQG 64

10 Query: 436 HADHTLSSFCQWQSGLMGKDGTRHDHAILLTGLDICSWKNEPCDTLGFAPISGMCSKYRS 495  
 | + | | | +|+ + | | | +| | | | +| +| + | | | | |  
 Sbjct: 65 DANDTLHRFLEWRETDLLKR-KSHDNAQLLTGIDF-----DGNTIGAAAYVGGMCSPKRS 117

15 Query: 496 CTINED---TGLGLAFTIAHESGHNFMIHDGEGN-MCKKSEGNIMSPTLAGRNGVFSWS 551  
 + + | | + | +| | | | | | | | | + | | | + | +|  
 Sbjct: 118 VGVVQDHSPIVLLVAVTMAHELGHNLGMTHDDINKCTCEGGGGCIMNPVASSSPGKK-FS 176

Query: 552 PCSRQYLHKFLSTAQAICLADQ 573  
 | | +| | + + | | ++  
 Sbjct: 177 NCSMDDYQQFLTQKGPQCLLNK 198

**Table 1M. Domain Analysis of NOV1**

gnl|Smart|smart00209, TSP1, Thrombospondin type 1 repeats; Type 1 repeats in thrombospondin-1 bind and activate TGF-beta. (SEQ ID NO:77)  
 CD-Length = 51 residues, 100.0% aligned  
 Score = 63.5 bits (153), Expect = 5e-11

20 Query: 668 WSDWSSWSPCSRTCGGGVSHRSRLCTNPNSHGGKFCEGSTRTLKLCNSQKCP 720  
 | +| | | | | | | | | | +| | | | | | | | + | | | | |  
 Sbjct: 1 WGEWSEWSPCSVTCTGGGVQTRTRCCNPPN--GGGPCTGPDTEACNEQPCP 51

**Table 1N. Domain Analysis of NOV1**

gnl|Pfam|pfam00090, tsp\_1, Thrombospondin type 1 domain. (SEQ ID NO:78)  
 CD-Length = 48 residues, 100.0% aligned  
 Score = 54.7 bits (130), Expect = 2e-08

25 Query: 669 SDWSSWSPCSRTCGGGVSHRSRLCTNPNSHGGKFCEGSTRTLKLCNSQKC 719  
 | | | | | | | | + | | | +| | | | | + + | | |  
 Sbjct: 1 SPWSEWSPCSVTCTGKGI RTRQRTCNSP---AGGK PCTGDAQETEACMMDP 48

**Table 10. Domain Analysis of NOV1**

gnl|Pfam|pfam01562, Pep\_M12B\_propep, Reprolysin family propeptide.  
This region is the propeptide for members of peptidase family M12B.  
The propeptide contains a sequence motif similar to the "cysteine  
switch" of the matrixins. This motif is found at the C terminus of the  
alignment but is not well aligned. repeats. (SEQ ID NO:79)  
CD-Length = 117 residues, only 39.3% aligned  
Score = 38.5 bits (88), Expect = 0.002

Query: 180 LRTSSSLVAPGFIVQTLGKTGTSVQTLPPEDFCFYQGSLSRSHRNS 225  
| + ||+|| | | || + +| ||| + + ||  
Sbjct: 2 LEKNRSLAPDFTVTTYDDGTLVTEHPLIQDHCYQGYVEGYPN 47

5

Thrombospondin-1 (THBS1) associates with the extracellular matrix and inhibits angiogenesis in vivo. In vitro, THBS1 blocks capillary-like tube formation and endothelial cell proliferation. The antiangiogenic activity is mediated by a region that contains 3 type 1 (properdin or thrombospondin) repeats. By searching an EST database for sequences containing the antiangiogenic motif of THBS1, Vazquez et al. (1999) identified heart and lung cDNAs encoding ADAMTS1 and ADAMTS8, which they called METH1 and METH2, respectively. Sequence analysis predicted that the 890-amino acid ADAMTS8 protein shares 52% amino acid identity with ADAMTS1. ADAMTS8 is a secreted protein that has an N-terminal signal peptide, a zinc metalloprotease domain containing a zinc-binding site, and a cysteine-rich region containing 2 putative disintegrin loops. The C terminus of ADAMTS8 has 2 heparin-binding thrombospondin repeats with 6 cys and 3 trp residues. Southern blot analysis showed that ADAMTS8 is a single-copy gene distinct from that encoding ADAMTS1. Northern blot analysis detected highest expression of a 3.7-kb ADAMTS8 transcript in adult and fetal lung, with lower expression in brain, placenta, heart, and stomach, as well as fetal brain and kidney. Expression was also detected in a colon carcinoma cell line. SDS-PAGE analysis demonstrated that ADAMTS8 is expressed as a 98-kD protein, a 79-kD protein after cleavage at the subtilisin site, or as a 64-kD protein, which is most abundant, generated by an additional processing event. Functional analysis determined that ADAMTS8 disrupts angiogenesis in vivo and in vitro more efficiently than THBS1 or endostatin but somewhat less efficiently than ADAMTS1.

By interspecific backcross analysis, Georgiadis et al. (1999) mapped the mouse Adamts8 gene to chromosome 9 in a region showing homology of synteny with human 11q23-qter. They mapped the human ADAMTS8 gene to 11q25 by PCR analysis of a radiation hybrid mapping panel. The authors noted that a number of disorders have been mapped in the vicinity of the ADAMTS8 gene in mice and humans, most notably, given the expression and functional analyses, lung neoplasms.

The novel protein described here contains thrombospondin type I domains and Reprolysin domain. It is homologous to ADAM genes. Thrombospondin type I domain [IPR000884; (TSP1)] was found in the thrombospondin protein where it is repeated 3 times. Now a number of proteins involved in the complement pathway (properdin, C6, C7, C8A, C8B, C9) as well as extracellular matrix protein like mindin, F-spondin, SCO-spondin and even the circumsporozoite surface protein 2 and TRAP proteins of Plasmodium contain one or more instance of this repeat. It has been involved in cell-cell interaction, inhibition of angiogenesis, apoptosis. The intron-exon organisation of the properdin gene confirms the hypothesis that the repeat might have evolved by a process involving exon shuffling. A study of properdin structure provides some information about the structure of the thrombospondin type I repeat.

Reprolysin family propeptide [IPR002870; (Pep\_M12B\_propep)] domain is contained in the propeptide for members of peptidase family M12B. The propeptide contains a sequence motif similar to the 'cysteine switch' of the matrixins. This motif is found at the C terminus of the alignment but is not well aligned.

Through a subtractive hybridization approach to identify genes specifically expressed in the caput epididymidis, the mouse homologue of a member of the ADAM (a disintegrin and metalloprotease) family of proteins was identified. This rapidly growing gene family encodes cell surface proteins that possess putative adhesion and protease domains. Northern blot analyses demonstrated that the mouse ADAM gene, termed ADAM7, is expressed in the caput region of the epididymis and in the anterior pituitary gonadotropes with no detectable expression in the twenty-six other tissues examined. Furthermore, in situ hybridization experiments revealed that the ADAM7 messenger RNA (mRNA) exhibited an apical localization within the proximal caput epididymal epithelium that may correlate with an unusual sparsely granulated endoplasmic reticulum uniquely present in the proximal region of the epididymidis and to which no known function has been ascribed. Hormonal, surgical, and genetic strategies demonstrated that ADAM7 gene expression requires, in a region-dependent manner, androgens as well as testicular factors for expression. Interestingly, the apical localization of ADAM7 mRNA is dependent upon an intact testis, because in situ hybridization analyses of the proximal caput epididymidis from a testosterone maintained castrate mouse did not show the apical localization of ADAM7 mRNA. Finally, chromosomal mapping demonstrated that the ADAM7 gene maps to the central region of mouse Chromosome 14, approximately 4-5 cM distal from the fertilin beta locus, which encodes another reproductive-specific ADAM protein (1).

Because of the presence of the domain and the homology to the , we anticipate that the novel sequence described here will have useful properties and functions similar to these genes.

The disclosed NOV1 nucleic acid of the invention encoding a ADAM-TS 7 -like protein includes the nucleic acid whose sequence is provided in Table 1A, 1C, 1E, and 1G or a  
5 fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 1A, 1C, 1E and 1G while still encoding a protein that maintains its ADAM-TS 7-like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments  
10 that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical  
15 stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 33% percent of the bases may be so changed.

The disclosed NOV1 protein of the invention includes the ADAM-TS 7-like protein whose sequence is provided in Table 1B, 1D, 1F, or 1H. The invention also includes a mutant  
20 or variant protein any of whose residues may be changed from the corresponding residue shown in Table 1B, 1D, 1F, or 1H while still encoding a protein that maintains its ADAM-TS 7 -like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 62% percent of the residues may be so changed.

The invention further encompasses antibodies and antibody fragments, such as F<sub>ab</sub> or  
25 (F<sub>ab</sub>)<sub>2</sub>, that bind immunospecifically to any of the proteins of the invention.

The above defined information for this invention suggests that this ADAM-TS 7-like protein (NOV1) may function as a member of a "ADAM-TS 7 family". Therefore, the NOV1 nucleic acids and proteins identified here may be useful in potential therapeutic applications implicated in (but not limited to) various pathologies and disorders as indicated below. The  
30 potential therapeutic applications for this invention include, but are not limited to: protein therapeutic, small molecule drug target, antibody target (therapeutic, diagnostic, drug targeting/cytotoxic antibody), diagnostic and/or prognostic marker, gene therapy (gene delivery/gene ablation), research tools, tissue regeneration *in vivo* and *in vitro* of all tissues and cell types composing (but not limited to) those defined here.



The NOV1 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in cancer including but not limited to various pathologies and disorders as indicated below. For example, a cDNA encoding the ADAM-TS 7-like protein (NOV1) may be useful in gene therapy, and the ADAM-TS 7-like protein (NOV1) may be useful when administered to a subject in need thereof. By way of nonlimiting example, the compositions of the present invention will have efficacy for treatment of patients suffering from adrenoleukodystrophy, Alzheimer's disease, autoimmune disease, allergies, addiction, anxiety, ataxia-telangiectasia, asthma, ARDS, atherosclerosis, behavioral disorders, aortic stenosis, atrial septal defect (ASD), atrioventricular (A-V) canal defect, ductus arteriosus, allergy, cerebral palsy, congenital adrenal hyperplasia, cirrhosis, cardiomyopathy, congenital heart defects, diabetes, diverticular disease, epilepsy, emphysema, endometriosis, endocrine dysfunctions, graft versus host disease, glomerulonephritis, graft versus host disease (GVHD), growth and reproductive disorders, hemophilia, hypercoagulation, hypercalcaemia, Huntington's disease, hypertension, hypogonadism, fertility, idiopathic thrombocytopenic purpura, immunodeficiencies, interstitial nephritis, IgA nephropathy, lymphoedema, inflammatory bowel disease, Lesch-Nyhan syndrome, leukodystrophies, multiple sclerosis, muscular dystrophy, myasthenia gravis, neurodegeneration, neuroprotection, obesity, Parkinson's disease, pain, polycystic kidney disease, pulmonary stenosis, pancreatitis, renal artery stenosis, renal tubular acidosis, stroke, systemic lupus erythematosus, scleroderma, subaortic stenosis, transplantation, tuberous sclerosis, Von Hippel-Lindau (VHL) syndrome, ventricular septal defect (VSD), valve diseases, Von Hippel-Lindau (VHL) syndrome, ulcers, bacterial and viral infections, neonatal apnea, eagle's syndrome, atherosclerosis, metabolic disorders such as familial amyloidotic polyneuropathy, hyperkinetic diseases, muscular dystrophy, cerebral vascular disease, hypertension, cardiovascular diseases, renal fibrogenesis, inflammatory bowel disease, Meckel syndrome, colorectal cancer, papillomavirus infection and cervical carcinoma, liver malignancies, skin disorders, connective tissue disorders such as type VIIC Ehlers-Danlos syndrome, osteoarthritis, rheumatoid arthritis, or other pathologic or conditions. The NOV1 nucleic acid encoding the ADAM-TS 7-like protein of the invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

NOV1 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immuno-specifically to the novel NOV1 substances for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the

art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. The disclosed NOV1a and b proteins have multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated NOV1a and b epitope is from about amino acids 20 to 70. In another embodiment, a NOV1a and b epitope is from about amino acids 80 to 180. In additional embodiments, NOV1a and b epitopes are from about amino acids 200 to 280, from about amino acids 300 to 360, from about amino acid 400 to 530, from about amino acid 540 to 550, from about amino acid 580 to 610, from about amino acid 630 to 680, and from about amino acids 710 to 750. Also, the disclosed NOV1c protein has multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated NOV1c epitope is from about amino acids 1 to 50. In another embodiment, a NOV1c epitope is from about amino acids 80 to 100. In additional embodiments, NOV1c epitopes are from about amino acids 110 to 260, from about amino acids 280 to 320, from about amino acid 350 to 400, from about amino acid 420 to 470, from about amino acid 480 to 620, from about amino acid 700 to 750, and from about amino acids 770 to 810. Also, the disclosed NOV1d protein has multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated NOV1d epitope is from about amino acids 20 to 180. In another embodiment, a NOV1d epitope is from about amino acids 190 to 280. In additional embodiments, NOV1d epitopes are from about amino acids 300 to 360, from about amino acids 400 to 530, from about amino acid 540 to 550, from about amino acid 580 to 610, from about amino acid 630 to 680, and from about amino acids 710 to 750. These novel proteins can be used in assay systems for functional analysis of various human disorders, which will help in understanding of pathology of the disease and development of new drug targets for various disorders.

## NOV2

A disclosed NOV2 nucleic acid of 4488 nucleotides (also referred to as SC\_78316254\_A) encoding a novel alpha-2-macroglobulin precursor-like protein is shown in Table 2A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 1-3 and ending with a TGA codon at nucleotides 4477-4479. A putative untranslated region upstream from the initiation codon and downstream from the termination codon is underlined in Table 2A, and the start and stop codons are in bold letters.

Table 2A. NOV2 nucleotide sequence (SEQ ID NO:9).

**ATG**TGGGCTCAGCTCCTTCTAGGAATGTTGGCCCTATCACCAGCCATTGCAGAAGAACTTCCAAACTACCTG  
 GTGACATTACCAGCCCGGCTAAATTTCCCTCCGTTTCTGAGAGGTTGTTTGGACCTGAGCCCTGGGTACAGT  
 GATGTTAAATTCACGGTTACTCTGGAGACCAAGGACAAGACCCAGAAGTTGCTAGAACTACTCTGGACTGAAG  
 AAGAGGCACCTACATTGTATCTCCTTTCTGTACCACCTCCTGCTGGTGGCACAGAAGAAGTGCCACAATC  
 CGGGTGTGCGGAGTTGGAAATAACATCAGCTTTGAGGAGAAGAAAAAGGTTCTAATTCAGAGGCAGGGGAAC  
 GGCACCTTTGTACAGACTGACAAACCTCTCTACACCCAGGGCAGCAAGTGTATTTCCGCATTGTCAACATG  
 GATAGCAACTTCGTTCCAGTGAATGACAAGTACTCCATGGTGGAACTACAGGATCCAAATAGCAACAGGATT  
 GCACAGTGGCTGGAAGTGGTACCTGAGCAAGGCATTGTAGACCTGTCTTCCAACCTGGCACAGAGGCAATG  
 CTGGGCACCTACACTGTGGCAGTGGCTGAGGGCAAGACCTTTGGTACTTTTCTGAGTGGAGGAATATGTGCTT  
 TCTCCATTTCTCCTTTTACTCTCTTCTCAGTGCTGCCGAAGTTTAAAGTGGAAGTGGTGGAAACCCAGAGTTA  
 TCAACGGTGCAGGAATCTTTCTTAGTAAAAATTTGTTGTAGGTACACCTATGGAAGCCCATGCTAGGGGCA  
 GTGCAGGTATCTGTGTGTCAGAAGGCAAACTTACTGTTATCGAGAGGTGGAACGGGAACAGCTTCTGAC  
 AAATGCAGGAACCTCTCTGGACAGACTGACAAAAACAGGATGTTTCTCAGCACCTGTGGACATGGCCACCTTT  
 GACCTCATTGGATATGCGTACAGCCATCAAATCAATATTGTGGTACTGTTGTGGAGGAAGGACAGGTGTG  
 GAGGCCAATGCCACTCAGAATATCTACATTTCTCCACAAATGGGATCAATGACCTTTGAAGACACCAGCAAT  
 TTTTACCATCCAAATTTCCCTTCTCAGTGGGAAGATGCTGCTCAAGTTTCCGCAAGGCGGTGTGCTCCCTTGC  
 AAGAACCATCTAGTGTCTTCTGGTGATTATGGCACAAATGGAACCTTCAACCAGACCTGGTACTGATAAC  
 AATGGCCTAGCTCCCTTTACCTTGGAGACATCCGGTTGGAATGGGACAGACGTTTCTCTGGAGGGAAAGTTT  
 CAAATGGAAGACTTAGTATATAATCCGGAACAAGTGCCACGTTACTACCAAATGCCTACCTGCACCTGCCA  
 CCTTTCTACAGCACAAACCCGAGCTTCTTGGCATCCACCGGCTAAACGGCCCCCTTGAATGTGGCCAGCCC  
 CAGGAAGTGTGGTGGATTATTACATCGACCCGGCCGATGCAAGCCCTGACCAAGAGATCAGCTTCTCCTAC  
 TATTTAATAGGGAAAGGAAGTTTGGTGATGGAGGGGCAGAAACACCTGAACCTAAGAGAAAGGACTGAAA  
 GCCTCCTTCTCTCTCTCACTGACCTTCACTTCGAGACTGGCCCCGATCCTTCCCTGGTGTATGATGCCATT  
 TTTCCAGTGGAGGTGTTGTAGCTGACAAAAATTCAGTTCTCAGTCGAGATGTGCTTTGACAACTCAGCAGCTT  
 CCAGGAGCAGAAGTGGAGCTGCAGCTGCAGGCAGCTCCCGGATCCCTGTGTGCGCTCCGGGCGGTGGATGAG  
 AGTGTCTTACTGCTTAGGCCAGACAGAGAGCTGAGCAACCGCTCTGTCTATGGGATGTTTCCATTCTGGTAT  
 GGTCATACCCCTATCAAGTGGCTGAGTATGATCAGTGTCCAGTGTCTGGCCCATGGGACTTTCTCAGCCC  
 CTCATTGACCCCAATGCCCCAAGGGCATTGAGCCAGCGTTCATTATCTGGAGGCCCTGTTCTCTGAAGGC  
 ACGGACCTTTTTCAGCTTTTCCGGGACGTGGGCTGAAAATACTGTCCAATGCCAAAATCAAGAAGCCAGTA  
 GATTGCAGTCACAGATCTCCAGAATACAGCACTGCTATGGGTGGCGGTGGTCACTCAGAGGCTTTTGAGTCA  
 TCAACTCCTTTACATCAAGCAGAGGATTCTCAGGTCCGCCAGTACTTCCCAGAGACCTGGCTCTGGGATCTG  
 TTTCTTATTGGTAACCTCGGGGAAGGAGGCGGTCCACGTCACAGTTTCTGACGCCATCACCGAGTGAAGGCG  
 ATGAGTTTCTGCACTTCCAGTCAAGAGGCTTCGGGCTTCAACCACTGTTGGACTAAGCTGCTTTCAAGCCG  
 TTCTTTGTGACCTGACTCTCCCTTACTCAGTAGTCCGTGGGAATCCTTTCTGTTACTGCCACCATTCTC  
 AATTACCTAAAGGATTGCATCAGGGTTCAGACTGACCTGGCTAAATCGCATGAGTACCAGCTAGAATCATGG  
 GCAGATTCTCAGACCTCCAGTTGTCTCTGTGTGATGACGCAAAACCCACCACTGGAACATCACAGCTGTC  
 AAATTTGGGTACATTAACCTTACTATTAGTACAAAGATTCTGGACAGCAATGAACCATGTGGGGGCCAGAAG  
 GGGTTTGTTCCTCCCAAAAGGGCCGAAGTGACACGCTCATCAAGCCAGTTCTCGTCAAACTGAGGAGCTCTG  
 GTGGAGAAGACACACAGCTCATTGCTGTGCCCAAAAGGAGGAAAGGTGGCATCTGAATCTGTCTCCCTGGAG  
 CTCCCAGTGGACATGTTCCTGACTCGACCAAGGCTTATGTTACGTTCTGGGAGACATTATGGGCACAGCC  
 CTGCAGAACCTGGATGGTCTGGTGCAGATGCCAGTGGCTGTGGCGAGCAGAACATGGTCTTGTGTTGCTCCC  
 ATCATCTATGTCTTGCAGTACCTGGAGAAGGAGGCTGCTGACGGAGGAGATCAGGTCTCGGGCAGTGGGT  
 TTCCTGGAAATAGGTTACCAGAAGGAGCTGATGTACAAACACAGCAATGGCTCATACTAGTGCCTTTGGGGAG  
 CGAGATGGAATGGAACACATGGCTGACAGCGTTTGTACAAATGCTTTGGCCAAGCTCAGAAATTCATC  
 TTCATTGATCCCAAGAACATCCAGGATGCTCTCAAGTGGATGGCAGGAAACAGCTCCCCAGTGGCTGCTAT  
 GCCAACGTGGGAAATCTCCTTACACAGCTATGAAGGGTGGTGTGATGATGAGGTCTCCTTGACTGCGTAT  
 GTCACAGCTGCATTGCTGGAGATGGGAAAGGATGTAGATGACCCAATGGTGAGTCAGGCTACCGGTGTCTC  
 AAGAATTCGGCCACCTCCACGACCAACCTCTACACACAGGCCCTGTGGCTTACATTTTCTCCCTGGCTGGG  
 GAAATGGACATCAGAAACATTTCTCCTTAAACAGTTAGATCAACAGGCTATCATCTCAGGAGAATCCATTTAC  
 TGGAGCCAGAAACCTACTCCATCATCGAACGCCAGCCCTTGGTCTGAGCCTGCGGCTGTAGATGTGGAACCTC  
 ACAGCATATGCATTGTTGGCCCAGCTTACCAAGCCCAGCCTGACTCAAAAGGAGATAGCGAAGGCCACTAGC  
 ATAGTGGCTTGGTTGGCCAAGCAACACAATGCATATGGGGGCTTCTCTTACTCAGGATACTGTAGTTGCT  
 CTCCAAGCTCTTGCCAAATATGCCACTACCGCTACATGCCATCTGAGGAGATCAACCTGGTTGTAAATCC  
 ACTGAGAATTTCCAGCGCACATTCAACATACAGTCAGTTAAACAGATTGGTATTTTCAAGGATACCCCTGCCC  
 AATGTCCCTCGAATGTACACGTTGGAGGCCTCAGGCCAGGGCTGTGTCTATGTGCAGACGGTGTGAGATAC  
 AATATTTCTCCCTCCACAAAATATGAAGACCTTTAGTCTTAGTGTGGAATAGGAAAAGCTAGATGTGAGCAA  
 CCGACTTCACCTCGATCCTTGACTCTCACTATTACACACAGTTATGTGGGGAGCCGTAGCTCTTCCAATATG  
 GCTATTGTGGAAGTGAAGATGCTATCTGGGTTTCACTCCATGGAGGGCACCAATCAGTTACTTCTCCAGCAA  
 CCCCTGGTGAAGAAGGTTGAATTTGGAACGTACACACTTAACATTTACTTGGATGAGCTCATTAGAACACT  
 CAGACTTACACCTTACCATCAGCCAAAGTGTGCTGGTACCAACTTGAACCAGCAACCATCAAGGTCTAT  
 GACTACTACCTACCAGGTTCTTTAAATTATCTCAGTACACAATTTGTGTGGTCCATGAGCAATGAGCATA  
 GTGGACTCTGTGGCACGGCACCCAGAACCCCTTTCAAGACAGAAGCATTTATTCCTTCACTTCTGGG  
 AGTGTTAAACAAC**GT**ATAGCTACCA

The disclosed NOV2 nucleic acid sequence, localized to chromosome 12, has 840 of 1324 bases (63%) identical to a *Rattus norvegicus* Alpha-2-Macroglobulin Precursor mRNA (GENBANK-ID: RATA2M) ( $E = 1.3e^{-119}$ ).

A NOV2 polypeptide (SEQ ID NO:10) encoded by SEQ ID NO:9 has 1492 amino acid residues and is presented using the one-letter code in Table 2B. Signal P, Psort and/or Hydropathy results predict that NOV2 does contain a signal peptide and is likely to be localized extracellularly with a certainty of 0.3703. In other embodiments, NOV2 may also be localized to the lysosome (lumen) with a certainty of 0.1900, the microbody (peroxisome) with a certainty of 0.1585, or the endoplasmic reticulum (membrane) with a certainty of 0.1000.

The most likely cleavage site for a NOV2 peptide is between amino acids 17 and 18, at: AIA-EE.

**Table 2B. Encoded NOV2 protein sequence (SEQ ID NO:10).**

```
MWAQLLGLMLALSPAIAEELPNYLVTLPARLNFPSVQKVLCLDLSPGYSDVKFTVTLETDKDKTKLLEYSGLK
KRHLHCISFLVPPAGGTEEVATIRVSGVGNNISFEEKKKVLIQRQNGTFVQTDKPLYTPGQQVYFRIVTM
DSNFPVNDKYSMVELQDPNSNRIQWLEVPEQGIVDLSFQLAPEAMLGTYTVAVAEGKTFGTFSVEEVVL
SPFLLLLSSVLPKFKVEVVEPKELSTVQESFLVKICCRYTYGKPMLGAVQSVQCQKANTYWYREVEREQLPD
KCRNLSSGQTDKTCGFSAPVDMATFDLIGYAYSHQINIVATVVEEGTGVEANATQNIYISPMGMSMTFEDTSN
FYHPNFPFSGKMLLKFPQGGVLPCKNHLVFLVIYGTNGTFNQTLVTDNGLAPFTLETSGWNGTDVSLGKGF
QMEDLVNPEQVPRYYQNAYLHLRPFYSTTRSFLGIHRLNGPLKCGQPQEVLDVYYIDPADASPDQEIISFSY
YLIGKGLVMGEGKHLNSKKKGLKASFSLSTFTSRLAPDPSLVIYAIFPSGGVVADKIQFSVMCFDNNQQL
PGAEEVLQQLQAAPGSLCALRAVDESLLLLRPDRELSNRSVYGMFPFWYGHYPYQVAEYDQCPVSGPWDFPQP
LIDPMPQGHSSQRSIIWRPSFSEGTDLFSFFRDVGLKILSNAKIKKPVDCSHRSPEYSTAMGGGGHPEAFES
STPLHQAEDSQVRQYFPETWLWDLFPIGNSGKEAVHVTVPDAITEWKAMSFCTSQSRGFGLSPTVGLTAFKP
FFVDLTLPYSVVRGESFRLTATIFNYLKDCIRVQTDLAKSHEYQLESWADSQTSSCLCADDKTHHWNITAV
KLGHINFITISTKILDSNEPCGGQKGFVPQKGRSDTLIKPVLVKPEGVLVEKTHSSLLCPKGGKVASESVSLE
LPVDIVPDSTKAYVTVLGDMGTALQNLQDGLVQMPSCGCEQNMVLFAPIIYVLQYLEKAGLLTEEIRSRVAVG
FLEIGYQKELMYKHSNGSYSAFGERDGNNTWLTAFVTKCFGQAQKFIIDPKNIQDALKWMAGNQLPSGCV
ANVGNLLHTAMKGGVDDEVS LTAYVTAALLEMGKDVEDPMVSQGLRCLKNSATSTTNLYTQALLAYIFSLAG
EMDIRNILLKQLDQQAIIISGESIYWSQKPTPSSNASPWSEPAAVDVLTAYALLAQLTKPSLTQKEIAKATS
IVAWLAKQHNAVGGSSTQDVTVALQALAKYATTAYMPSEEINLVVKSTENFQRTFNIQSVNRLVFQDQTLN
NVPGMYLEASGQGCYVYQTVLRYNLPPTNMKTFSLSVEIGKARCEQPTSPRSLTLTIHTSYVGSRSSNM
AIVEVKMLSGFSPMEGTNQLLLQQLPVKKVEFGTDLNIIYLDLTKNTQTYTFTISQSVLVTNLKPATIKVY
DYYLPGSFGLSKYQTIWVSMNNDISIVDSVARHPEPPPFKTEAFIPSLPGSVNN
```

The disclosed NOV2 amino acid sequence has 595 of 1450 amino acid residues (41%) identical to, and 873 of 1450 residues (60 %) positive with, the 1474 amino acid residue Alpha-2-Macroglobulin Precursor protein from *Homo sapiens* (ptnr:SPTREMBL-ACC:P01023) ( $E = 2.0e^{-279}$ ).

NOV2 is expressed in at least the following tissues: Hematopoietic tissues, blood plasma, fetal lung, and the coronary artery.

NOV2 also has homology to the amino acid sequences shown in the BLASTP data listed in Table 2C.

Table 2C. BLAST results for NOV2

Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 14765710 ref XP_006925.4	alpha 2 macroglobulin precursor [Homo sapiens]	1474	593/1486 (39%)	870/1486 (57%)	0.0
gi 4557225 ref NP_000005.1	alpha 2 macroglobulin precursor [Homo sapiens]	1474	591/1486 (39%)	869/1486 (57%)	0.0
gi 224053 prf  1009174A	macroglobulin alpha2 [Homo sapiens]	1450	585/1471 (39%)	861/1471 (57%)	0.0
gi 6978425 ref NP_036620.1	alpha-2-macroglobulin [Rattus norvegicus]	1472	578/1483 (38%)	867/1483 (57%)	0.0
gi 2144118 pir  JC5143	alpha-macroglobulin precursor - guinea pig	1476	570/1495 (38%)	858/1495 (57%)	0.0

The homology of these sequences is shown graphically in the ClustalW analysis shown in Table 2D.

Table 2D. ClustalW Analysis of NOV2

- 1) NOV2 (SEQ ID NO 10)
- 2) gi|14765710|ref|XP\_006925.4| alpha 2 macroglobulin precursor [Homo sapiens] (SEQ ID NO:35)
- 2) gi|4557225|ref|NP\_000005.1| alpha 2 macroglobulin precursor [Homo sapiens] (SEQ ID NO:36)
- 3) gi|224053|prf||1009174A macroglobulin alpha2 [Homo sapiens] (SEQ ID NO:37)
- 4) gi|6978425|ref|NP\_036620.1| alpha-2-macroglobulin [Rattus norvegicus] (SEQ ID NO:38)
- 5) gi|2144118|pir||JC5143 alpha-macroglobulin precursor - guinea pig (SEQ ID NO:39)

	570	580	590	600
NOV2	SLVLIYALP	SGGVADKIQ	SVEMCFDN	-----QQL
gi 14765710	RLLIYAVLP	GVIGDSAKY	VENCLANKVDLSFSP	QSL
gi 4557225	RLLIYAVLP	GVIGDSAKY	VENCLANKVDLSFSP	QSL
gi 224053	RLLIYAVLP	GVIGDSAKY	VENCLANKVDLSFSP	QSL
gi 6978425	RLVLYAVLP	GVIGDSAKY	VENCLANKVDLSFSP	QSL
gi 2144118	RLLIYAVLP	GVIGDSAKY	VENCLANKVDLSFSP	QSL
	610	620	630	640
NOV2	PGAEVELQ	CAAPCS	CALRAVD	SVLLLPDRELSNRSV
gi 14765710	PASHAHLRV	TAAPOS	CALRAVDQSVLLMKPDAELSASSV	
gi 4557225	PASHAHLRV	TAAPOS	CALRAVDQSVLLMKPDAELSASSV	
gi 224053	PASHAHLRV	TAAPOS	CALRAVDQSVLLMKPDAELSASSV	
gi 6978425	PATRALLSV	MASPOS	CGLRAVDQSVLLMKPDAELSASSV	
gi 2144118	PASKTHLRV	TAAPOS	CALRAVDQSVLLMKPDAELSASSV	
	650	660	670	680
NOV2	VGMFP	FWYGHY	PYQVAYD	QCPVSGPWDFPQPLIDPMPQ

gi|14765710| YLLPEKDLTGFPGLDQDND-----CINR  
gi|4557225| YLLPEKDLTGFPGLDQDND-----CINR  
gi|224053| YLLPEKDLTGFPGLDQDND-----CINR  
gi|6978425| YLLPVKDLTGFPGLDQDND-----CVKQ  
gi|2144118| YALLPVKDLTGFPGLDQDND-----CVSL

690 700 710 720

NOV2 GHSSQRSITIRPSFSEG-TDL SFFRDYGLKILEN KIK  
gi|14765710| ENVYINGITY PVSSSTNEKDMYSFL DMGLKFTNSKIRK  
gi|4557225| ENVYINGITY PVSSSTNEKDMYSFL DMGLKFTNSKIRK  
gi|224053| ENVYINGITY PVSSSTNEKDMYSFL DMGLKFTNSKIRK  
gi|6978425| NDTYINGILYSPVQ TNE DMYGFL DMGLKFTNSKIRK  
gi|2144118| NDTYINGILYSPVQ TNE DMYGFL DMGLKFTNSKIRK

730 740 750 760

NOV2 PVDGSHRSPEYST----AMGCGGHPAFESSTP-----  
gi|14765710| PKCPOLQOYE IHGPEGLRVCYESDVMGRGHARLVHV--  
gi|4557225| PKCPOLQOYE IHGPEGLRVCYESDVMGRGHARLVHV--  
gi|224053| PKCPOLQOYE IHGPEGLRVCYESDVMGRGHARLVHV--  
gi|6978425| PKVCELDNDKIPAAAYHLSQSHMDAFLE-----  
gi|2144118| PKCAHYQAEVP---TMAYSTSESSFRSGPRVPVAVGI

770 780 790 800

NOV2 ---LHQEDSQVRYFPETWIWDLFPKGNEGKEAVHVTVP  
gi|14765710| ---EPPHTETVR YFPETWIWDLVVV SAGVAEVGVTVP  
gi|4557225| ---EPPHTETVR YFPETWIWDLVVV SAGVAEVGVTVP  
gi|224053| ---EPPHTETVR YFPETWIWDLVVV SAGVAEVGVTVP  
gi|6978425| S---SESPHTETRRSYFPETWIWDLVVV SAGVAEVEVTVP  
gi|2144118| AATYSEPPKETVRYSFPETWIWDLVVV SAGVAEVEVTVP

810 820 830 840

NOV2 DAITWKAMSFCTSSSRGFGSPITVGLTAFKPPFVOLTVP  
gi|14765710| DTITWKAGAFCLS DAGLGTSSTASLRAFQPPFVELTMP  
gi|4557225| DTITWKAGAFCLS DAGLGTSSTASLRAFQPPFVELTMP  
gi|224053| DTITWKAGAFCLS DAGLGTSSTASLRAFQPPFVELTMP  
gi|6978425| DTITWKAGAFCLSNDTGLGSPVVQFAFQPPFVELTMP  
gi|2144118| DTITWKAGAFCLSNDTGLGSPPTASLRAFQPPFVELTMP

850 860 870 880

NOV2 YSVIRGEAFTLKATVNLNPKCIRVSVQLEASPAFLAVPV  
gi|14765710| YSVIRGEAFTLKATVNLNPKCIRVSVQLEASPAFLAVPV  
gi|4557225| YSVIRGEAFTLKATVNLNPKCIRVSVQLEASPAFLAVPV  
gi|224053| YSVIRGEAFTLKATVNLNPKCIRVSVQLEASPAFLAVPV  
gi|6978425| YSVIRGEAFTLKATVNLNPKCIRVSVQLEASPAFLAVPV  
gi|2144118| YSVIRGEAFTLKATVNLNPKCIRVSVQLEASPAFLAVPV

890 900 910 920

NOV2 ADSQTSSCLCADDATTHWNITAVKLGINFTLSTILLS  
gi|14765710| EKEQAPHICANGRQTVSWAVTPKSLGNVNFTVSAEALS  
gi|4557225| EKEQAPHICANGRQTVSWAVTPKSLGNVNFTVSAEALS  
gi|224053| EKEQAPHICANGRQTVSWAVTPKSLGNVNFTVSAEALS  
gi|6978425| EKEQASHCICMNORETTASWAVTPKSLGNVNFTVSAEALS  
gi|2144118| AKEQESHCVCCNERQTVSWAVTPKSLGNVNFTVSAEALS

930 940 950 960

NOV2 NEPCGGKGFVPCGRSDTIKPLVPEGVLVERKTHESL  
gi|14765710| CELCGTEVPSVPE GRKDTIKPLVPEGLEKETTFNSL  
gi|4557225| CELCGTEVPSVPE GRKDTIKPLVPEGLEKETTFNSL  
gi|224053| CELCGTEVPSVPE GRKDTIKPLVPEGLEKETTFNSL  
gi|6978425| CELCGTEVPSVPE GRKDTIKPLVPEGLEKETTFNSL  
gi|2144118| SELCGTEVPSVPE GRKDTIKPLVPEGLEKETTFNSL

970 980 990 1000

	NOV2	LCPSGGVLSSESLLPVDPVPISTAMVVLGDI.G.A
	gi   14765710	LCPSGGGEVS-E LSLKLPNNVVEESARASVSVLGDILGSA
5	gi   4557225	LCPSGGGEVS-E LSLKLPNNVVEESARASVSVLGDILGSA
	gi   224053	LCPSGGGEVS-E LSLKLPNNVVEESARASVSVLGDILGSA
	gi   6978425	LCPSGGGEVS-E LSLKLPNNVVEESARASVSVLGDILGSA
	gi   2144118	LRVSDTTVS-E LHL LPSNVVQ SARAVSVLGDILGSA
		1010 1020 1030 1040
10	NOV2	YQNLIDGLQMPSSGCGEONMVLFAPIIXVLOYLEAGLLTE
	gi   14765710	MQNTQNLQMPYGCGEONMVLFAPIIXVLDYLNQQLTP
	gi   4557225	MQNTQNLQMPYGCGEONMVLFAPIIXVLDYLNQQLTP
	gi   224053	MQNTQNLQMPYGCGEONMVLFAPIIXVLDYLNQQLTP
15	gi   6978425	MQNTQNLQMPYGCGEONMVLFAPIIXVLDYLNQQLTP
	gi   2144118	MQNTQNLQMPYGCGEONMVLFAPIIXVLDYLNQQLTP
		1050 1060 1070 1080
20	NOV2	EIKSKAIGYLNLTGYQRLNYKHMDGSYSTFGERYGRNQGN
	gi   14765710	EIKSKAIGYLNLTGYQRLNYKHMDGSYSTFGERYGRNQGN
	gi   4557225	EIKSKAIGYLNLTGYQRLNYKHMDGSYSTFGERYGRNQGN
	gi   224053	EIKSKAIGYLNLTGYQRLNYKHMDGSYSTFGERYGRNQGN
	gi   6978425	EIKSKAIGYLNLTGYQRLNYKHMDGSYSTFGERYGRNQGN
25	gi   2144118	EIKSKAIGYLNLTGYQRLNYKHMDGSYSTFGERYGRNQGN
		1090 1100 1110 1120
30	NOV2	TWLTAFVLKTFQAQARYIFIDEAHITQALWLSQOKDNG
	gi   14765710	TWLTAFVLKTFQAQARYIFIDEAHITQALWLSQOKDNG
	gi   4557225	TWLTAFVLKTFQAQARYIFIDEAHITQALWLSQOKDNG
	gi   224053	TWLTAFVLKTFQAQARYIFIDEAHITQALWLSQOKDNG
	gi   6978425	TWLTAFVLKTFQAQARYIFIDEAHITQALWLSQOKDNG
35	gi   2144118	TWLTAFVLKTFQAQARYIFIDEAHITQALWLSQOKDNG
		1130 1140 1150 1160
40	NOV2	CFRSSGSLNNAIKGGVEDEVTL SAYITIALLEPLTVTH
	gi   14765710	CFRSSGSLNNAIKGGVEDEVTL SAYITIALLEPLTVTH
	gi   4557225	CFRSSGSLNNAIKGGVEDEVTL SAYITIALLEPLTVTH
	gi   224053	CFRSSGSLNNAIKGGVEDEVTL SAYITIALLEPLTVTH
	gi   6978425	CFRSSGSLNNAIKGGVEDEVTL SAYITIALLEPLTVTH
	gi   2144118	CFRSSGSLNNAIKGGVEDEVTL SAYITIALLEPLTVTH
45		1170 1180 1190 1200
50	NOV2	PVVRNALFCLESARKAEGDHGSHVYTKALLAYAFALAG
	gi   14765710	PVVRNALFCLESARKAEGDHGSHVYTKALLAYAFALAG
	gi   4557225	PVVRNALFCLESARKAEGDHGSHVYTKALLAYAFALAG
	gi   224053	PVVRNALFCLESARKAEGDHGSHVYTKALLAYAFALAG
	gi   6978425	PVVRNALFCLESARKAEGDHGSHVYTKALLAYAFALAG
	gi   2144118	PVVRNALFCLESARKAEGDHGSHVYTKALLAYAFALAG
55		1210 1220 1230 1240
60	NOV2	NQDNRKEVLKSL EEA VK DNSVHWERPQKPKPVGHYE
	gi   14765710	NQDNRKEVLKSL EEA VK DNSVHWERPQKPKPVGHYE
	gi   4557225	NQDNRKEVLKSL EEA VK DNSVHWERPQKPKPVGHYE
	gi   224053	NQDNRKEVLKSL EEA VK DNSVHWERPQKPKPVGHYE
	gi   6978425	NQDNRKEVLKSL EEA VK DNSVHWERPQKPKPVGHYE
	gi   2144118	NQDNRKEVLKSL EEA VK DNSVHWERPQKPKPVGHYE
65		1250 1260 1270 1280
70	NOV2	PQAPSAEVENTSYVLLAYLTA PAPTSEDLT SAT IVKWW
	gi   14765710	PQAPSAEVENTSYVLLAYLTA PAPTSEDLT SAT IVKWW
	gi   4557225	PQAPSAEVENTSYVLLAYLTA PAPTSEDLT SAT IVKWW
	gi   224053	PQAPSAEVENTSYVLLAYLTA PAPTSEDLT SAT IVKWW
	gi   6978425	PQAPSAEVENTSYVLLAYLTA PAPTSEDLT SAT IVKWW
	gi   2144118	PQAPSAEVENTSYVLLAYLTA PAPTSEDLT SAT IVKWW

			1290	1300	1310	1320
5	NOV2		AKQHNAAGGFSSTQDTVVALHALSKYGAATFTRTKAAQV			
	gi   14765710		TKQQNAAGGFSSTQDTVVALHALSKYGAATFTRTKAAQV			
	gi   4557225		TKQQNAAGGFSSTQDTVVALHALSKYGAATFTRTKAAQV			
	gi   224053		TKQQNAAGGFSSTQDTVVALHALSKYGAATFTRTKAAQV			
	gi   6978425		TKQQNAAGGFSSTQDTVVALHALSKYGAATFTRTKAAQV			
10	gi   2144118		TKQQNAAGGFSSTQDTVVALHALSKYGAATFTRTKAAQV			
			1330	1340	1350	1360
15	NOV2		VLTSTENFORTENQVNRLLQOQVSLPEPGYKVTG			
	gi   14765710		VLTSTENFORTENQVNRLLQOQVSLPEPGYKVTG			
	gi   4557225		VLTSTENFORTENQVNRLLQOQVSLPEPGYKVTG			
	gi   224053		VLTSTENFORTENQVNRLLQOQVSLPEPGYKVTG			
	gi   6978425		VLTSTENFORTENQVNRLLQOQVSLPEPGYKVTG			
	gi   2144118		VLTSTENFORTENQVNRLLQOQVSLPEPGYKVTG			
20			1370	1380	1390	1400
	NOV2		EGCVYLQTSILKYNILPEKEEPPFALGVQTLPTQCDPKAH			
	gi   14765710		EGCVYLQTSILKYNILPEKEEPPFALGVQTLPTQCDPKAH			
	gi   4557225		EGCVYLQTSILKYNILPEKEEPPFALGVQTLPTQCDPKAH			
25	gi   224053		EGCVYLQTSILKYNILPEKEEPPFALGVQTLPTQCDPKAH			
	gi   6978425		EGCVYLQTSILKYNILPEKEEPPFALGVQTLPTQCDPKAH			
	gi   2144118		EGCVYLQTSILKYNILPEKEEPPFALGVQTLPTQCDPKAH			
30			1410	1420	1430	1440
	NOV2		TSFQISLVSYTGSRSSNMAIVDKMVSGFIPLKPTVKM			
	gi   14765710		TSFQISLVSYTGSRSSNMAIVDKMVSGFIPLKPTVKM			
	gi   4557225		TSFQISLVSYTGSRSSNMAIVDKMVSGFIPLKPTVKM			
	gi   224053		TSFQISLVSYTGSRSSNMAIVDKMVSGFIPLKPTVKM			
35	gi   6978425		TSFQISLVSYTGSRSSNMAIVDKMVSGFIPLKPTVKM			
	gi   2144118		TSFQISLVSYTGSRSSNMAIVDKMVSGFIPLKPTVKM			
40			1450	1460	1470	1480
	NOV2		LERSNHVSRTEVSNHVLIIYLDKVSNOTLSLFTVQDVP			
	gi   14765710		LERSNHVSRTEVSNHVLIIYLDKVSNOTLSLFTVQDVP			
	gi   4557225		LERSNHVSRTEVSNHVLIIYLDKVSNOTLSLFTVQDVP			
	gi   224053		LERSNHVSRTEVSNHVLIIYLDKVSNOTLSLFTVQDVP			
	gi   6978425		LERSNHVSRTEVSNHVLIIYLDKVSNOTLSLFTVQDVP			
45	gi   2144118		LERSNHVSRTEVSNHVLIIYLDKVSNOTLSLFTVQDVP			
50			1490	1500	1510	1520
	NOV2		VRDLKPAIVKVYDYETEFATAEYAPCSKDLGNA----			
	gi   14765710		VRDLKPAIVKVYDYETEFATAEYAPCSKDLGNA----			
	gi   4557225		VRDLKPAIVKVYDYETEFATAEYAPCSKDLGNA----			
	gi   224053		VRDLKPAIVKVYDYETEFATAEYAPCSKDLGNA----			
	gi   6978425		VRDLKPAIVKVYDYETEFATAEYAPCSKDLGNA----			
55	gi   2144118		VRDLKPAIVKVYDYETEFATAEYAPCSKDLGNA----			
60	NOV2		1530	1540		
	gi   14765710		ARHPEPPPFKTEAFIPSLPGSVNN			
	gi   4557225		-----			
	gi   224053		-----			
	gi   6978425		-----			
	gi   2144118		-----			
65						



Tables 2E-F list the domain description from DOMAIN analysis results against NOV2. This indicates that the NOV2 sequence has properties similar to those of other proteins known to contain this domain.

**Table 2E Domain Analysis of NOV2**

gnl|Pfam|pfam00207, A2M, Alpha-2-macroglobulin family. This family includes the C-terminal region of the alpha-2-macroglobulin family.  
(SEQ ID NO:80)  
CD-Length = 751 residues, 98.5% aligned  
Score = 563 bits (1451), Expect = 2e-161

5	Query: 728	EDSQVRQYFPETWLWDLFPIGNSGKEAVHVTVPDAITEWKAMSFACTSQRGFLSPTVGL	787
	Sbjct: 4	DDITIRSYFPESWLWEVEEVDSPVLTVNITLPDSITTWEILAVSLSNKGLCVADPVEL	63
10	Query: 788	TAFKPFVVDLTLPSVVRGESFRLTATIFNYL-KDCIRVQTDLAKSHEYQLESWADSQTS	846
	Sbjct: 64	TVFQDFLELRPLPSVVRGEQVELRAVLYNYLPSQDIKV-----VVQLEVEPLCQAG	115
15	Query: 847	SCLCADDAKTHHWNITAVKLGHINFITSTKILDSNEPCGGQKGFVPQGRSDTLIKPVLV	906
	Sbjct: 116	FCSLATQRTRSSQSVRPKSLSSVSFPVVVPLASGLSLVEVVASVPEFFVKDAVVKTLKV	175
20	Query: 907	KPEGVLVEKTHSSLLCP---KGGKVASESVSLELPVDIVPD-STKAYVTVLGDIMGTALQ	962
	Sbjct: 176	EPEGARKEETVSSLLLPPEHLGGGLEVSEVPALKLPDDVPDTEAEAVISVQGDPVAQAIQ	235
25	Query: 963	N-----LDGLVQMPSGCGEQNMVLFAPIIYVLQYLEKAGLLTE---EIRSRAVGFLSIG	1013
	Sbjct: 236	NTLSGEGLNLLRLPSGCGEQNMIYMAPTVYVLHYLDETQWEKPGTKKKQKAIDLINKG	295
30	Query: 1014	YQKELMYKHSNGSYSAFGERDNGNTWLTAFVTKCFGQAQKFIFIDPKNIQDALKW-MAG	1072
	Sbjct: 296	YQRQLNYRKADGSYAAFLHRA--SSTWLTAFVLKVFSSQARNYVFIDEHICGAVKWLILN	353
35	Query: 1073	NQLPSGCIYANVGNLLHTAMKGGVDD---EVSLTAYVTAALLEMGKDVDPMVSGQLRCL	1128
	Sbjct: 354	QQKDDGVFRESGPVIHNEKGGVGDDAEVEVTLTAFITIALLEAKLVCISPVVANALSIL	413
40	Query: 1129	KNSATSTN-----LYTQALLAYIFSLAGEMDIRNILLKQLDQQAISGESIYWS--QK	1180
	Sbjct: 414	KASDYLLENYANGQVRVYTLALTAYALALAGVLHKLKEILKSLKEELYKALVKGHWERPQK	473
45	Query: 1181	PTPSSNASPWSEPAVDVELTAYALLAQLTKPSLTQKEIAKATSIVAWLAKQHAYGGFS	1240
	Sbjct: 474	PKDAPGHPYSPQPQAAAVEMTSYALLALLT--LLFPKVMAPKVVKWLTEQQYYGGGFG	531
50	Query: 1241	STQDITVALQALAKYATTAYMPSE-EINLVVKSTEN-FQRTFNIQSVNRLVFQDITLP-N	1297
	Sbjct: 532	STQDITVALQALSKYGIATPHKEKNLSVTIQSPSGSFKSHFQILNNAFLLRPVELPLN	591
55	Query: 1298	VPGMYTLEASGQGCYVQTVLRYNLPPTNMKTFSLSVEIGKARCEQPTSPR-SLTLTIH	1356
	Sbjct: 592	EGFTVTAKVTGQGTLTLTVTTRYKVLDDKNTFCFDLKIETVPDTCVEPKGAKNSDYLISIC	651
	Query: 1357	TSYVGSRSSNMAIVEVKMLSGFSPMEGT--NQLLLQQLVKKVEFGTDTLNIYLDDELIC	1414
	Sbjct: 652	TRYAGSRSDSGMAIADISMLTGFIPLKPDLLKLENGVDRYVSKYEIDGNHVLVLYLDKVS	711
	Query: 1415	-NTQTYTFTISQSVLVTNLKPAIKVYDYYLP	1445
	Sbjct: 712	SETECVGFKIHQDFEVGLLQPASVKVYDYYEP	743

**Table 2F Domain Analysis of NOV2**

gnl|Pfam|pfam01835, A2M\_N, Alpha-2-macroglobulin family N-terminal region. This family includes the N-terminal region of the alpha-2-macroglobulin family. (SEQ ID NO:81)  
CD-Length = 620 residues, 98.4% aligned  
Score = 236 bits (603), Expect = 5e-63

Query:	5	LLLGMLALSPAIAEEL--PNYLVTLPARLNFPSVQKVCLDLSPGYSDVKFTVTLETDKDT	62
		+           +   +   +   + +         +     +   +	
5	Sbjct:	2	LLWLLLLLLLLFFDSSLQKPRYMVIVPSILRTETPEKVCVQLHDLNETVTVTVSLHSFPGK
			61
Query:	63	QKLLEYSGLK---KRHLHCISFLVPPPA---GGTEEVATIRVSGVGNNISFEEKKKVLIQ	116
		+   +   +                     + +     +   +     +	
10	Sbjct:	62	RNLSSLFTVLLSSKDLFHCVSFTVPQGLFKSSKGEESFVVVQVKGPTHFTFEKVTVLVS
			121
Query:	117	RQNGTFVQTDKPLYTPGQQVYFRIVTMDSNFVPVNDKYSMVELQDPNSNRIAQWLEVVP	176
		+   +         +           + + + +     +   +   +   +	
	Sbjct:	122	SRRGLVFIQTDKPIYTPGQTVRYRVFSVDENLRPLNELI-LVYIEDPEGNRVDQWEVNKL
			180
15	Query:	177	EQGIVDLSFQLAPEAMLGTYTAV---AEGKTFGT--FSVEEYVLSPLLLLSSVLPKFK
		+   +   +   + + + +         +     +	231
	Sbjct:	181	EGGIFQLSFPIPISEPIQGTWKIVARYESGPESNYTHYFEVKEY-----VLPSFEVS
			231
20	Query:	232	VEVVEPKELSTVQESFLVKICCRYTYGKPMGLGAVQVSVCQKANTYWYREVEREQLPDKCR
		+ +   +               +         + + +	291
	Sbjct:	232	ITPPKPFIIYDNFKEFEVTICARYTYGKPVPGVAYVRFVGVK-----DEDGKKELLAGLE
			285
Query:	292	NLSGQTDKTG--CFSAPVDMATFDLIGYAY-SHQINIVATVVEEGTGVEANA-TQNIYIS	347
		+           +     + +   +	
25	Sbjct:	286	ERAKLLDGNGEICLSQEVLLKELQLKNEDLEGKSLYVAVAVIESEGGDMEEAELGGIKIV
			345
Query:	348	PQMGSMTFEDTSNFYHPNFPFSGKMLLKFPQGGVLPCKNHLVFLVIYGTNGTGNQTLVTD	407
		+     + +       +   +         + + +	
30	Sbjct:	346	RSPYKLFVKTPSHFKPGIPFLLKVLVVDPDGS--PAPNVPVK--VSAQDASYSSNGTTD
			401
Query:	408	NNGLAPFTLETSGWNGTDVSLLEGKFQMEDLVYNPEQVPRYYQNAYLHLRPFYSTTRSFLG	467
		+       + +     + +   + + +   +     +   +	
	Sbjct:	402	EDGLAQFSINTS--GISSLSITVRTNHKELPEEVQAHAEQAATAYSTVSL--SKSYIHLS
			457
35	Query:	468	IHRLNGPLKCGQPQEVLDVYYIDPADASPDQEISFSYYLIGKGLVMEGQKHLNSKKKGL
		+ + + + +     + +     +   + + + +	527
	Sbjct:	458	IER---TLPCGPGVGEQANFILRGKSLGELKILHFFYYLIMSKGKIVKTGRE-----PREPG
			510
40	Query:	528	KASFSLSLTFTSRLAPDPSLVIAIFPSGGVVADKIQFSVEMCFDN-----QQL
		+       +                     +         +	576
	Sbjct:	511	QGLFSLSIPVTPDLAPSFRLVAYYILPQGEVVADSVWIDVEDCCANKLDLSFSPSKDYRL
			570
Query:	577	PGAELVELQLQAAPGSLCALRAVDESLLLLRPDRELSNRSVY	617
		+   + + +             + +     +   +	
45	Sbjct:	571	PAQOVKLRVEADPQSLVALRAVDQAVYLLKPKAKLSMSKVY
			611

The proteinase-binding alpha-macroglobulins (A2M) [1] are large glycoproteins found in the plasma of vertebrates, in the hemolymph of some invertebrates and in reptilian and avian egg white. A2M-like proteins are able to inhibit all four classes of proteinases by a 'trapping' mechanism. They have a peptide stretch, called the 'bait region', which contains specific cleavage sites for different proteinases. When a proteinase cleaves the bait region, a

conformational change is induced in the protein, thus trapping the proteinase. The entrapped enzyme remains active against low molecular weight substrates, whilst its activity toward larger substrates is greatly reduced, due to steric hindrance. Following cleavage in the bait region, a thiol ester bond, formed between the side chains of a cysteine and a glutamine, is  
5 cleaved and mediates the covalent binding of the A2M-like protein to the proteinase.

Alpha2-Macroglobulin (A2M) is a proteinase inhibitor found in association with senile plaques (SP) in Alzheimer's disease (AD). A2M has been implicated biochemically in binding and degradation of the amyloid beta (Abeta) protein which accumulates in SP. We studied the relationship between Alzheimer's disease and a common A2M polymorphism, Val1000  
10 (GTC)/Ile1000 (ATC), which occurs near the thiolester active site of the molecule. In an initial exploratory data set (90 controls and 171 Alzheimer's disease) we noted an increased frequency of the G/G genotype from 0.07 to 0.12. We therefore tested the hypothesis that the G/G genotype is over-represented in Alzheimer's disease in an additional independent data set: a group of 359 controls and 566 Alzheimer's disease patients. In the hypothesis testing cohort,  
15 the G/G genotype increased from 0.07 in controls to 0.12 in Alzheimer's disease ( $P < 0.05$ , Fisher's exact test). The odds ratio for Alzheimer's disease associated with the G/G genotype was 1.77 (1.16-2.70,  $P < 0.01$ ) and in combination with APOE4 was 9.68 (95% CI 3.91-24.0,  $P < 0.001$ ). The presence of the G allele was associated with an increase in Abeta burden in a small series. The A2M receptor, A2M-r/LRP, is a multifunctional receptor whose ligands  
20 include apolipoprotein E and the amyloid precursor protein. These four proteins have each been genetically linked to Alzheimer's disease, suggesting that they may participate in a common disease pathway.

Six alpha 2-macroglobulin cDNA clones were isolated from two liver cDNA libraries produced from rats undergoing acute inflammation. The coding sequence for rat alpha 2-  
25 macroglobulin including its 27-residue signal peptide and the 3' - and part of the 5' nontranslated regions were determined. The mature protein consisting of 1445 amino acids is coded for by a 4790 +/- 40 nucleotide messenger RNA. It contains a typical internal thiol ester region and 25 cysteine residues which are conserved between rat and human alpha 2-macroglobulin. Although the amino acid sequences of rat and human alpha 2-macroglobulin  
30 share 73% identity, two small divergent areas of 17 and 38 residues were found, corresponding to 29 and 11% identity, respectively. These areas are located in the bait region and, therefore, may confer specific proteinase recognition capabilities on rat alpha 2-macroglobulin. Following an inflammatory stimulation, rat alpha 2-macroglobulin mRNA levels increased 214-fold over control values and reached a maximum at 18 h. By 24 h the

levels had decreased to less than 30% of the maximum value. Transcription rates from the alpha 2-macroglobulin gene as measured in nuclear run-on experiments showed a less than 3-fold increase in nuclei from acutely inflamed rats as compared to controls. These results suggest that the accumulation of alpha 2M mRNA is due to the combined effects of increased transcription rates and post-transcriptional processing.

The alpha 2-macroglobulin (alpha 2M) receptor complex as purified by affinity chromatography contains three polypeptides: a 515-kDa heavy chain, an 85-kDa light chain, and a 39-kDa associated protein. Previous studies have established that the 515/85-kDa components are derived from a 600-kDa precursor whose complete sequence has been determined by cDNA cloning (Herz, J., Hamann, U., Rogne, S., Myklebost, O., Gassepohl, H., and Stanley, K. (1988) *EMBO J.* 7,4119-4127). We have now determined the primary structure of the human 39-kDa polypeptide, termed alpha 2M receptor-associated protein, by cDNA cloning. The deduced amino acid sequence contains a putative signal sequence that precedes the 323-residue mature protein. Comparative sequence analysis revealed that alpha 2M receptor-associated protein has 73% identity with a rat protein reported to be a pathogenic domain of Heymann nephritis antigen gp 330 and 77% identity to a mouse heparin-binding protein termed HBP-44. The high overall identity suggests that these molecules are interspecies homologues and indicates that the pathogenic domain, previously thought to be a portion of gp 330, is in fact a distinct protein. Further, the 120-residue carboxyl-terminal region of alpha 2M receptor-associated protein has 26% identity with a region of apolipoprotein E containing the low density lipoprotein receptor binding domain. Pulse-chase experiments revealed that the newly formed alpha 2M receptor-associated protein remains cell-associated, while surface labeling experiments followed by immunoprecipitation suggest that this protein is present on the cell surface forming a complex with the alpha 2M receptor heavy and light chains.

Alpha-2-macroglobulin is a serum pan-protease inhibitor. A possible correlation to this gene is the following reference on Rhinovirus infections causing exacerbations of eosinophilic airway disease. The acute effects of allergen-challenge on nasal interleukin-8 (IL-8), eosinophil cationic protein (ECP), and alpha2-macroglobulin were examined in atopic subjects with common cold symptoms. Twenty-three patients with seasonal allergic rhinitis were inoculated with human rhinovirus 16 outside the pollen season. Diluent and allergen challenges, followed by nasal lavages, were carried out about 3 months before and 4 days after virus inoculation. Seventeen patients developed significant common cold symptoms with increased nasal lavage fluid levels of alpha2-macroglobulin, IL-8, and ECP at baseline

( $p < 0.001$ - $0.05$  versus before inoculation), and were further increased by allergen challenge ( $p < 0.001$ - $0.05$ ); IL-8 and ECP levels were correlated. See *Eur Respir J* 1999 Jan;13(1):41-7.

Allergen challenge-induced acute exudation of IL-8, ECP and alpha2-macroglobulin in human rhinovirus-induced common colds. Greiff L, Andersson M, Svensson C, Linden M, Myint S, Persson CG Dept of Otorhinolaryngology, Head and Neck Surgery, University Hospital, Lund, Sweden.

In the CNS area Alpha-2-macroglobulin has been implicated in Alzheimer disease (AD) based on its ability to mediate the clearance and degradation of A-beta, the major component of amyloid beta deposits. Blacker et al. (1998) analyzed a deletion in the A2M gene at the 5-prime splice site of 'exon II' of the bait region (exon 18) and found that inheritance of the deletion, designated A2M-2, conferred increased risk for AD.

The disclosed NOV2 nucleic acid of the invention encoding a Alpha-2-macroglobulin precursor-like protein includes the nucleic acid whose sequence is provided in Table 2A or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 2A while still encoding a protein that maintains its Alpha-2-macroglobulin precursor-like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 40% percent of the bases may be so changed.

The disclosed NOV2 protein of the invention includes the Alpha-2-macroglobulin precursor-like protein whose sequence is provided in Table 2B. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 2B while still encoding a protein that maintains its Alpha-2-macroglobulin precursor-like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 62% percent of the residues may be so changed.

The NOV2 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in asthma, allergy and psoriasis, Alzheimer disease, Emphysema, pulmonary disease, immune disorders and Cancer and/or other pathologies and disorders. The NOV2 nucleic acid encoding Alpha-2-macroglobulin precursor-like protein, and the Alpha-2-macroglobulin precursor-like protein of the invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

NOV2 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. The disclosed NOV2 protein has multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated NOV2 epitope is from about amino acids 80 to 100. In another embodiment, a NOV2 epitope is from about amino acids 110 to 200. In additional embodiments, NOV2 epitopes are from about amino acids 290 to 340, from about 380 to 400, from about amino acids 410 to 580, from about amino acids 620 to 730, from about amino acids 810 to 900, from about amino acids 990 to 1100, from about amino acids 1190 to 1210, from about amino acids 1240 to 1320, and from about amino acids 1400 to 1450. These novel proteins can be used in assay systems for functional analysis of various human disorders, which are useful in understanding of pathology of the disease and development of new drug targets for various disorders.

### NOV3

A disclosed NOV3 nucleic acid of 987 nucleotides (also referred to as GMAC079237\_A) encoding a novel Ileal Sodium/Bile Acid Cotransporter-like protein is shown in Table 3A. An open reading frame was identified beginning with a ATG initiation codon at nucleotides 1-3 and ending with a TGA codon at nucleotides 979-981. The start and stop codons are in bold letters, and the 3' untranslated region is underlined.

**Table 3A. NOV3 Nucleotide Sequence (SEQ ID NO:11)**

**ATG**AGAGCCAATTGTTCCAGCAGCTCAGCCTGCCCTGCCAACAGTTCAGAGGAGGAGCTGCCAGTGGGACT  
 GGAGGTGCATGGAAACCTGGAGCTCGTTTTTCACAGTGGTGTCCACTATCATGATGGGGCTGCTCATGTTCT  
 CTTTGGGATGTTCCGTGGAGATCCGGAAGCTGTGGTCGCACATCAGGAGACCCCTGGGGCATTGCTGTGGGA  
 CTGCTCTGCCAGTTTGGGCTCATGCCTTTTACAGCTTATCTCCTGGCCATTAGCTTTTCTCTGAAGCCAGT  
 CCAAGCTATTGCTGTTCTCATCATGGGCTGCTGCCGGGGGGCACCATCTTAACATTTTACCTTCTGGGT  
 TGATGGAGATATGGATCTCAGGTGCCCTGGGAATGATGCCACTCTGCATTATCTCTACACCTGGTCTCTGG  
 AGTCTTCAGCAGAATCTCACCATTCTTATCAGAACATAGGTCTGTCTTTAGGAATTACCCTTGTGTGCCT  
 GACCATTCTGTGGCCTTGGTGTCTATGTGAATTACAGATGGCCAAAACAATCCAAAATCATTCTCAAGG  
 CCGTTGTTGGTGGGGTCTCCTTCTGGTGGTGCAGTTGCTGGTGTGGTCTGGCGAAAGGATCTTGAAT  
 TCAGACATCACCTTCTGACCATCAGTTTTCATCTTTCTTGGATTGGCCATGTACACGGGTTTTCTGCTGGC  
 ACTTTTTACCCACCACTTGGCAAAGGACCTTGCCTATCTTTTAGGTTTAGCTTTCAAGACACCTGTG  
 ATACCCTACTCGCAATGACTTCGTGTCTGAAATGTTCCAGGCTCATCTATGCCTTCATTCTCTGTATAT  
 GGACTCTTCAGCTGATAGATGGATTCTTATTGTTGAAGAGAGAACAGAAGATACAGACTGCGATGGTTC  
 ACCTTTACCTGAGTATTTTACTGAGGTAACAATAACCTAAACAACCTAGGATATGACAGCTT

The disclosed NOV3 nucleic acid sequence maps to the q33 region of chromosome 13 and has 257 of 382 bases (67%) identical to a *O.cuniculus* ileal sodium-dependent bile acid transporter mRNA from *O.cuniculus* (GENBANK-ID: Z54357) ( $E = 4.9e^{-34}$ ).

5 A disclosed NOV3 protein (SEQ ID NO:12) encoded by SEQ ID NO:11 has 326 amino acid residues, and is presented using the one-letter code in Table 3B. Signal P, Psort and/or Hydropathy results predict that NOV3 does not have a signal peptide, and is likely to be localized to the plasma membrane with a certainty of 0.6000. In other embodiments NOV3 is also likely to be localized to Golgi body with a certainty of 0.4000, to the mitochondrial inner  
 10 membrane with a certainty of 0.3815, or to the endoplasmic reticulum (membrane) with a certainty of 0.3000.

**Table 3B. Encoded NOV3 protein sequence (SEQ ID NO:12).**

MRANCSSSSACPANSSEELPVGLEVHGNLELVFTVVSTIMMGLLMFSLGCSVEIRKLWSHIRRPWGIAVG  
 LLCQFGLMPFTAYLLAISFSLKPVQAI AVLIMGCCRGAPSLTFSPSGLMEIWI SGALGMMPLCIYLYTWSW  
 SLQQNLTI PYQNI GLSLGITLVCLTIPVAFGVYVNYRWPQSKII LKAVVGGVLLLVAVAGVVLAKGSWN  
 SDITLLTISFIFPLIGHVTGFLALFTHQSWQRTLP IFLGLAFKTPCDTLLAMTSCPECSRLIYAFIPLY  
 GLFQLIDGFLIVEERTEDTDCDGSLPEYFTEVTIIPKQPRI

15 The disclosed NOV3 amino acid has 129 of 331 amino acid residues (38%) identical to, and 198 of 331 amino acid residues (59%) similar to, the 348 amino acid residue ileal sodium/bile acid cotransporter protein from *Cricetulus griseus* (Chinese hamster) (Q60414) ( $E = 1.6e^{-56}$ ).

The NOV3 sequence is predicted to be expressed in ileum because of the expression pattern of (GENBANK-ID: NTCI\_HUMAN) a closely related ILEAL SODIUM/BILE ACID  
 20 COTRANSPORTER homolog in species *Homo sapien*.

TaqMan data for NOV3 can be found below in Example 2. NOV3 also has homology to the amino acid sequences shown in the BLASTP data listed in Table 3E.

**Table 3E. BLAST results for NOV3**

Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 12858115 dbj BAB 31203.1  (AK018423)	putative [Mus musculus]	373	187/310 (60%)	226/310 (72%)	6e-87
gi 3024224 sp Q2872 7 NTCI_RABIT	ILEAL SODIUM/BILE ACID COTRANSPORTE R (ILEAL NA(+)/BILE ACID COTRANSPORTE R) (NA+ DEPENDENT ILEAL BILE ACID TRANSPORTER) (ILEAL SODIUM- DEPENDENT BILE ACID TRANSPORTER) (ISBT) (SODIUM/TAUR OCHOLATE COTRANSPORTI NG POLYPEPTIDE, ILEAL)	347	116/279 (41%)	173/279 (61%)	1e-52
gi 8394281 ref NP_0 58918.1	solute carrier family 10, member 2 [Rattus norvegicus]	348	130/344 (37%)	195/344 (55%)	2e-52
gi 6755530 ref NP_0 35518.1	solute carrier family 10, member 2 [Mus musculus]	348	125/313 (39%)	191/313 (60%)	4e-52
gi 6755530 ref NP_0 35518.1	solute carrier family 10, member 2 [Mus musculus]	348	125/313 (39%)	191/313 (60%)	4e-52



gi 2842631 sp Q60414 NTCI_CRIGR	ILEAL SODIUM/BILE ACID COTRANSPORTE R (ILEAL NA(+)/BILE ACID COTRANSPORTE R) (NA+ DEPENDENT ILEAL BILE ACID TRANSPORTER) (ILEAL SODIUM- DEPENDENT BILE ACID TRANSPORTER) (ISBT) (SODIUM/TAUR OCHOLATE COTRANSPORTI NG POLYPEPTIDE, ILEAL)	348	121/306 (39%)	185/306 (59%)	4e-52
---------------------------------	---	-----	------------------	------------------	-------

The homology of these sequences is shown graphically in the ClustalW analysis shown in Table 3F.

5

**Table 3F. ClustalW Analysis of NOV3**

- 1) NOV3 (SEQ ID NO:12)
- 3) gi|12858115|dbj|BAB31203.1| (AK018423) putative [Mus musculus] (SEQ ID NO:40)
- 4) gi|3024224|sp|Q28727|NTCI\_RABIT ILEAL SODIUM/BILE ACID COTRANSPORTER (ILEAL NA(+)/BILE ACID COTRANSPORTER) (NA+ DEPENDENT ILEAL BILE ACID TRANSPORTER) (ILEAL SODIUM-DEPENDENT BILE ACID TRANSPORTER) (ISBT) (SODIUM/TAUROCHOLATE COTRANSPORTING POLYPEPTIDE, ILEAL) (SEQ ID NO:41)
- 5) gi|8394281|ref|NP\_058918.1| solute carrier family 10, member 2 [Rattus norvegicus] (SEQ ID NO:42)
- 6) gi|6755530|ref|NP\_035518.1| solute carrier family 10, member 2 [Mus musculus] (SEQ ID NO:43)
- 7) gi|6755530|ref|NP\_035518.1| solute carrier family 10, member 2 [Mus musculus] (SEQ ID NO:44)
- 8) gi|2842631|sp|Q60414|NTCI\_CRIGR ILEAL SODIUM/BILE ACID COTRANSPORTER (ILEAL NA(+)/BILE ACID COTRANSPORTER) (NA+ DEPENDENT ILEAL BILE ACID TRANSPORTER) (ILEAL SODIUM-DEPENDENT BILE ACID TRANSPORTER) (ISBT) (SODIUM/TAUROCHOLATE COTRANSPORTING POLYPEPTIDE, ILEAL) (SEQ ID NO:45)

		10	20	30	40
25	NOV3	MRANCS-SSSACPANSS	EEELPVGLEV	HGNLEI	VFTVVS
	gi 12858115	MSTDCA-GNSTCPVNST	EDPPVGMEGF	ANLKLFTVLSA	
	gi 3024224	MSNLTVGCLANATV	CEGASCVAPESN	FNAILSVV	STVLT
	gi 8394281	MDNSSV-CSPNATV	CEGASCVLPESN	FNAILSTV	STVLT
30	gi 6755530	MDNSSV-CPPNATV	CEGASCVLPESN	FNAILSTV	STVLT
	gi 2842631	MDNSSV-CPPNATV	CEGASCVLPESN	FNAILSVV	STVLT
		50	60	70	80
35	NOV3	LLMGLMFS	SGCCEVIRKLWS	HRRPWGI	AVGLCQFGIM
	gi 12858115	VAVGLVMFS	SGCCEVSQKLWL	HRRPWGI	AVGLCQFGIM
	gi 3024224	ILLALVMFS	MGCNVEIKKFLGH	IIRPWGIF	IGFLCQFGIM
	gi 8394281	ILLA VMFS	MGCNVEIKKFLGH	IIRPWGIF	VGFLCQFGIM
	gi 6755530	ILLA VMFS	MGCNVEIKKFLGH	IIRPWGIF	VGFLCQFGIM

gi|2842631| ILLALVMFSMGCNVEGKFLGHRRPWGIVVGFLCQFGIM

90 100 110 120

5 NOV3 PFTAYLLATSFSSKPVQAAVLIWGCCRGL---APSLTSP  
gi|12858115| PLTAYLLATGFGKPFQAAVLIWGSCPGGTISNLTWV  
gi|3024224| PLTGFLVAVFGIPIQAVVLIWGCCPGGTASNILAYW  
gi|8394281| PLTGFLVAVSGIIPVQAVVLIWGCCPGGTASNILAYW  
gi|6755530| PLTGFLVAVSGIIPVQAVVLIWGCCPGGTASNILAYW  
10 gi|2842631| PLTGFLVAVFGIIPVQAVVLIWGCCPGGTASNILAYW

130 140 150 160

15 NOV3 SGGLMFWISG-----ALGMPLCLSYTWSLSQONT  
gi|12858115| DGDMDLSVSMTTCTSTLALGMPLCLSYTKMWVDSGTIV  
gi|3024224| DGDMDLSVSMTTCTSTLALGMPLCLSYTKMWVDSGTIV  
gi|8394281| DGDMDLSVSMTTCTSTLALGMPLCLSYTKMWVDSGTIV  
gi|6755530| DGDMDLSVSMTTCTSTLALGMPLCLSYTKMWVDSGTIV  
20 gi|2842631| DGDMDLSVSMTTCTSTLALGMPLCLSYTKMWVDSGTIV

170 180 190 200

25 NOV3 IPYONIGLSIGILVLVLTIPVAFGVVNNWPQAKIILK  
gi|12858115| IPYQ----SIGILVLVLPVAFGVVNNWPQAKIILK  
gi|3024224| IPYD----SIGISLVALVIPVSIGMVNHNKWPQAKIILK  
gi|8394281| IPYD----SIGISLVALVIPVSIGMVNHNKWPQAKIILK  
gi|6755530| IPYD----SIGISLVALVIPVSIGMVNHNKWPQAKIILK  
30 gi|2842631| IPYD----SIGISLVALVIPVSIGMVNHNKWPQAKIILK

210 220 230 240

35 NOV3 --AVVCGVLLIVVAVAGVVLAKGWNSTITLLTSFIFPP  
gi|12858115| VGIILGGILLIVVAVTGMVLAKGWNSTITLLTSFIFPP  
gi|3024224| VGSIIAGAILIVLIAVGGILYQSAWIIEPKLWIIGTIFP  
gi|8394281| VGSIIAGAILIVLIAVGGILYQSAWIIEPKLWIIGTIFP  
gi|6755530| VGSITGVLLIVLIAVGGILYQSAWIIEPKLWIIGTIFP  
40 gi|2842631| VGSIIAGAILIVLIAVGGILYQSAWIIEPKLWIIGTIFP

250 260 270 280

45 NOV3 IGVVTGFLALPFTQSQW---RTLPFLGLAFKTPCDTLIA  
gi|12858115| VGVVTGFLALPFTQSQRCTSIETCAONIQLCIAMTQ  
gi|3024224| AGYSLGFFFLARAGQPWYRCRTVALETGQNTQLCSTIVQ  
gi|8394281| AGYSLGFFFLARAGQPWYRCRTVALETGQNTQLCSTIVQ  
45 gi|6755530| AGYSLGFFFLARAGQPWYRCRTVALETGQNTQLCSTIVQ  
gi|2842631| AGYCLGFFFLARAGQPWYRCRTVALETGQNTQLCSTIVQ

290 300 310 320

50 NOV3 MTSCPECSRLTYAFIPLLYGTFQLEDGFLIVEE  
gi|12858115| LSFSAEYLVQLLNFA-LAYGLFQVLEHGLLIYAAYQAYRR  
gi|3024224| LSFSPEDLTYVFTFP-LIYTFQIAFAAIFLGIYVAYKRC  
gi|8394281| LSFSPEDLNLVFTFP-LIYTFQIAFAAIFLGIYVAYKRC  
gi|6755530| LSFSPEDLNLVFTFP-LIYTFQIAFAAIFLGIYVAYKRC  
55 gi|2842631| LSFSPEDLNLVFTFP-LIYTFQIAFAAIFLGIYVAYKRC

330 340 350 360

60 NOV3 ---RTEDTDC---DGSP-LPEYFT---EVITIIP  
gi|12858115| QKSKCRQHPPDCPDVCYKQPRETSALDKGDEAAVTLGP  
gi|3024224| EG-KNOAEFPDIKDT-KTEP-ESSFHO-----MNGG  
gi|8394281| EG-KNOAEFLEKTDN-DMDP-MPSFOI-----TNKG  
gi|6755530| EG-KNOAEFLEKTDN-DMDP-MPSFOI-----TNKG  
65 gi|2842631| EG-KNNTELOEKTDN-DMDP-RSSFQI-----TNKG

370 380

70 NOV3 KQPRI-----  
gi|12858115| VQPIQHHRAAELTSHIPSCE  
gi|3024224| FQPI-----

```
gi|8394281|FOPDK-----
gi|6755530|FOPDK-----
gi|2842631|FOPDK-----
```

5

Table 3G list the domain description from DOMAIN analysis results against NOV3. This indicates that the NOV3 sequence has properties similar to those of other proteins known to contain this domain.

**Table 3G Domain Analysis of NOV3**

gnl|Pfam|pfam01758, SBF, Sodium Bile acid symporter family. This family consists of Na<sup>+</sup>/bile acid co-transporters. These transmembrane proteins function in the liver in the uptake of bile acids from portal blood plasma a process mediated by the co-transport of Na<sup>+</sup>. Also in the family is ARC3 from *S. cerevisiae* this is a putative transmembrane protein involved in resistance to arsenic compounds. (SEQ ID NO:82) CD-Length = 186 residues, 80.1% aligned Score = 75.9 bits (185), Expect = 3e-15

10

```
Query: 39 TIMMGLLMFSLGCSVEIRKLWSHIRRPWGIAVGLLCQFGLMPFTAYLLAIS-FSLKPVQA 97
      + + | + | + | | | + | + | + | + | + | + | + |
Sbjct: 1 ALGLFLMMFSGMGLKVRFDLKEALRRPKALILGLLQWIIMPLLMFILAWLLRLPPELA 60
```

15

```
Query: 98 IAVLIMGCCRG---APSLTFSPSGLMEIWIS-----GALGMMPLCIYLYTWSWSLQQN 147
      + + + + | | | + | + | + | + | + | + |
Sbjct: 61 TGLILVGCAPGGAMSNVWTYLAKGDVELSVVMVALSTLLAPLVTPLLSFLLAGLL----- 115
```

20

```
Query: 148 LTIPYQNIGLSLGITLVCLTIPVAFGVVNYRWP 181
      + + + + | | + | + | + | + |
Sbjct: 116 VHVDVSPWSLIKSVLVYVVIPLIAGMLTRYFLP 149
```

Bile acids are synthesized from cholesterol in the liver and secreted into the small intestine, where they facilitate absorption of fat-soluble vitamins and cholesterol. Wong et al. (1996) noted that, rather than being excreted, the majority of bile acids are reabsorbed from the intestine and returned to the liver via the portal circulation. In the liver, bile acids are quantitatively extracted and resecreted into bile, thereby eliminating the need for substantial de novo hepatic bile acid synthesis. The ileum is the major site of active uptake of bile acids from the intestine, where the sodium-gradient-driven transporter has been identified in the ileal enterocyte. Mutations in the transporter or other ileal genes that participate in the transepithelial transport of bile acids are predicted to affect bile acid and cholesterol metabolism significantly. For example, Heubi et al. (1982) described an apparent familial defect in active ileal bile acid transport.

Using homologous sequences from hamster and rat, Wong et al. (1995) cloned a cDNA encoding an ileal sodium/bile acid cotransporter gene (designated ISBT by them). They also isolated a genomic clone for human ISBT. The gene encodes a 348-amino acid polypeptide with 7 predicted transmembrane domains and a predicted molecular mass of 38 kD. The native

human protein has a relative molecular mass of 40 kD on SDS gel electrophoresis due to N-linked glycosylation. Wong et al. (1995) demonstrated a dysfunctional mutation (P290S) in the ileal sodium-bile acid cotransporter gene in the course of cloning the human cDNA.

Wong et al. (1996) mapped the SLC10A2 gene to chromosome 13 by study of a human/rodent cell hybrid mapping panel and refined the localization to 13q33 by fluorescence in situ hybridization. The ileal sodium-bile acid cotransporter gene is clearly distinct from the hepatic sodium-bile acid cotransporter gene (SLC10A1; 182396) which maps to chromosome 14. Lammert et al. (1998) mapped the Slc10a2 gene to mouse chromosome 8 in a region homologous to chromosome 13q33.

Primary bile acid malabsorption (PBAM) is an idiopathic intestinal disorder associated with congenital diarrhea, steatorrhea, interruption of the enterohepatic circulation of bile acids, and reduced plasma cholesterol levels. Oelkers et al. (1997) screened the SLC10A2 gene for PBAM-associated mutations using SSCP analysis. Four polymorphisms were identified and sequenced in a family with congenital PBAM. One allele encoded an A171S missense mutation and a mutated donor splice site for exon 3 (601295.0002). The other allele encoded 2 missense mutations at conserved amino acid positions, L243P and T262M (601295.0001). In transfected COS cells, the L243P, T262M, and double mutant (L243P/T262M) did not affect transporter protein expression or trafficking to the plasma membrane; however, transport of taurocholate and other bile acids was abolished. In contrast, the A171S mutation had no effect on taurocholate uptake. The dysfunctional mutations were not detected in 104 unaffected control subjects, whereas the A171S was present in 28% of that population. The findings of Oelkers et al. (1997) established that SLC10A2 mutations can cause PBAM and underscored the ileal Na(+)/bile acid cotransporter's role in intestinal reclamation of bile acids. Autosomal recessive inheritance had been suggested by earlier studies of PBAM patients and was supported by the findings that the proband's son, who inherited an SLC10A2 allele encoding the L243P and T262M missense mutations, was free of clinical symptoms. The authors stated this was the second reported defect of a Na(+)/solute cotransporter, the first having been SLC5A1 (182380), the cotransporter defective in glucose/galactose malabsorption.

As reviewed by Small (1997), the enterohepatic circulation (EHC) is an in vivo ecologic system for the conservation of bile salts, allowing them to be used over and over for the absorption of fat. The EHC confines the bile salt pool to the liver, bile ducts, gallbladder, small intestine, and portal vein. There is a virtual absence of bile salts in the systemic circulation. A minimum of 6 (known or postulated) molecules (3 in the ileal mucosal cells and 3 analogous molecules in the hepatocytes) constitutes the active players in the EHC (Dawson

and Oelkers, 1995). The polar ileal enterocytes and the hepatocytes each have 3 unique molecules: a receptor that binds bile salts on one surface and translocates them into the cell, a cellular bile salt binding protein that moves them across the cell, and an exit molecule that moves bile salts out of the other side of the cell. In the intestine, ileal sodium/bile salt transporter (ISBT) is present on the brush boarders of the ileum but not the jejunum. It binds bile salts in the gut lumen and transports them across the brush boarder membrane and hands them to the ileal lipid-binding protein (ILBP; 600422) which binds bile acid in the cytoplasm of the cell. ILBP allows the bile salts to move through the cytoplasm to the basolateral membrane of the ileal intestinal epithelial cell, where a sodium-independent organic ion exchange system secretes bile salts into portal capillaries. Bile salts in portal capillaries bind to albumin and flow to the liver. There they are recognized by a transporter with high homology to ISBT, the sodium taurocholate cotransporting polypeptide (SLC10A1). Small et al. (1972) suggested that a genetic defect in the predicted bile acid receptor in the ileum would lead to diarrhea and/or steatorrhea and suggested that bile acid turnover and fecal bile acid excretion be studied in patients with unexplained diarrhea. Hess Thaysen and Pedersen (1976) described several patients who had diarrhea and excessive bile acid loss, without other ileal pathology. Heubi et al. (1979, 1982) reported a case study of a boy who presented 48 hours after birth with severe diarrhea, steatorrhea, and malabsorption. Intestinal absorption of bile acid was nearly absent and resulted in a small pool size, a low interluminal bile acid concentration, and severe malabsorption of water and fat. Ileal biopsies had no active bile acid transport. Parenteral nutrition was necessary to sustain the child. At the other extreme, a child with marked bile acid malabsorption but with almost normal development, nearly normal fat absorption, and a moderately well-maintained bile acid pool, was described by Jonas et al. (1986). This patient had a 15-fold increase in bile acid synthesis that was adequate to maintain pool size, interluminal bile acids, and fat absorption. Thus, the clinical phenotype apparently can vary from severe diarrhea, fat malabsorption, and malnutrition, to modest diarrhea without significant fat malabsorption. Small (1997) suggested that the bile acid malabsorption and the variable severity could represent mutations in any of the 3 main players in the ileal transport.

The disclosed NOV3 nucleic acid of the invention encoding a Ileal Sodium/Bile Acid Cotransporter-like protein includes the nucleic acid whose sequence is provided in Table 3A or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 3A while still encoding a protein that maintains its Ileal Sodium/Bile Acid Cotransporter-like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes

nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 33% percent of the bases may be so changed.

The disclosed NOV3 protein of the invention includes the Ileal Sodium/Bile Acid Cotransporter-like protein whose sequence is provided in Table 3B or 3D. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 3B or 3D while still encoding a protein that maintains its Ileal Sodium/Bile Acid Cotransporter-like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 61% percent of the residues may be so changed.

The protein similarity information, expression pattern, and map location for the Ileal Sodium/Bile Acid Cotransporter-like protein and nucleic acid (NOV3) disclosed herein suggest that NOV3 may have important structural and/or physiological functions characteristic of the citron kinase-like family. Therefore, the NOV3 nucleic acids and proteins of the invention are useful in potential diagnostic and therapeutic applications. These include serving as a specific or selective nucleic acid or protein diagnostic and/or prognostic marker, wherein the presence or amount of the nucleic acid or the protein are to be assessed, as well as potential therapeutic applications such as the following: (i) a protein therapeutic, (ii) a small molecule drug target, (iii) an antibody target (therapeutic, diagnostic, drug targeting/cytotoxic antibody), (iv) a nucleic acid useful in gene therapy (gene delivery/gene ablation), and (v) a composition promoting tissue regeneration in vitro and in vivo.

The NOV3 nucleic acids and proteins of the invention are useful in potential diagnostic and therapeutic applications implicated in various diseases and disorders described below. For example, the compositions of the present invention will have efficacy for treatment of patients suffering from Primary bile acid malabsorption (PBAM, an idiopathic intestinal disorder), congenital diarrhea, steatorrhea, refractory infantile diarrhea, interruption of the enterohepatic circulation of bile acids, reduced plasma cholesterol levels, crohn's disease, Inflammatory

bowel disease, Diverticular disease, Hirschsprung's disease, Cirrhosis, Transplantation, Hypercalcaemia, Ulcers, growth failure and/or other pathologies. The NOV3 nucleic acid, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

5           NOV3 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. For example the disclosed NOV3 protein have multiple hydrophilic regions,  
10 each of which can be used as an immunogen. In one embodiment, contemplated NOV3 epitope is from about amino acids 5 to 30. In another embodiment, a NOV3 epitope is from about amino acids 55 to 60. In additional embodiments, NOV3 epitopes are from about amino acids 140 to 150, from about amino acids 180 to 190, and from about amino acids 280 to 330. This novel protein also has value in development of powerful assay system for functional  
15 analysis of various human disorders, which will help in understanding of pathology of the disease and development of new drug targets for various disorders.

#### NOV4

20           A disclosed NOV4 nucleic acid of 850 nucleotides (designated CuraGen Acc. No. AL161453\_A) encoding a novel Prohibitin -like protein is shown in Table 4A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 6-8 and ending with a TGA codon at nucleotides 822-824. A putative untranslated regions upstream of the initiation codon and downstream from the termination codon are underlined in Table 4A, and the start and stop codons are in bold letters.

25

**Table 4A. NOV4 Nucleotide Sequence (SEQ ID NO:13)**

AAAACATGGCTGCCAAAATGTTTGTAGTTCATCGGCAAGTTTGGCCTGGCCTTAGTTGTTGCAG  
GAGGCGTGGTGAACCTCTGCCTTATATAGTGTGGATGCTGGGCACAGAGCTGTCGTCTTTGACC  
GATTCCGTGGAGTGCAGGACATTGTGGTAGGCAAAGGGACTCACTGTCTCATCCCATGGTTAC  
AGAAATCAATAATCTTTGACTGCCGTTCTCAGCCACGTAATGTGCCAGTCATCACCGGTAGCA  
AAGATTTACAGAATGTCAACCTCACACTGCGCATCATCTCCGGCCCGTAGCTAGCCAGCTTC  
CTCACATCTTCACCAGCAGCGGAGAGGACCATGATGAGCGTGTGCCGCCGTCCATCAGAACAGA  
TCCTCAAGTCAGTGGTGGCTCGCTTGAAGCTGGAGAACTAATCACCAGAGAGAGCAGATCTCCAG  
GCAGGTGAGCGATGACCTTACGGAGCCAGCAGCCACCTTTGGGCTCATTCTGGACGACGTGTC  
CTTGACATATCTGACCTTCGGGAAGGAGTTCATAGAAGCGGTGGAAGCCAAACAGATAGCTCA  
GCAGGAAGCAGAGAGGGCCAGATTTGTGGTGGAAAAGGCTGAGCAGCAGAAAAAGGCGGCCAT  
CATTTCTGCTGAGGGCGACTCCAAGGTGGCCGAGCTGATCACCAACTCACTGGCCACAGCAGG  
GGACGCCCTGATCGAGCTGCGCAAGCTGGAAGCTGTGGAGGACATCACCTACCAGCTTTTACG  
CTCTCGGAACATCACCTACCTGCGGGCAGGCGAGTCCATGCCCTGCAGCTGCGCTGGTGAGG  
GCCCCACCCTGCCTGCACCTCCGAG

The nucleic acid sequence of NOV4, localized on chromosome 9, has 782 of 849 bases (92%) identical to a Prohibitin mRNA from *Homo sapiens* (GENBANK-ID: S85655) ( $E = 7.2e^{-158}$ ).

5 A NOV4 polypeptide (SEQ ID NO:14) encoded by SEQ ID NO:13 is 272 amino acid residues and is presented using the one letter code in Table 4B. Signal P, Psort and/or Hydropathy results predict that NOV4 has a signal peptide and is likely to be localized at the endoplasmic reticulum (membrane) with a certainty of 0.5500. In other embodiments, NOV4 may also be localized to the lysosome (lumen) with a certainty of 0.2631, the endoplasmic  
10 reticulum (lumen) with a certainty of 0.1000, or extracellularly with a certainty of 0.1000. The most likely cleavage site for a NOV4 peptide is between amino acids 25 and 26, at: VNS-AL.

**Table 4B. NOV4 protein sequence (SEQ ID NO:14)**

MAAKMFEFIGKFGGLLVVAGGVVNSALYSVDAGHRAVVFDRFRGVQDIVVGKGTCLIPWLQKSIIF  
DCRSQPRNPVITGSKDLQNVNLTLLRIIFRPVASQLPHIFTSSGEDHDERVPPSITNKILKSVVARF  
EAGELITQREQISRQVSDDLTEPAATFGLILDVSLTYLTFGKEFIEAVEAKQIAQQAERARFVVE  
KAEQQKKAIIISAEGDSKVAELITNSLATAGDALIELRKLEAVEDITYQLLSRNITYLRAQSMPL  
QLRW

15 The full amino acid sequence of the protein of the invention was found to have 236 of 270 amino acid residues (87%) identical to, and 251 of 270 amino acid residues (92%) similar to, the 272 amino acid residue Prohibitin protein from *Homo sapiens* ACC: P35232) ( $E = 3.0e^{-117}$ ).

The disclosed NOV4 protein is widely expressed in a variety of tissues.

20 TaqMan data for NOV4 can be found below in Example 2. NOV4 also has homology to the amino acid sequences shown in the BLASTP data listed in Table 4C.



Table 4C. BLAST results for NOV4

Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 4505773 ref NP_002625.1	prohibitin [Homo sapiens]	272	236/270 (87%)	251/270 (92%)	e-123
gi 92643 pir  A39682	prohibitin - rat	272	235/270 (87%)	251/270 (92%)	e-123
gi 7298546 gb AAF53765.1  (AE003661)	l(2)37Cc gene product [Drosophila melanogaster]	276	178/270 (65%)	220/270 (80%)	8e-96
gi 2055454 gb AAB53231.1  (U97204)	prohibitin-like molecule TC-PRO-1 [Toxocara canis]	274	153/264 (57%)	209/264 (78%)	9e-80
gi 13491275 gb AAK27865.1  (AC087079)	Hypothetical protein Y37E3.9 [Caenorhabditis elegans]	275	154/270 (57%)	210/270 (77%)	6e-79

The homology of these sequences is shown graphically in the ClustalW analysis shown in Table 4D.

Table 4D ClustalW Analysis of NOV4

- 5 1) NOV4 (SEQ ID NO:14)  
 2) gi|4505773|ref|NP\_002625.1| prohibitin [Homo sapiens] (SEQ ID NO:46)  
 3) gi|92643|pir||A39682 prohibitin - rat (SEQ ID NO:47)  
 4) gi|7298546|gb|AAF53765.1| (AE003661) l(2)37Cc gene product [Drosophila melanogaster] (SEQ ID NO:48)  
 10 5) gi|2055454|gb|AAB53231.1| (U97204) prohibitin-like molecule TC-PRO-1 [Toxocara canis] (SEQ ID NO:49)  
 6) gi|13491275|gb|AAK27865.1| (AC087079) Hypothetical protein Y37E3.9 [Caenorhabditis elegans] (SEQ ID NO:50)

15		10	20	30	40
	NOV4	---MAAKPEFFIGFGLALVAGGVVNSALYSVDAGHRAV			
	gi 4505773	---MAAKVESIGFGLALVAGGVVNSALYNVDAGHRAV			
	gi 6679299	---MAAKVESIGFGLALVAGGVVNSALYNVDAGHRAV			
20	gi 7298546	---MAAFENRIGMGLCAVLGGVVNSALYNVCGHRAV			
	gi 2055454	-MAGAOKLGRIGGICGLAVTGGVVNSALYNVDGGORAV			
	gi 13491275	MAASAOKLGRIGTVGGLGLAGGIAQTALYNVDGGORAV			
25		50	60	70	80
	NOV4	VFDRFRGVQDIIVVGEGTHCLIPWQKPIIFDCRS*PRNV			
	gi 4505773	IFDRFRGVQDIIVVGEGTHFLIPWVQKPIIFDCRS*PRNV			
	gi 6679299	IFDRFRGVQDIIVVGEGTHFLIPWVQKPIIFDCRS*PRNV			
	gi 7298546	IFDRFTGLINNVVGEGTHFFIPWVQKPIIFDIRS*PRNV			
30	gi 2055454	IFDRFTGVIPDVVGEGTHFLIPWVQKPIIFDIRSTPRAIS			
	gi 13491275	IFDRFSGVIEVVVGEGTHFLIPWVQKPIIFDIRSTPRAVT			
35		90	100	110	120
	NOV4	VITGSKDLQNVNITLRILRPVASQLPRIITSSG*DYDER			
	gi 4505773	VITGSKDLQNVNITLRILRPVASQLPRIITSSG*DYDER			
	gi 6679299	VITGSKDLQNVNITLRILRPVASQLPRIITSSG*DYDER			
	gi 7298546	VITGSKDLQNVNITLRILRPVDPQLPRIITITG*DYDER			
	gi 2055454	VITGSKDLQNVNITLRILRPVDPQLPRIITITG*DYDER			
40	gi 13491275	VITGSKDLQNVNITLRILRPVDPQLPRIITITG*DYDER			
45		130	140	150	160
	NOV4	VLPSITN*ILKSVVAF*FAGELITOREQISROVSDDLTEP			
	gi 4505773	VLPSITN*ILKSVVAF*FAGELITOREQISROVSDDLTER			

```

gi|6679299| VLPSTITNE LK VVA FDA GELITQRE V S Q V S D D L T E R
gi|7298546| VLPSTIAPE LK VVA FDA GELITQRE V S Q V S Q L T V R
gi|2055454| VLPSTITNE LK VVA FDA HE ITQRESVSH VSV L S E R
gi|13491275| VLPSTITNE LK VVA FDA HE ITQRE V S Q A S V A L R E R
5
170 180 190 200
NOV4 AATFGLILDD SLT LTFG E F I E A V E A K Q I A Q Q E A E A R
gi|4505773| AATFGLILDD SLT LTFG E F T E A V E A K Q V A Q Q E A E A R
10 gi|6679299| AATFGLILDD SLT LTFG E F T E A V E A K Q V A Q Q E A E A R
gi|7298546| A A Q F G F I L D D S L T H L T F G E F T L A V E A K Q V A Q Q E A E A R
gi|2055454| A R C F G I L D D S L T H L S F G E F T E A V E A K Q V A Q Q E A E A R
gi|13491275| A A Q F G L L D D S L T H L N F G E F T E A V E A K Q V A Q Q E A E A R
15
210 220 230 240
NOV4 FVVEKAEQQK K A A I I S A E G D S I A A L I N S L A A G D A L I E
gi|4505773| FVVEKAEQQK K A A I I S A E G D S I A A L I A N S L A A G D G L I E
20 gi|6679299| FVVEKAEQQK K A A I I S A E G D S I A A L I A N S L A A G D G L I E
gi|7298546| FVVEKAEQQK L A S I I S A E G D S I A A C L A K S F G E A G D G L I E
gi|2055454| F V E I A E Q M K I A A I T A E G D A C A A L L A Q A F K D A G D G L I E
gi|13491275| FVVEKAEQ M K I A A I T A E G D A C A A L L A K A F A S A G D G L I E
25
250 260 270
NOV4 L R K I E A V E D I I Y Q L R S R N I T Y L R A G S M P L Q L R W ---
gi|4505773| L R K I E A A E D I A Y Q L S R S R N I T Y L P A G S V L L Q L P Q ---
30 gi|6679299| L R K I E A A E D I A Y Q L S R S R N I T Y L P A G S V L L Q L P Q ---
gi|7298546| L R K I E A A E D I A Y Q L S R S R G V A Y L P S G S T L L N L P S T I A Q
gi|2055454| L R K I E A A E I A E R A S T A R N V I Y L P G N Q T L F N L P A ---
gi|13491275| L R K I E A A E I A E R A L A T N V I Y L P G N Q T L L N L Q S ---

```

Tables 4E-F list the domain description from DOMAIN analysis results against NOV4. This indicates that the NOV4 sequence has properties similar to those of other proteins known to contain this domain.

**Table 4E. Domain Analysis of NOV4**

gnl|Pfam|pfam01145, Band\_7, SPFH domain / Band 7 family. This family also includes proteins with high blast scores to known Band 7 protein: HflC from E. coli HflK from E. coli, and Prohibitin family members (SEQ ID NO:83)  
 CD-Length = 191 residues, 91.6% aligned  
 Score = 157 bits (397), Expect = 7e-40

```

40 Query: 28 YSVDAGHRVVFDRFRGVQDIVVGKGTCHLIPWLQKSIIFDCRSQPRNVVPVITGSKDLQN 87
      | | + | | + | | | | + | | + | | + | |
Sbjct 17 KIVKEYERGVI FRLGRVVRQ-VVGPGLHFIIPFIDTVK FVDLRTVVYDVPSQEII TKDNV 75

Query 88 VNLTLR IIFRPVASQLPHIFTSSGEDHDERVPPSITNKILKSVVARFEAGELITQREQIS 147
      | + +++ | | ++ | | + | ++ + + | ++ | + | +
45 Sbjct 76 VVI VDAVVY YRVVDPLKAVYEVED---AERALPQLAQTTLRN VIGQFTLDEILTERERIN 132

Query 148 RQVSDDLTEPAATFGLILDDVSLTYLTFGKEFIEAVEAFQIAQQEAEARARFVVEKAEQQK 207
      | + + | | + + + + | + + + | | + + + + | +
50 Sbjct: 133 SQLREILDEATDPWG I KVERVEIKDIRLP EEVQRAMAAQMEAEREA-RAKILEAEGEQEA 191

```

**Table 4F. Domain Analysis of NOV4**

gnl|Smart|smart00244, PHB, prohibitin homologues; prohibitin homologues (SEQ ID NO:84)  
CD-Length = 160 residues, 98.8% aligned  
Score = 97.4 bits (241), Expect = 9e-22

5 Query: 28 YSV DAGHRA VV FDR FRGV QDI VVG KGT HCL IPWL QKS IIFDCRSQPRNPVITG-SKDLQ 86  
+ | | + | | + | | + | | + | | + | | + | |  
Sbjct: 3 FYVIGEGERG VVERLGRVLK-VLGPGLHFVIPFID DVKRVDLRAQTDDVPPQEVITKDNV 61

10 Query: 87 NVNLT LRIIFRPV ASQLPHIFTSSGEDHDERVPPSITNKILKSVVARFEAGELIT-QREQ 145  
| ++ + + | | | ++ | | | + | | + + | | + | | +  
Sbjct: 62 TVSVDAVVYR-VLDPLKAVYGV--LDADYRALRQLAQTTLRSVIGKRTLDELLTDEREK 118

10 Query: 146 ISRQVSDDLTEPAATFGLILDDVSLTYLTFGKEFIEAVEAKQ 187  
| | + ++ | | | + | + ++ | | + + + | | | + |  
Sbjct: 119 ISENIREELNEAAEPWGIEVEDVEIKDIRLPEEIKEAMEAQQ 160

15 Genes that negatively regulate proliferation inside the cell are of considerable interest because of the implications in processes such as development and cancer. Prohibitin, a novel cytoplasmic anti- proliferative protein widely expressed in a variety of tissues, inhibits DNA synthesis. Studies have suggested that Prohibitin may be a suppressor gene and is associated with tumor development and/or progression of at least some breast cancers. Sequence  
20 comparisons suggest that the Prohibitin gene is an analogue of Cc, a Drosophila gene that is vital for normal development.

Prohibitin is a 30-kD intracellular, antiproliferative protein. White et al. (1991) mapped the gene to chromosome 17 by analysis of human-mouse somatic cell hybrid cell lines using a genomic fragment of human Prohibitin DNA isolated from a library using the rat Prohibitin  
25 cDNA clone. By a study of cell lines containing portions of human chromosome 17, they determined that the PHB gene was located in the 17q11.2-q23 region. By in situ hybridization, they localized the gene to 17q21. Sato et al. (1992) isolated the human homolog of the rat Prohibitin gene and mapped it to 17q12-q21 by in situ hybridization. Sato et al. (1993) showed that the human Prohibitin gene family consists of 1 functional PHB gene on 17q21 and 4  
30 processed pseudogenes, each on a different chromosome: PHBP1 on 6q25, PHBP2 on 11p11.2, PHBP3 on 1p31.3, and PHBP4 on 2q21.

By DNA sequence analysis of 2 exons in the PHB gene in 23 sporadic breast cancers that showed loss of heterozygosity on 17q or developed in patients 35 years old or younger, they identified 4 cases of somatic mutation: 2 of these were missense mutations, 1 showed a 2-  
35 bp deletion resulting in truncation of the gene product due to frameshift, and the fourth had a

C-to-T transition in an intron adjacent to an intron-exon boundary. Sato et al. (1993) found no mutations in the PHB gene in other forms of tumors, namely, those of ovary, liver, and lung.

The disclosed NOV4 nucleic acid of the invention encoding a Prohibitin -like protein includes the nucleic acid whose sequence is provided in Table 4A or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 4A while still encoding a protein that maintains its Prohibitin -like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 10% percent of the bases may be so changed.

The disclosed NOV4 protein of the invention includes the Prohibitin -like protein whose sequence is provided in Table 4B. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 4B while still encoding a protein that maintains its Prohibitin -like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 43% percent of the residues may be so changed.

The protein similarity information, expression pattern, and map location for the Prohibitin-like protein and nucleic acid (NOV4) disclosed herein suggest that this NOV4 protein may have important structural and/or physiological functions characteristic of the Prohibitin family. Therefore, the NOV4 nucleic acids and proteins of the invention are useful in potential diagnostic and therapeutic applications. These include serving as a specific or selective nucleic acid or protein diagnostic and/or prognostic marker, wherein the presence or amount of the nucleic acid or the protein are to be assessed, as well as potential therapeutic applications such as the following: (i) a protein therapeutic, (ii) a small molecule drug target, (iii) an antibody target (therapeutic, diagnostic, drug targeting/cytotoxic antibody), (iv) a nucleic acid useful in gene therapy (gene delivery/gene ablation), and (v) a composition promoting tissue regeneration in vitro and in vivo.

The NOV4 nucleic acids and proteins of the invention are useful in potential diagnostic and therapeutic applications implicated in various diseases and disorders described below. For example, the compositions of the present invention will have efficacy for treatment of patients suffering from breast cancer (In a sporadic breast cancer, Sato et al. (1992) found a missense mutation from valine (GTC) to alanine (GCC) at codon 88 of the PHB gene), and/or other pathologies. The NOV4 nucleic acids, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

NOV4 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. For example, the disclosed NOV4 protein has multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated NOV4 epitope is from about amino acids 40 to 45. In another embodiment, a NOV4 epitope is from about amino acids 60 to 75. In additional embodiments, NOV4 epitopes are from about amino acids 100 to 130, from about amino acids 140 to 160,, from about amino acids 180 to 220, and from about amino acids 240 to 260. These novel proteins can be used in assay systems for functional analysis of various human disorders, which will help in understanding of pathology of the disease and development of new drug targets for various disorders.

### NOV5

A disclosed NOV5 nucleic acid of 2011 nucleotides (also referred to as dj1182a14\_da1) encoding a novel Macrophage Stimulating Protein Precursor -like protein is shown in Table 5A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 1-3 and ending with a TAG codon at nucleotides 1999-2001. A putative untranslated region downstream from the termination codon is underlined in Table 5A, and the start and stop codons are in bold letters.

<b>Table 5A. NOV5 Nucleotide Sequence (SEQ ID NO:15)</b>
--

[illegible]

5

10

**Table 5B. Encoded NOV5 protein sequence (SEQ ID NO:16)**

<p>MGWLPLLLLLTQCLGVPGQRSPLNDFQVLRGTELQHLLHAVVPGPWQEDVADAEECAGRCGPLMDCRAFHYN  VSSHGCGQLLPWTQHSPHTRLRRSGRCDLFQKKDYVRTCIMNNGVG YRGTMATTVGGLPCQAWSHKFPNDHKY  TPTLRNGLEENFCRNP DGP GGPWPCYTTPDAVRFQSCGIKSCREAA C VWCNGEYRGAVDRTESGRECQRWD  LQHPHQHPFEPGKFLDQGLDDNYCRNPDGSERPWCYTTPDQIEREFCDLPRCGSEAQPRQEATTVSCFRGKG  EGYRGRTANTTTAGVPCQRWDAQIPHQHRFTPEKYACKDLRENFCRNPDGSEAPWCFTLRPGMRAAFICYQIRR  CTDDVRPQTATTAQGSSTAARSARPARVSSASAGPLRRRTSRSSRLPNRMHNWRRTSAGTQMGIAMGPAT  RWTQGP HSTTTPCDAALMTRSHQSWTPQTRCSLRSVARGWIGWISGVPSCAWLGAIRATHPGQSACGIGMLP</p>
---

LTGYEVWLGTLFQNPQHGEPSLQRPVPAKMCVCGPSGSQLVLLKLEKSVTLNQRVALICLPPEWYVVPPTKCEIAGWGETKGTGNDTVLNVALLNVISNQECNIKHRGRGDYGGPLACFTHNCWVLEGIIPNRVCARSCWPAVFTRVSVFVDWIHKVMRLG

The disclosed NOV5 amino acid sequence has 368 of 368 amino acid residues (100%) identical to, and 368 of 368 amino acid residues (100%) similar to, the 711 amino acid residue ptnr:SPTREMBL-ACC:Q14870 protein from *Homo sapiens* (Human) (Macrophage-Stimulating Protein Precursor ( $E = 9.9e^{-310}$ )).

NOV5 is expressed in at least HepG2 (liver) In addition, the sequence is predicted to be expressed in the following tissues because of the expression pattern of (GENBANK-ID: HUMMST1A|acc:L11924) a closely related {Homo sapiens macrophage-stimulating protein (MST1) mRNA, complete cds homolog in species Homo sapiens : HepG2, and liver.

TaqMan data for NOV5 can be found below in Example 2. NOV5 also has homology to the amino acid sequences shown in the BLASTP data listed in Table 5C.

Table 5C. BLAST results for NOV5					
Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 15294659 ref XP_054070.1	macrophage stimulating 1 (hepatocyte growth factor-like) [Homo sapiens]	711	561/720 (77%)	577/720 (79%)	0.0
gi 10337615 ref NP_066278.1	macrophage stimulating 1 (hepatocyte growth factor-like) [Homo sapiens]	711	560/720 (77%)	576/720 (79%)	0.0
gi 123114 sp P26927 HGFL_HUMAN	HEPATOCYTE GROWTH FACTOR-LIKE PROTEIN PRECURSOR (MACROPHAGE STIMULATORY PROTEIN) (MSP) (MACROPHAGE STIMULATING PROTEIN)	711	560/720 (77%)	576/720 (79%)	0.0
gi 15299258 ref XP_017811.2	hypothetical protein XP_017811 [Homo sapiens]	529	440/532 (82%)	456/532 (85%)	0.0
gi 90615 pir  A40332	macrophage-stimulating protein 1 precursor - mouse	716	449/725 (61%)	507/725 (69%)	0.0

The homology of these sequences is shown graphically in the ClustalW analysis shown in Table 5D.

Table 5D Clustal W Sequence Alignment

- 1) NOV5 (SEQ ID NO:16)  
 2) gi|15294659|ref|XP\_054070.1| macrophage stimulating 1 (hepatocyte growth factor-like) [Homo sapiens] (SEQ ID NO:51)  
 3) gi|10337615|ref|NP\_066278.1| macrophage stimulating 1 (hepatocyte growth factor-like) [Homo sapiens] (SEQ ID NO:52)  
 4) gi|123114|sp|P26927|HGFL\_HUMAN HEPATOCYTE GROWTH FACTOR-LIKE PROTEIN PRECURSOR (MACROPHAGE STIMULATORY PROTEIN) (MSP) (MACROPHAGE STIMULATING PROTEIN) (SEQ ID NO:53)  
 5) gi|15299258|ref|XP\_017811.2| hypothetical protein XP\_017811 [Homo sapiens] (SEQ ID NO:54)  
 6) gi|90615|pir||A40332 macrophage-stimulating protein 1 precursor - mouse (SEQ ID NO:55)

		10	20	30	40
NOV5		.....	.....	.....	.....
gi 15294659	MGWLP	LLLLLTQCLGV	PGQ	RSP	LNDFQVLRGTELQHLLHA
gi 10337615	MGWLP	LLLLLTQCLGV	PGQ	RSP	LNDFQVLRGTELQHLLHA
gi 123114	MGWLP	LLLLLTQCLGV	PGQ	RSP	LNDFQVLRGTELQHLLHA
gi 15299258	-----	-----	-----	-----	-----
gi 90615	MGWLP	LLLLLVQCSR	ALG	Q	RSP
		50	60	70	80
NOV5		.....	.....	.....	.....
gi 15294659	VVPGPWQEDV	ADAEECAGRC	GPLMD	CR	AFHYNVSSHGCQL
gi 10337615	VVPGPWQEDV	ADAEECAGRC	GPLMD	CR	AFHYNVSSHGCQL
gi 123114	VVPGPWQEDV	ADAEECAGRC	GPLMD	CR	AFHYNVSSHGCQL
gi 15299258	-----	-----	-----	-----	-----
gi 90615	AVPGPWQEDV	ADAEECAR	RCGPL	DC	RAHYNVSSHGCQL
		90	100	110	120
NOV5		.....	.....	.....	.....
gi 15294659	LPWTQHS	PHTLR	RRSGRCDL	FQKKDY	VRTCIMNNGVGYRG
gi 10337615	LPWTQHS	PHTLR	RRSGRCDL	FQKKDY	VRTCIMNNGVGYRG
gi 123114	LPWTQHS	PHTLR	RRSGRCDL	FQKKDY	VRTCIMNNGVGYRG
gi 15299258	LPWTQHS	PHTLR	RRSGRCDL	FQKKDY	VRTCIMNNGVGYRG
gi 90615	LPWTQHS	PHT	LYHS	SLCH	LFQKKDYVRTCIMNGVSYRG
		130	140	150	160
NOV5		.....	.....	.....	.....
gi 15294659	TMATTVGGLP	CQAWSHKFP	NDHKYTP	TLR	NGLEENFCRNP
gi 10337615	TMATTVGGLP	CQAWSHKFP	NDHKYTP	TLR	NGLEENFCRNP
gi 123114	TMATTVGGLP	CQAWSHKFP	NDHKYTP	TLR	NGLEENFCRNP
gi 15299258	TMATTVGGLP	CQAWSHKFP	NDHKYTP	TLR	NGLEENFCRNP
gi 90615	TMATTAGGLP	CQAWSR	FPNDHKYTP	TLR	NGLEENFCRNP
		170	180	190	200
NOV5		.....	.....	.....	.....
gi 15294659	DGDPGGPW	CYTTDPA	VR	FQSCGIK	SCREAA
gi 10337615	DGDPGGPW	CYTTDPA	VR	FQSCGIK	SCREAA
gi 123114	DGDPGGPW	CYTTDPA	VR	FQSCGIK	SCREAA
gi 15299258	DGDPGGPW	CYTTDPA	VR	FQSCGIK	SCREAA
gi 90615	DGDPGGPW	CYTTDPA	VR	FQSCGIK	SCREAA
		210	220	230	240
NOV5		.....	.....	.....	.....
gi 15294659	GAVDRTES	GRE	QRWDLQ	HPHQHP	FE
gi 10337615	GAVDRTES	GRE	QRWDLQ	HPHQHP	FE
gi 123114	GAVDRTES	GRE	QRWDLQ	HPHQHP	FE
gi 15299258	GAVDRTES	GRE	QRWDLQ	HPHQHP	FE
gi 90615	GAVDRTES	GRE	QRWDLQ	HPHQHP	FE



		250	260	270	280
5	NOV5	RNPDGSERPWCYTTDPQIEREFCDLPRCG-----SE			
	gi   15294659	RNPDGSERPWCYTTDPQIEREFCDLPRCG-----SE			
	gi   10337615	RNPDGSERPWCYTTDPQIEREFCDLPRCG-----SE			
	gi   123114	RNPDGSERPWCYTTDPQIEREFCDLPRCG-----SE			
	gi   15299258	RNPDGSERPWCYTTDPQIEREFCDLPRCG-----SE			
	gi   90615	RNPDGSERPWCYTTDPQIEREFCDLPSCCPNLPPTVKGS			
10	NOV5	290	300	310	320
	gi   15294659	AQPRQEATTVSCFRGKGEGYRGNTTTAGVPCQRWDAQI			
	gi   10337615	AQPRQEATTVSCFRGKGEGYRGNTTTAGVPCQRWDAQI			
15	gi   123114	AQPRQEATTVSCFRGKGEGYRGNTTTAGVPCQRWDAQI			
	gi   15299258	AQPRQEATTVSCFRGKGEGYRGNTTTAGVPCQRWDAQI			
	gi   90615	AQPRQEATTVSCFRGKGEGYRGNTTTAGVPCQRWDAQS			
20	NOV5	330	340	350	360
	gi   15294659	PHQHRFTPEKYACKDLRENFCRNPDGSEAPWCFTLRPGMR			
	gi   10337615	PHQHRFTPEKYACKDLRENFCRNPDGSEAPWCFTLRPGMR			
	gi   123114	PHQHRFTPEKYACKDLRENFCRNPDGSEAPWCFTLRPGMR			
25	gi   15299258	PHQHRFTPEKYACKDLRENFCRNPDGSEAPWCFTLRPGMR			
	gi   90615	PHQHRFTPEKYACKDLRENFCRNPDGSEAPWCFTLRPGMR			
30	NOV5	370	380	390	400
	gi   15294659	AAFCYQIRRCTDDVRPQDCTAQAQSSTAARSIPAIVSSA			
	gi   10337615	AAFCYQIRRCTDDVRPQDCYHGAGEQYRGTVSKTRKGVQC			
	gi   123114	AAFCYQIRRCTDDVRPQDCYHGAGEQYRGTVSKTRKGVQC			
	gi   15299258	AAFCYQIRRCTDDVRPQDCYHGAGEQYRGTVSKTRKGVQC			
35	gi   90615	AAFCYQIRRCTDDVRPQDCYHGAGEQYRGTVSKTRKGVQC			
40	NOV5	410	420	430	440
	gi   15294659	SAGPLRRRTSRSSRLPPNRMHNWRTSAGTQMGIIMGPGA			
	gi   10337615	QRWSAETPHKPQFTFTSEPHAQLEENFCRNPDGDSHGPP--			
	gi   123114	QRWSAETPHKPQFTFTSEPHAQLEENFCRNPDGDSHGPP--			
	gi   15299258	QRWSAETPHKPQFTFTSEPHAQLEENFCRNPDGDSHGPP--			
45	gi   90615	QRWSAETPHKPQFTFTSEPHAQLEENFCRNPDGDSHGPP--			
50	NOV5	450	460	470	480
	gi   15294659	TRNTQGPHTTVPDCAALMTSRHQS--WTPQITCSLRSVA			
	gi   10337615	WCYTDPRTDPEDYCALRCADDQPPSILDPDQVVEKCG			
	gi   123114	WCYTDPRTDPEDYCALRCADDQPPSILDPDQVVEKCG			
	gi   15299258	TRNTQGPHTTVPDCAALMTSRHQS--WTPQITCSLRSVA			
	gi   90615	WCYTDPRTDPEDYCALRCADDQPPSILDPDQVVEKCG			
55	NOV5	490	500	510	520
	gi   15294659	RGWIGWISGVVPSCAWLCAIRATHPGQSICGIG-----			
	gi   10337615	KRVDRLDQRRSKLRVVGHPGNSPWTVSLNRQGHFCGG			
60	gi   123114	KRVDRLDQRRSKLRVVGHPGNSPWTVSLNRQGHFCGG			
	gi   15299258	KRVDRLDQRRSKLRVVGHPGNSPWTVSLNRQGHFCGG			
	gi   90615	RGWIGWISGVVPSCAWLCAIRATHPGQSICGIGQGHFCGG			
65	NOV5	530	540	550	560
	gi   15294659	SLVKEQWILTARQCFSSCHMPLTGVEVWLGTLFQNPQHGE			
	gi   10337615	SLVKEQWILTARQCFSSCHMPLTGVEVWLGTLFQNPQHGE			
	gi   123114	SLVKEQWILTARQCFSSCHMPLTGVEVWLGTLFQNPQHGE			
70	gi   15299258	SLVKEQWILTARQCFSSCHMPLTGVEVWLGTLFQNPQHGE			

```

gi|90615| SLVKEQWYLTARQCIWSCHPLTGYEVWLGTHNONPOPGE
          570      580      590      600
5  NOV5      PSLQRPVPAKMVCGPSGSQLVLLKLESVTLNQRVALICL
   gi|15294659| PSLQRPVPAKMVCGPSGSQLVLLKLESVTLNQRVALICL
   gi|10337615| PSLQRPVPAKMVCGPSGSQLVLLKLESVTLNQRVALICL
   gi|123114| PSLQRPVPAKMVCGPSGSQLVLLKLESVTLNQRVALICL
   gi|15299258| PSLQRPVPAKMVCGPSGSQLVLLKLESVTLNQRVALICL
10 gi|90615| APLQRPVPAKMVCGPSGSQLVLLKLESVTLNQRVALICL
          610      620      630      640
15 NOV5      PPEWYVPPGKCEIAGWGETKGTGNDTVLNVALLNVISN
   gi|15294659| PPEWYVPPGKCEIAGWGETKGTGNDTVLNVALLNVISN
   gi|10337615| PPEWYVPPGKCEIAGWGETKGTGNDTVLNVALLNVISN
   gi|123114| PPEWYVPPGKCEIAGWGETKGTGNDTVLNVALLNVISN
   gi|15299258| PPEWYVPPGKCEIAGWGETKGTGNDTVLNVALLNVISN
20 gi|90615| PPEWYVPPGKCEIAGWGETKGTGNDTVLNVALLNVISN
          650      660      670      680
25 NOV5      QECNIKHRC-----RGDYGGPLACFT
   gi|15294659| QECNIKHRCGRV ESE CT GLAPVGACEGDYGGPLACFT
   gi|10337615| QECNIKHRCGRV ESE CT GLAPVGACEGDYGGPLACFT
   gi|123114| QECNIKHRCGRV ESE CT GLAPVGACEGDYGGPLACFT
   gi|15299258| QECNIKHRCGRV ESE CT GLAPVGACEGDYGGPLACFT
30 gi|90615| QECNIKHRCGRV ESE CT GLAPVGACEGDYGGPLACFT
          690      700      710      720
35 NOV5      HNCWVLEGIIPNRVCARSRWPAVFTRVSVFVDWIHKVMR
   gi|15294659| HNCWVLEGIIPNRVCARSRWPAVFTRVSVFVDWIHKVMR
   gi|10337615| HNCWVLEGIIPNRVCARSRWPAVFTRVSVFVDWIHKVMR
   gi|123114| HNCWVLEGIIPNRVCARSRWPAVFTRVSVFVDWIHKVMR
   gi|15299258| HNCWVLEGIIPNRVCARSRWPAVFTRVSVFVDWIHKVMR
40 gi|90615| HNCWVLEGIIPNRVCARSRWPAVFTRVSVFVDWIHKVMR
          730      740      750      760
45 NOV5      LG
   gi|15294659| LG
   gi|10337615| LG
   gi|123114| LG
   gi|15299258| LG
   gi|90615| LG

```

Tables 5E-N list the domain description from DOMAIN analysis results against NOV5. This indicates that the NOV5 sequence has properties similar to those of other proteins known to contain this domain.

**Table 5E. Domain Analysis of NOV5**

gnl|Pfam|pfam00051, kringle, Kringle domain. Kringle domains have been found in plasminogen, hepatocyte growth factors, prothrombin, and apolipoprotein A. Structure is disulfide-rich, nearly all-beta. (SEQ ID NO:85)  
 CD-Length = 79 residues, 100.0% aligned  
 Score = 117 bits (292), Expect = 3e-27

Query: 191 CWCNGEEYRGAVDRTESGRECORWDLQHPHQHPF-EPGKFLDQGLDDNYCRNPDGSRP 249  
 | ||| ||| |||| ||||| |||+| |++ +|| +||| |||

Sbjct 1 CYHNGENYRGTA TTESGAPCQRWDSQTPHRHSKYTPERYPAKGLGENYCRNPDGDERP 60

Query 250 WCYT TDPQIEREFCDLPRC 268  
 |||||++ |++|||

5 Sbjct: 61 WCYT TDPVRWEYCDIPRC 79

**Table 5F. Domain Analysis of NOV5**

gnl|Pfam|pfam00051, kringle, Kringle domain. Kringle domains have been found in plasminogen, hepatocyte growth factors, prothrombin, and apolipoprotein A. Structure is disulfide-rich, nearly all-beta. (SEQ ID NO:85)  
 CD-Length = 79 residues, 100.0% aligned  
 Score = 112 bits (279), Expect = 9e-26

10 Query: 283 CFRGKGEGYRGTA TTTAGVPCQRWDAQIPHQHRF-TPEKYACKDLRENFCRNPDGSEAP 341  
 |+ || || |||||++ |++|||

Sbjct: 1 CYHNGENYRGTA TTESGAPCQRWDSQTPHRHSKYTPERYPAKGLGENYCRNPDGDERP 60

Query: 342 WCFTLRPGMRAAFQYQIRRC 361  
 ||+| |++| +| |||

15 Sbjct: 61 WCYT TDPVRWEYC-DIPRC 79

**Table 5G. Domain Analysis of NOV5**

gnl|Pfam|pfam00051, kringle, Kringle domain. Kringle domains have been found in plasminogen, hepatocyte growth factors, prothrombin, and apolipoprotein A. Structure is disulfide-rich, nearly all-beta. (SEQ ID NO:85)  
 CD-Length = 79 residues, 100.0% aligned  
 Score = 104 bits (259), Expect = 2e-23

Query: 110 CIMNNGVG YRGTMATTVGG LPCQAWSHKFPNDH-KYTPT--LRNGLEENFCRNPDGDPGG 166  
 | || |||||++ | ||| | + |+ | ||| || |||+||| |||

20 Sbjct: 1 CYHNGENYRGTA TTESGAPCQRWDSQTPHRHSKYTPERYPAKGLGENYCRNPDGDE-R 59

Query: 167 PWCYT TDPAVRFQSCGIKSC 186  
 ||||| ||++ | |||

25 Sbjct: 60 PWCYT TDPVRWEYCDIPRC 79

**Table 5H. Domain Analysis of NOV5**

gnl|Smart|smart00130, KR, Kringle domain; Named after a Danish pastry. Found in several serine proteases and in ROR-like receptors. Can occur in up to 38 copies (in apolipoprotein(a)). Plasminogen-like kringles possess affinity for free lysine and lysine- containing peptides. (SEQ ID NO:86)  
 CD-Length = 83 residues, 97.6% aligned  
 Score = 111 bits (278), Expect = 1e-25

Query: 191 CVWCNGEEYRGAVDR TTESGREGCQRWDLQHPHQHPFEPGKFLDQGLDDNYCRNPDG-SERP 249  
 | ||| ||| |++| ||||| | ||| | | +| + ||+ ||||| |||

30 Sbjct: 3 CYAGNGESYRGTA TTKSGKPCQRWDSQTPHLHRFTPERFPGLGHEHNYCRNPDGDSEGP 62

Query: 250 WCYT TDPQIEREFCDLPRCGS 270

Sbjct: 63      ||||| + |++||  
WCYTDPNVRWEYCDIPQCES 83

**Table 5I. Domain Analysis of NOV5**

gnl|Smart|smart00130, KR, Kringle domain; Named after a Danish pastry. Found in several serine proteases and in ROR-like receptors. Can occur in up to 38 copies (in apolipoprotein(a)). Plasminogen-like kringles possess affinity for free lysine and lysine- containing peptides. (SEQ ID NO:86)  
CD-Length = 83 residues, 97.6% aligned  
Score = 106 bits (265), Expect = 4e-24

5

Query: 108 RTCIMNNGVGVRGTMTTVGGLPCQAWSHKFPNDHKYTPTLRN--GLEENFCRNPDPG 165  
|| || ||| +|| ||| | + | + ||| ||| +||| |||  
Sbjct: 1 RDCYAGNGESYRGTAATTKSGKPCQRWDSQTPHLHRFTPERFPPELGLEHNYCRNPDPGSE 60  
10 Query: 166 GPWCYTTPAVRFQSCGIKSC 186  
||| ||| ||++ || |  
Sbjct: 61 GPWCYTTPNVRWEYCDIPQC 81

**Table 5J. Domain Analysis of NOV5**

gnl|Smart|smart00130, KR, Kringle domain; Named after a Danish pastry. Found in several serine proteases and in ROR-like receptors. Can occur in up to 38 copies (in apolipoprotein(a)). Plasminogen-like kringles possess affinity for free lysine and lysine- containing peptides. (SEQ ID NO:86)  
CD-Length = 83 residues, 97.6% aligned  
Score = 104 bits (260), Expect = 1e-23

15

Query 283 CFRGKGEYRGTTANTTTAGVPCQRWDAQIPHQRFTPEKYACKDLRENFCRNPDPG-SEAP 341  
|+ | || |||+|| +| |||+|| || |||++ | | +||| || |  
Sbjct 3 CYAGNGESYRGTAATTKSGKPCQRWDSQTPHLHRFTPERFPPELGLEHNYCRNPDPGSEGP 62  
20 Query 342 WCFTLRPGMRAAFQYQIRRCTD 363  
||+| | +| +| | +|  
Sbjct 63 WCYTDPNVRWEYCD-IPQCES 83

**Table 5K. Domain Analysis of NOV5**

gnl|Smart|smart00020, Tryp\_Sp, Trypsin-like serine protease; Many of these are synthesised as inactive precursor zymogens that are cleaved during limited proteolysis to generate their active forms. A few, however, are active as single chain molecules, and others are inactive due to substitutions of the catalytic triad residues. (SEQ ID NO:87)  
CD-Length = 230 residues, 79.1% aligned  
Score = 110 bits (274), Expect = 3e-25

25

Query: 504 PLTGYEVWLGTLFQNPQHGEPSLQVRVPAKMVCGPSGSQ-----LVLLKLESVTLNQR 557  
+ | ||+ + | | +||+ | + + + ||| ||+  
Sbjct: 49 APSSIRVRLGSHDLSSGEET---QTVKSVKIVHPNYPSTYDNDIALLLKLEPVTLSDT 105

Query: 558 VALICLPPEWYVPPGKCEIAGWGETKGTG--NDTVLNVALLNVISNQECNIKHRGR-- 613  
 | ||| | | | | | ++||| | + | + ++|| | + |  
 Sbjct: 106 VRPICLPSSGYNVPAGTTCTVSGWGRTESSGSLPDTLQEVNVPPIVSNATCRRAYSGGPA 165

5 Query: 614 -----GDYGGPLACFTHNCWVLEGIIIPNRV-CARSCWPAVFTRVS 653  
 | | ||| | | | | | + | | | | | + ||||  
 Sbjct: 166 ITDNMLCAGGLEGGKDACQGDSSGGPLVC-NDPRWVLVGIVSWGSGYGCARPKNKPGVYTRVS 224

10 Query: 654 VFVDWI 659  
 ++|||  
 Sbjct: 225 SYLDWI 230

#### Table 5L. Domain Analysis of NOV5

gnl|Pfam|pfam00089, trypsin, Trypsin. Proteins recognized include all proteins in families S1, S2A, S2B, S2C, and S5 in the classification of peptidases. Also included are proteins that are clearly members, but that lack peptidase activity, such as haptoglobin and protein Z (PRTZ\*). (SEQ ID NO:88)  
 CD-Length = 217 residues, 77.9% aligned  
 Score = 92.0 bits (227), Expect = 9e-20

15 Query: 508 YEVWLGTLFQNPQHGEPSLQRPVAKMV---CGPSGSQLVLLKCLERSVTLNQRVALICL 563  
 | | | | | + | + | | ++ | + + ||| + | | | | |  
 Sbjct: 49 VRVVLGE--HNLGTTEGTEQKFDVKKIIIVHPNYNPDTNDIALLLKLSKSPVTLGDTVIRPICL 106

20 Query: 564 PPEWYVPPGKCEIAGWGETKGTGNDTVLNVALLNVISNQECNIKHRG----- 612  
 | + | | | ++||| | | | | ++ ++| + | + |  
 Sbjct: 107 PSASSDLPVGTTCVSGWGRTKNLGSDTLQEVVPIVSRETCRSAYGGTVTDTMICAGA 166

25 Query: 613 -----RGDYGGPLACFTHNCWVLEGIIIPNRVCARSCWPAVFTRVSFVVDWI 659  
 + || ||| | | | | + | | + | + |||| ++|||  
 Sbjct: 167 LGGKDACQGDSSGGPLVCSGD---ELVGIVSWGYGCAVGNYPGVYTRVSRYLWI 217

#### Table 5M. Domain Analysis of NOV5

gnl|Smart|smart00473, PAN\_AP, divergent subfamily of APPLE domains; Apple-like domains present in Plasminogen, C. elegans hypothetical ORFs and the extracellular portion of plant receptor-like protein kinases. Predicted to possess protein- and/or carbohydrate-binding functions. (SEQ ID NO:89)  
 CD-Length = 79 residues, 94.9% aligned  
 Score = 52.0 bits (123), Expect = 1e-07

30 Query: 25 DFQVLRGTEHQHLLHAVVPGPWQEDVADAEACAGRC-GPLMDCRAFHYNVSSHGCQLLPW 83  
 | | | + | | | | | + | | + | | + | | |  
 Sbjct: 5 CFVRLPNTKL-----PDFSPIVISVASLEECAQKCLNSNCSCRSFTYNNDTKGCLLWSE 58

35 Query: 84 TQHSPHTRLRRSGRCDLFQKK 104  
 + + | | | ++|  
 Sbjct: 59 SSLGDARQLLPSSGGVDYYEKI 79

**Table 5N. Domain Analysis of NOV5**

gnl|Pfam|pfam00024, PAN, PAN domain. The PAN domain contains a conserved core of three disulphide bridges. In some members of the family there is an additional fourth disulphide bridge the links the N and C termini of the domain. The domain is found in diverse proteins, in some they mediate protein-protein interactions, in others they mediate protein-carbohydrate interactions. (SEQ ID NO:159)  
CD-Length = 78 residues, 76.9% aligned  
Score = 50.1 bits (118), Expect = 4e-07

Query: 45 PWQEDVADAEACAGRCGPLMD-CRAFHYNVSSHGCQLLPWTQHSPHTRLRRSGRCDLFQK 103  
+| |||| | ||+| || | | + | | + | ++|  
Sbjct: 17 ISVVNVPSLEECAALCLEEPRVCRSFTYNNKSKQCLLKSESSGSLPRLKRPSQKVDYYEK

5

Macrophage-stimulating protein (MSP) is an 80-kD serum protein with homology to hepatocyte growth factor (HGF)(Sakamoto O, et.al.; J Clin Invest 1997 Feb 15;99(4):701-9). Its receptor, RON tyrosine kinase, is a new member of the HGF receptor family. The MSP-RON signaling pathway has been implicated in the functional regulation of mononuclear phagocytes. However, the function of this pathway in other types of cells has not been elucidated. Here we show that in contrast to the HGF receptor, which was expressed at the basolateral surface, RON was localized at the apical surface of ciliated epithelia in the airways and oviduct. In addition, MSP was found in the bronchoalveolar space at biologically significant concentrations. MSP bound to RON on normal human bronchial epithelial cells with a high affinity ( $K_d = 0.5$  nM) and induced autophosphorylation of RON. Activation of RON by MSP led to a significant increase in ciliary beat frequency of human nasal cilia. These findings indicate that the ciliated epithelium of the mucociliary transport apparatus is a novel target of MSP. Ciliary motility is critical for mucociliary transport. Our findings suggest that the MSP-RON signaling pathway is a novel regulatory system of mucociliary function and might be involved in the host defense and fertilization.

Macrophage stimulating protein (MSP; 142408), also known as hepatocyte growth factor-like protein (HGFL), is structurally related to hepatocyte growth factor. Gaudino et al. (EMBO J. 13: 3524-3532, 1994) showed that the RON gene is expressed at the cell surface of several epithelial cell types in addition to granulocytes and monocytes. The RON mRNA is translated into a glycosylated precursor that is cleaved into a 185-kD heterodimer of 35-kD (alpha) and 150-kD (beta) subunits joined by the predicted disulfide linkage. Gaudino et al. (1994) further demonstrated that the beta chain undergoes tyrosine phosphorylation upon stimulation by MSP. By isotopic in situ hybridization, Ronsin et al. (Oncogene 8: 1195-1202, 1993) mapped the RON gene to 3p21, with the most probable location being 3p21.3. The gene

ed on 3p21, a region of frequent deletion or mutation in small cell

g hepatocyte growth factor (HGF; 142409) and its MET receptor are of ligand and receptor on the same chromosome, and the structural and HGF suggested that the ligand for RON might be MSP. Wang (1994) showed that this is indeed the case. Their experiments showed that the gene product is a specific cell surface receptor for MSP. A human cDNA library was screened with an oligonucleotide probe for the protein (MSP) to clone an MSP cDNA (Yoshimura T, et.al.; J Biol Chem 269:15461-8). Deduced sequences of isolated clones were compared with the sequences of MSP. MSP9 cDNA encoded most of the known sequence of MSP, a segment of the 5' end of the open reading frame. Consequently, a cDNA that encoded the complete MSP amino acid sequence was isolated. Culture fluid from COS-7 cells transfected with this full-length cDNA showed biological activity, and the expressed MSP was detected by Western blotting with an antibody against native MSP. The deduced amino acid sequence of MSP contains domains, which have been found in hepatocyte growth factor and the blood coagulation system. Among them, MSP has the highest sequence homology with hepatocyte growth factor (45% identity). The MSP cDNA hybridized strongly to human liver mRNA, to a lesser extent to mRNA from kidney and pancreas, suggesting that liver is the source of MSP. Several cloned and sequenced MSP cDNAs had different 5' ends, suggesting that alternatively spliced MSP mRNAs may occur. This was confirmed by Northern blotting probed with an MSP cDNA, which showed more than one mRNA species. Although the gene coding for MSP is on chromosome 3, the sequence of the cDNA is identical with a unique sequence in chromosome 1, indicating that there are two MSP genes, located on chromosomes 3 and 1.

Hepatocyte growth factor-like protein (HLP) shares a 50% homology with hepatocyte growth factor, the biological function of HLP has remained unknown (S Lett 1993 Oct 25;333(1-2):61-6). Addition of conditioned medium to macrophages transfected with the expression plasmid for HLP cDNA to cultures of macrophages induced specific activation of macrophages, and the factor was purified from the conditioned medium. The purified factor has a molecular weight of 70 kDa on SDS-PAGE, and this M(r) is in agreement with that of hepatocyte growth factor (MSP) previously purified from human serum, as well as with

the predicted M(r) of HLP. Amino acid composition of the purified protein coincided with the compositions of human HLP and MSP. Together with the finding that the partial amino acid sequences of MSP are highly homologous to that of HLP, we conclude that the biological function of HLP is to activate macrophages and that HLP and MSP are identical molecules.

5           The disclosed NOV5 nucleic acid of the invention encoding a Macrophage Stimulating Protein Precursor -like protein includes the nucleic acid whose sequence is provided in Table 5A or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 5A while still encoding a protein that maintains its Macrophage Stimulating Protein Precursor -like activities  
10           and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way  
15           of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 10% percent of the bases may be so  
20           changed.

          The disclosed NOV5 protein of the invention includes the Macrophage Stimulating Protein Precursor -like protein whose sequence is provided in Table 5B. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 5B while still encoding a protein that maintains its  
25           Macrophage Stimulating Protein Precursor -like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 39% percent of the residues may be so changed.

          The NOV5 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in Aicardi-Goutieres syndrome 1; Brugada syndrome;  
30           Deafness, autosomal recessive 6; Heart block, nonprogressive; Heart block, progressive, 2; Ichthyosiforme erythroderma, congenital, nonbullous; Long QT syndrome-3; Night blindness; congenital stationary; Pituitary ACTH-secreting adenoma; Small-cell cancer of lung; Ventricular fibrillation, idiopathic; entricular tachycardia, idiopathic; HIV infection, susceptibility/resistance to; Von Hippel-Lindau (VHL) syndrome; Cirrhosis; Transplantation



as well as other diseases, disorders and conditions. The NOV5 nucleic acid, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

NOV5 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. For example the disclosed NOV5 protein have multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, contemplated NOV5 epitope is from about amino acids 20 to 80. In other embodiments, NOV5 epitope is from about amino acids 90 to 120, from about amino acids 140 to 180, from about amino acids 190 to 340, from about amino acids 350 to 460, from about amino acids 500 to 530, from about amino acids 570 to 590, and from about amino acids 600 to 620, or from about amino acids 420 to 460. This novel protein also has value in development of powerful assay system for functional analysis of various human disorders, which will help in understanding of pathology of the disease and development of new drug targets for various disorders.

## NOV6

A disclosed NOV6 nucleic acid of 634 nucleotides (also referred to as GM382a20\_A) encoding a novel Fatty Acid-Binding Protein-like protein is shown in Table 6A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 58-60 and ending with a TAA codon at nucleotides 460-462. A putative untranslated region upstream from the initiation codon and downstream from the termination codon is underlined in Table 6A, and the start and stop codons are in bold letters.

**Table 6A. NOV6 Nucleotide Sequence (SEQ ID NO:17)**

<p>CAAGCTGCCACGCCGACGGCAACCCTGCTCTGCATGCCCGCCCGCCCGTGCCACCA<b>AT</b>GCCACAGTTC  AGCAGCTGGGAGGAAGATGGCGCCTGGTGGACAGCAAACGCTTTGATGAATACATGAAGGAGGGAGGAGT  GGGAAC'TGCTTTGCGAAAAATGGACGCAATGGCCAAGCCAGATTGTATCATCACTTGTGATGCCAAAAAC  CTCACCATAAAAACCGAGAGCACTTTGAAAAACACAGTTTCTTGTACCCTGGGAGAGAAGTTTGAAGAAA  CCACAGCTGATGGCAGAAAACTCAGACTGTGTGCAGCTTTCAGATGGTGCATTGGTTCAGCATCAGGA  GTGGGATGGGAAGGAAAAACAATAACAAGAAAACTGAAAGATGGGAAATTAGTGGTGTACTGTGTCATG  AACAATGTCGCCTGTACTCGGATCTATGAAAAAGTAGAAT<b>TAAAAAT</b>CCATCATCACTTTGGACAGGAGT  TAACTAATAGAATGATCAAGCTCAGTTCAATGAGCAAATCTCCATAGTGTTTTTTTTCATTACTGTGTTT  AATTATCTTTATCACAAACGTTTCACATGCAGCTATTTCAAAGTGTCTTGGATTAATTAGGATCATCCCT  TTGG</p>
---

The disclosed NOV6 nucleic acid sequence , located on chromosome 15, has 537 of 589 bases (91 %) identical to a Fatty Acid-Binding Protein mRNA from *Homo sapiens* (GENBANK-ID: HUMFABPHA) ( $E = 3.7e^{-102}$ ).

A disclosed NOV6 polypeptide (SEQ ID NO:18) encoded by SEQ ID NO:17 is 134 amino acid residues and is presented using the one-letter amino acid code in Table 6B. Signal P, Psort and/or Hydropathy results predict that NOV6 contains no signal peptide and is likely to be localized in the cytoplasm with a certainty of 0.4500. In other embodiments, NOV6 is also likely to be localized to the mitochondrial matrix space with a certainty of 0.3600, or to the lysosome (lumen) with a certainty of 0.1000.

**Table 6B. Encoded NOV6 protein sequence (SEQ ID NO:18).**

MATVQQLGGRWRLVDSKRFDEYMKEGGVGTALRKMDAMAKPDCIITCDGKNLTIKTESTLTKTQFSCTLGEKFEET TADGRKTQTVCSFADGALVQHQEWDGKENTITRKLKDGKLVVYCVMMNVACTRIYEKVE
---

The disclosed NOV6 amino acid sequence has 124 of 135 amino acid residues (91 %) identical to, and 126 of 135 amino acid residues (93 %) similar to, the 135 amino acid residue Fatty Acid-Binding protein from *Homo sapiens* (Q01469) ( $E = 2.1e^{-61}$ ).

NOV6 is expressed in Sensory System.Skin; Nervous System.Brain; Male Reproductive System.Testis; Respiratory System.Lung, Larynx; Female Reproductive System; .Placenta; Whole Organism; Cardiovascular System.Heart; Endocrine System.Parathyroid Gland; Hematopoietic and Lymphatic System, Hematopoietic Tissues, Liver, Tonsils; Gastro-intestinal/Digestive System.Large Intestine, Colon, Stomach, Oesophagus; Urinary System.Kidney. In addition, the sequence is predicted to be expressed in the following tissues because of the expression pattern of (GENBANK-ID: ACC:Q05816) a closely related Fatty Acid-Binding Protein homolog in species *Mus musculus* : Sensory System.Skin; Nervous System.Brain; Male Reproductive System.Testis; Respiratory System.Lung, Larynx; Female Reproductive System; .Placenta; Whole Organism; Cardiovascular System.Heart; Endocrine System.Parathyroid Gland; Hematopoietic and Lymphatic System, Hematopoietic Tissues, Liver, Tonsils; Gastro-intestinal/Digestive System.Large Intestine, Colon, Stomach, Oesophagus; Urinary System.Kidney.

TaqMan data for NOV6 can be found below in Example 2. NOV6 also has homology to the amino acid sequences shown in the BLASTP data listed in Table 6C.

**Table 6C. BLAST results for NOV6**

Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
---------------------------	-------------------	----------------	-----------------	------------------	--------

gi 13651468 ref XP_016351.1	similar to GASTRIN/CHOLECYSTOKININ TYPE B RECEPTOR (CCK-B RECEPTOR) (CCK-BR) (H. sapiens) [Homo sapiens]	135	122/135 (90%)	126/135 (92%)	1e-58
gi 4557581 ref NP_01435.1	fatty acid binding protein 5 (psoriasis-associated); E-FABP [Homo sapiens]	135	124/135 (91%)	126/135 (92%)	1e-58
gi 13651882 ref XP_011655.5	fatty acid binding protein 5 (psoriasis-associated) [Homo sapiens]	135	119/135 (88%)	124/135 (91%)	6e-57
gi 13651563 ref XP_015760.1	similar to GASTRIN/CHOLECYSTOKININ TYPE B RECEPTOR (CCK-B RECEPTOR) (CCK-BR) (H. sapiens) [Homo sapiens]	135	118/135 (87%)	125/135 (92%)	5e-56
gi 6648071 sp P55052 FABE_BOVIN	FATTY ACID-BINDING PROTEIN, EPIDERMAL (E-FABP)	135	117/135 (86%)	124/135 (91%)	1e-55

The homology of these sequences is shown graphically in the ClustalW analysis shown in Table 6D.

**Table 6D Information for the ClustalW proteins**

- 1) NOV6 (SEQ ID NO:18)
- 2) gi|13651468|ref|XP\_016351.1| similar to GASTRIN/CHOLECYSTOKININ TYPE B RECEPTOR (CCK-B RECEPTOR) (CCK-BR) (H. sapiens) [Homo sapiens] (SEQ ID NO:56)
- 3) gi|4557581|ref|NP\_001435.1| fatty acid binding protein 5 (psoriasis-associated); E-FABP [Homo sapiens] (SEQ ID NO:57)
- 4) gi|13651882|ref|XP\_011655.5| fatty acid binding protein 5 (psoriasis-associated) [Homo sapiens] (SEQ ID NO:58)
- 5) gi|13651563|ref|XP\_015760.1| similar to GASTRIN/CHOLECYSTOKININ TYPE B RECEPTOR (CCK-B RECEPTOR) (CCK-BR) (H. sapiens) [Homo sapiens] (SEQ ID NO:59)
- 6) gi|6648071|sp|P55052|FABE\_BOVIN FATTY ACID-BINDING PROTEIN, EPIDERMAL (E-FABP) (SEQ ID NO:60)

		10	20	30	40
NOV6		..... ..... ..... ..... ..... .....			
20	gi 13651468	MATVQQLCGRWRLVDSKRFDEYMKELGVGALRKMDAMAK			
	gi 4557581	MATVQQLCGRWRLVDSKGFDEYMKELGVGIALRKMGAMAK			
	gi 13651882	MATVQQLCGRWRLVDSKGFDEYMKELGVGIALRKMDTAK			
	gi 13651563	MATVQQLCGRWRLVDSKGFDEYMKELGVGIALRKMGAMAK			
25	gi 6648071	MATVQQLCGRWRLVDSKGFDEYMKELGVGALRKMGAMAK			
		50	60	70	80
		..... ..... ..... ..... ..... .....			
NOV6		PDCIITCDGKNLTIKTESTLKTQFSCITLGEKFEETTADG			
30	gi 13651468	PDCIITCDGKNLTIKTESTLKTQFSCITLGEKFEETTADG			
	gi 4557581	PDCIITCDGKNLTIKTESTLKTQFSCITLGEKFEETTADG			
	gi 13651882	PDCIITCDGKNLTIKTESTLKTQFSCITLGEKFEETTADG			
	gi 13651563	PDCIITCDGKNLTIKTESTLKTQFSCITLGEKFEETTADG			
	gi 6648071	PDCIITCDGKNLTIKTESTLKTQFSCITLGEKFEETTADG			
35		90	100	110	120



Epidermal fatty acid binding protein (E-FABP) was cloned by as a novel keratinocyte protein by Madsen et al (1992, PMID: 1512466) from skin of psoriasis patients. Later using quantitative Western blot analysis, Kingma et al. (1998, PMID: 9521644) have shown that in addition to the skin, bovine E-FABP is expressed in retina, testis, and lens. Since E-FABP was originally identified from the skin of psoriasis patients, it is also known as psoriasis-associated fatty acid-binding protein (PA-FABP). PA-FABP is a cytoplasmic protein, and is expressed in keratinocytes. It is highly up-regulated in psoriatic skin. It shares similarity to other members of the fatty acid-binding proteins and belongs to the fabp/p2/crbp/crabp family of transporter. PA-FABP is believed to have a high specificity for fatty acids, with highest affinity for c18 chain length. Decreasing the chain length or introducing double bonds reduces the affinity. PA-FABP may be involved in keratinocyte differentiation.

Immunohistochemical localization of the expression of E-FABP in psoriasis, basal and squamous cell carcinomas has been carried out in order to obtain indirect information, at the cellular level, on the transport of the fatty acidss. (Masouye et al, 1996, PMID: 8726632). E-FABP was localized in the upper stratum spinosum and stratum granulosum in normal and non-lesional psoriatic skin. In contrast, lesional psoriatic epidermis strongly expressed E-FABP in all suprabasal layers, like nonkeratinized oral mucosa. The basal layer did not express E-FABP reactivity in any of these samples. Accordingly, basal cell carcinomas were E-FABP negative whereas only well-differentiated cells of squamous cell carcinomas expressed E-FABP. This suggests that E-FABP expression is related to the commitment of keratinocyte differentiation and that the putative role of E-FABP should not be restricted to the formation of the skin lipid barrier. Since the pattern of E-FABP expression mimics cellular FA transport, our results suggest that lesional psoriatic skin and oral mucosa have a higher metabolism/transport for FAs than normal and non-lesional psoriatic epidermis.

The disclosed NOV6 nucleic acid of the invention encoding a Fatty acid binding protein-like protein includes the nucleic acid whose sequence is provided in Table 6A or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 6A while still encoding a protein that maintains its Fatty acid binding protein-like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example,

modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 10% percent of the bases may be so changed.

The disclosed NOV6 protein of the invention includes the Fatty acid binding protein-like protein whose sequence is provided in Table 6B. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 6B while still encoding a protein that maintains its Fatty acid binding protein-like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 14% percent of the residues may be so changed.

The above defined information for this invention suggests that these Fatty acid binding protein-like proteins (NOV6) may function as a member of a "Fatty acid binding proteinfamily". Therefore, the NOV6 nucleic acids and proteins identified here may be useful in potential therapeutic applications implicated in (but not limited to) various pathologies and disorders as indicated below. The potential therapeutic applications for this invention include, but are not limited to: protein therapeutic, small molecule drug target, antibody target (therapeutic, diagnostic, drug targeting/cytotoxic antibody), diagnostic and/or prognostic marker, gene therapy (gene delivery/gene ablation), research tools, tissue regeneration *in vivo* and *in vitro* of all tissues and cell types composing (but not limited to) those defined here.

The nucleic acids and proteins of NOV6 are useful in Cardiomyopathy, Atherosclerosis, Hypertension, Congenital heart defects, Aortic stenosis, Atrial septal defect (ASD), Atrioventricular (A-V) canal defect, Ductus arteriosus, Pulmonary stenosis, Subaortic stenosis, Ventricular septal defect (VSD), valve diseases, Tuberous sclerosis, Scleroderma, Transplantation, Endometriosis, Inflammatory bowel disease, Diverticular disease, Hirschsprung's disease, Crohn's Disease, Hemophilia, hypercoagulation, Idiopathic thrombocytopenic purpura, immunodeficiencies, Osteoporosis, Hypercalceimia, Arthritis, Von Hippel-Lindau (VHL) syndrome, Alzheimer's disease, Stroke, Tuberous sclerosis, hypercalceimia, Parkinson's disease, Huntington's disease, Cerebral palsy, Epilepsy, Asthma, allergy, ARDS, Lesch-Nyhan syndrome, Multiple sclerosis, Leukodystrophies, Behavioral disorders, Addiction, Anxiety, Pain, Neuroprotection Fertility, psoriasis, cancer including but not limited to basal and squamous cell carcinomas, obesity, diabetes, and/or other pathologies and disorders involving fatty acid transport of skin, oral mucosa as well as other organs. The novel NOV6 nucleic acid encoding NOV6 protein., or fragments thereof, may further be

useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed. These materials are further useful in the generation of antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods.

5 NOV6 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. For example the disclosed NOV6 protein have multiple hydrophilic regions,  
10 each of which can be used as an immunogen. In one embodiment, contemplated NOV6 epitope is from about amino acids 10 to 35. In other embodiments, NOV6 epitope is from about amino acids 40 to 45, from about amino acids 50 to 85, or from about amino acids 90 to 110. This novel protein also has value in development of powerful assay system for functional analysis of various human disorders, which will help in understanding of pathology of the  
15 disease and development of new drug targets for various disorders.

#### NOV7

A disclosed NOV7 nucleic acid of 822 nucleotides (also referred to  
sggc\_draft\_dj895c5\_20000819) encoding a novel Gap junction beta-5 protein-like protein is  
20 shown in Table 7A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 1-3 and ending with a TAA codon at nucleotides 800-802. In Table 7A, the 5' and 3' untranslated regions are underlined and the start and stop codons are in bold letters.

**Table 7A. NOV7 Nucleotide Sequence (SEQ ID NO:19)**

<p><u>CATGA</u>ACTGGGCATTTCTGCAGGGCCTGCTGAGTGCGGTGAACAAGTACTCCACAGTGCTGAGCCGCATCT GGCTGTCTGTGGTGTTCATCTTTTCGTGTGCTGGTGACGTGGTGGCAGCGGAGGAGGTGTGGGACGATGAG CAGAAGGACTTTGTCTGCAACACCAAGCAGCCCGGCTGCCCCAACGTCTGCTATGACGAGTTCTTCCCCGT GTCCACGTCGCCTCTGGGCCCTACAGCTCATCCTGGTCACGTGCCCTCACTGCTCGTGGTCATGCACG TGGCCTACCGCGAGGAACGCGAGCGCAAGCACCACCTGAAACACGGGCCCAATGCCCGTCCCTGTACGAC AACCTGAGCAAGAAGCGGGCGGACTGTGGTGGACGTACTTGCTGAGCCTCATCTTCAAGGCCCGCTGGA TGCTGGCTTCCTCTATATCTTCCACCGCCTCTACAAGGATTATGACATGCCCCGCGTGGTGGCTGCTCCG TGGAGCCTTGCCCCACACTGTGGACTGTTACATCTCCCGGCCACGGAGAAGAAGGTCTTCACCTACTTC ATGGTGACCACAGCTGCCATCTGCATCCTGCTCAACCTCAGTGAAGTCTTCTACCTGGTGGGCAAGAGGTG CATGGAGATCTTCGGCCCCAGGCACCGGCGGCTCGGTGCCGGGAATGCCTACCCGATACGTGCCCCACCAT ATGTCCTCTCCAGGGAGGCACCCTGAGGATGGGAACCTCTGTCTTAATGAAGGCTGGGTGGCCCCAGTG GATGCAGGTGGGTATCCAT<b>TAA</b>CCTGCGAGATCAGCAGATAA</p>
--

25 The disclosed NOV7 nucleic acid sequence, localized to the p34.3-36.11 region of chromosome 1, has 682 of 807 bases (84%) identical to a gb:GENBANK-

ID:RNCON303|acc:X76168 mRNA from *Rattus norvegicus* (*R.norvegicus* mRNA for connexin 30.3) ( $E = 3.7e^{-125}$ ).

A disclosed NOV7 polypeptide (SEQ ID NO:20) encoded by SEQ ID NO:19 is 266 amino acid residues and is presented using the one-letter amino acid code in Table 7B. Signal P, Psort and/or Hydropathy results predict that NOV7 has a signal peptide and is likely to be localized in the plasma membrane with a certainty of 0.6000. In other embodiments, NOV7 is also likely to be localized to the Golgi body with a certainty of 0.4000, to the endoplasmic reticulum (membrane) with a certainty of 0.3000, or the nucleus with a certainty of 0.2400. The most likely cleavage site for a NOV7 peptide is between amino acids 40 and 41, at: VAA-EE.

**Table 7B. Encoded NOV7 protein sequence (SEQ ID NO:20).**

```
MNWAFLQGLLSGVNKYSTVLSRIWLSVVFIFRVLVYVVAEEVWDDEQKDFVCNTKQPGCPNVCYDEFFPV
SHVRLWALQLILVTCPSLLVMHVAYREERERKHLKHGPNAPSLYDNLSSKRGGLWWTYLLSLIFKAAVD
AGFLYIFHRLYKDYDMPRVVACSVPCPHTVDCYISRPTEKKVFTYFMVTTAAICILLNLSEVFYLVGKRC
MEIFGPRHRRPRCRECLPDTCPFYVLSQGGHPEDGNSVLMKAGSAPVDAGGYP
```

The disclosed NOV7 amino acid sequence has 230 of 266 amino acid residues (86%) identical to, and 244 of 266 amino acid residues (91%) similar to, the 266 amino acid residue ptnr:SWISSPROT-ACC:Q02738 protein from *Mus musculus* (Mouse) (Gap Junction Beta-5 Protein (Connexin 30.3) (CX30.3) ( $4.4e^{-129}$ ).

NOV7 is expressed in at least kidney tissue. The sequence is predicted to be expressed in the following tissues because of the expression pattern of (GENBANK-ID: RNCON303|acc:X76168) a closely related {*R.norvegicus* mRNA for connexin 30.3 homolog in species *Rattus norvegicus* : kidney and thymus.

SNP data for NOV7 can be found below in Example 3. NOV7 also has homology to the amino acid sequence shown in the BLASTP data listed in Table 7C.

**Table 7C. BLAST results for NOV7**

Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 12229761 sp Q9NTQ9 CXB4_HUMAN	GAP JUNCTION BETA-4 PROTEIN (CONNEXIN 30.3) (CX30.3)	266	266/266 (100%)	266/266 (100%)	e-155
gi 544118 sp P36380 CXB5_RAT	GAP JUNCTION BETA-5 PROTEIN (CONNEXIN 30.3) (CX30.3)	265	230/266 (86%)	245/266 (91%)	e-132



gi 6680015 ref NP_032153.1	gap junction membrane channel protein beta 4; connexin 30.3 [Mus musculus]	266	230/266 (86%)	244/266 (91%)	e-131
gi 10835079 ref NP_005259.1	gap junction protein, beta 5 (connexin 31.1) [Homo sapiens]	273	153/226 (67%)	177/226 (77%)	3e-88
gi 4009522 gb AAC95472.1  (AF099731)	connexin 31.1 [Homo sapiens]	273	152/226 (67%)	176/226 (77%)	1e-87

The homology of these sequences is shown graphically in the ClustalW analysis shown in Table 7D.

**Table 7D. Information for the ClustalW proteins**

- 1) NOV7 (SEQ ID NO:20)
- 2) gi|12229761|sp|Q9NTQ9|CXB4\_HUMAN GAP JUNCTION BETA-4 PROTEIN (CONNEXIN 30.3) (CX30.3) (SEQ ID NO:61)
- 3) gi|544118|sp|P36380|CXB5\_RAT GAP JUNCTION BETA-5 PROTEIN (CONNEXIN 30.3) (CX30.3) (SEQ ID NO:62)
- 4) gi|6680015|ref|NP\_032153.1| gap junction membrane channel protein beta 4; connexin 30.3 [Mus musculus] (SEQ ID NO:63)
- 5) gi|10835079|ref|NP\_005259.1| gap junction protein, beta 5 (connexin 31.1) [Homo sapiens] (SEQ ID NO:64)
- 6) gi|4009522|gb|AAC95472.1| (AF099731) connexin 31.1 [Homo sapiens] (SEQ ID NO:65)

		10	20	30	40
NOV7		..... ..... ..... ..... ..... .....			
gi 12229761		MNWFLQGLLSGVNKYSTVL	SRIWLSVVFIFRVLVYVVA		
gi 544118		MNWFLQGLLSGVNKYSTVL	SRIWLSVVFIFRVLVYVVA		
gi 6680015		MNWFLQGLLSGVNKYSTAL	GRIWLSVVFIFRVLVYVVA		
gi 10835079		MNWSIFGLLSGVNKYSTAF	GRIWLSVVFIFRVLVYVVA		
gi 4009522		MNWSIFGLLSGVNKYSTAF	GRIWLSVVFIFRVLVYVVA		
		50	60	70	80
NOV7		..... ..... ..... ..... ..... .....			
gi 12229761		EEVWDDQKDFVCNTKQPGCPNVCYDEFFPVSHVRLWALQ			
gi 544118		EEVWDDQKDFVCNTKQPGCPNVCYDEFFPVSHVRLWALQ			
gi 6680015		EEVWDDQKDFVCNTKQPGCPNVCYDEFFPVSHVRLWALQ			
gi 10835079		EEVWDDQKDFVCNTKQPGCPNVCYDEFFPVSHVRLWALQ			
gi 4009522		EEVWDDQKDFVCNTKQPGCPNVCYDEFFPVSHVRLWALQ			
		90	100	110	120
NOV7		..... ..... ..... ..... ..... .....			
gi 12229761		LILVTCPSLLVVMHVAYREERERKHLKHGPNAPSLYDNL			
gi 544118		LILVTCPSLLVVMHVAYREERERKHLKHGPNAPSLYDNL			
gi 6680015		LILVTCPSLLVVMHVAYREERERKHLKHGPNAPSLYDNL			
gi 10835079		LILVTCPSLLVVMHVAYREERERKHLKHGPNAPSLYDNL			
gi 4009522		LILVTCPSLLVVMHVAYREERERKHLKHGPNAPSLYDNL			
		130	140	150	160
NOV7		..... ..... ..... ..... ..... .....			
gi 12229761		SKKRGGWLWWTYLLSLIFKAAVD	GFLYIFHRYKDYDMPR		
gi 544118		SKKRGGWLWWTYLLSLIFKAAVD	GFLYIFHRYKDYDMPR		
gi 6680015		SKKRGGWLWWTYLLSLIFKAAVD	GFLYIFHRYKDYDMPR		
gi 10835079		SKKRGGWLWWTYLLSLIFKAAVD	GFLYIFHRYKDYDMPR		
gi 4009522		SKKRGGWLWWTYLLSLIFKAAVD	GFLYIFHRYKDYDMPR		
		170	180	190	200
		..... ..... ..... ..... ..... .....			



**Table 7F. Domain Analysis of NOV7**

gnl|Smart|smart00037, CNX, Connexin homologues; Connexin channels participate in the regulation of signaling between developing and differentiated cell types. (SEQ ID NO:160)  
CD-Length = 34 residues, 100.0% aligned  
Score = 79.0 bits (193), Expect = 3e-16

Query: 42 EVWDDEQKDFVCNTKQPGCPNVCYDEFFPVSHVR 75  
          || ||| || |||+||| |||||+|||+|||  
Sbjct: 1 SVWGDEQSDFTCNTQQPGCENVCYDQFFPISHVR 34

5

Gap junctions are conduits that allow the direct cell-to-cell passage of small cytoplasmic molecules, including ions, metabolic intermediates, and second messengers, and thereby mediate intercellular metabolic and electrical communication. Gap junction channels consist of connexin protein subunits, which are encoded by a multigene family. Richard et al. (Nature Genet. 20: 366-369,1998) identified 2 expressed sequence tags (ESTs) from the human EST database by their similarity to mouse Gjb3 (see 603324) and Gjb5. By radiation hybrid mapping, they placed them in proximity to a sequence tagged site (STS) that is linked to GJA4 (121012) at 1p35.1. Richard et al. (1998) determined the cDNA sequences of the genes from which both ESTs were derived. Sequence similarity to rodent connexin genes established them as human homologs of Gjb3 and Gjb5, encoding Cx32 and Cx31.1, respectively. Richard et al. (1998) excluded GJB5 as a candidate gene for erythrokeratoderma variabilis (133200) by sequence analysis. Gap junctions are intercellular channels which connect adjacent cells and allow direct exchange of molecules of low molecular weight between them (Alves LA et.al.; Braz J Med Biol Res 2000 Apr;33(4):457-65). Such a communication has been described as fundamental in many systems due to its importance in coordination, proliferation and differentiation. Recently, it has been shown that gap junctional intercellular communication (GJIC) can be modulated by several extracellular soluble factors such as classical hormones, neurotransmitters, interleukins, growth factors and some paracrine substances. Herein, we discuss some aspects of the general modulation of GJIC by extracellular messenger molecules and more particularly the regulation of such communication in the thymus gland. Additionally, we discuss recent data concerning the study of different neuropeptides and hormones in the modulation of GJIC in thymic epithelial cells. We also suggest that the thymus may be viewed as a model to study the modulation of gap junction communication by different extracellular messengers involved in non-classical circuits, since this organ is under bidirectional neuroimmunoendocrine control. The

intercellular signaling system mediated by connexin channels is crucial for maintaining tissue homeostasis, growth control, development, and synchronized response of cells to stimuli (Richard G; *Exp Dermatol* 2000 Apr;9(2):77-96). This review summarizes the structure, assembly, and properties of the components of the complex and diverse connexin system, and their biological functions in skin. The importance of gap junctional intercellular communication for normal development and differentiation of human epidermis as well as the hearing function of the inner ear is illustrated by the examples of erythrokeratoderma variabilis and palmoplantar keratoderma associated with hearing loss. These autosomal dominant inherited disorders are caused by germline mutations in the connexin genes GJB3 (encoding connexin-31) and GJB2 (encoding connexin-26), respectively. Recent functional studies of individual connexin mutations suggest that they may exert a dominant inhibitory effect on normal connexin channel function and perturb gap junctional intercellular communication, resulting in phenotypic manifestation in patients with these disorders. Gap junction channels are sites of cytoplasmic communication between contacting cells. In vertebrates, they consist of protein subunits denoted connexins (Cxs) which are encoded by a gene family (Saez JC et.al.; *Braz J Med Biol Res* 2000 Apr;33(4):447-55). According to their Cx composition, gap junction channels show different gating and permeability properties that define which ions and small molecules permeate them. Differences in Cx primary sequences suggest that channels composed of different Cxs are regulated differentially by intracellular pathways under specific physiological conditions. Functional roles of gap junction channels could be defined by the relative importance of permeant substances, resulting in coordination of electrical and/or metabolic cellular responses. Cells of the native and specific immune systems establish transient homo- and heterocellular contacts at various steps of the immune response. Morphological and functional studies reported during the last three decades have revealed that many intercellular contacts between cells in the immune response present gap junctions or "gap junction-like" structures. Partial characterization of the molecular composition of some of these plasma membrane structures and regulatory mechanisms that control them have been published recently. Studies designed to elucidate their physiological roles suggest that they might permit coordination of cellular events which favor the effective and timely response of the immune system. Antitumor suicide gene therapy is one of the emerging strategies against cancer (Mesnil M et. al.; *Cancer Res* 2000 Aug 1;60(15):3989-99). It consists of the introduction into cancer cells of a gene capable of converting a nontoxic prodrug into a cytotoxic drug. Because this therapeutic gene cannot be easily introduced into the whole cell population of a tumor, the successful eradication of tumors depends on a

phenomenon called the "bystander effect," by which the introduced gene can affect even cells in which it is not itself present. From a therapeutic point of view, it may be crucial to enhance this phenomenon through various means to achieve tumor eradication. One such suicide gene, the thymidine kinase gene from the herpes simplex virus, in combination with the prodrug ganciclovir, has been extensively and successfully used in some animal models exhibiting a strong bystander effect. Among the mechanisms involved in this phenomenon, gap junctional intercellular communication (GJIC) is directly involved in the transfer of the toxic metabolites of ganciclovir, which pass directly from herpes simplex virus thymidine kinase-expressing cells to surrounding cells that do not express it. Because GJIC appears to be a mediator of the bystander effect both in vitro and in vivo, here we review possible molecular strategies for enhancing the extent of tumor cell death by increasing the intratumoral GJIC capacity. Synapses are classically defined as close connections between two nerve cells or between a neuronal cell and a muscle or gland cell across which a chemical signal (i.e., a neurotransmitter) and/or an electrical signal (i.e., current-carrying ions) can pass (Rozental et.al.; Brain Res Brain Res Rev 2000 Apr;32(1):11-5). The definition of synapse was developed by Charles Sherrington and by Ramon y Cajal at the beginning of this century and refined by John Eccles and Bernard Katz 50 years later; in this collection of papers, the definition of synapses is discussed further in the chapter by Mike Bennett. who provided the first functional demonstration of electrical transmission via gap junction channels between vertebrate neurons. As is evidenced by the range of topics covered in this issue, research dealing with gap junctions in the nervous system has expanded enormously in the past decade, major findings being that specific cell types in the brain expresses specific types of connexins and that expression patterns coincide with tissue compartmentalization and function and that these compartments change during development. Connexins, the protein molecules forming gap junction channels, are reduced in number or redistributed from intercalated disks to lateral cell borders in a variety of cardiac diseases (Jongsma HJ et.al.; Circ Res 2000 Jun 23;86(12):1193-7). This "gap junction remodeling" is considered to be arrhythmogenic. Using a simple model of human ventricular myocardium, we found that quantitative remodeling data extracted from the literature gave rise to only small to moderate changes in conduction velocity and the anisotropy ratio. Especially for longitudinal conduction, cytoplasmic resistivity (and thus cellular geometry) is much more important than commonly realized. None of the remodeling data gave rise to slow conduction on the order of a few centimeters per second. Physical signals, in particular mechanical loading, are clearly important regulators of bone turnover (Donahue HJ; Bone 2000 May;26(5):417-22). Indeed, the structural success of

the skeleton is due in large part to the bone's capacity to recognize some aspect of its functional environment as a stimulus for achievement and retention of a structurally adequate morphology. However, while the skeleton's ability to respond to its mechanical environment is widely accepted, identification of a reasonable mechanism through which a mechanical "load" could be transformed to a signal relevant to the bone cell population has been elusive. In addition, the downstream response of bone cells to load-induced signals is unclear. In this work, we review evidence suggesting that gap junctional intercellular communication (GJIC) contributes to mechanotransduction in bone and, in so doing, contributes to the regulation of bone cell differentiation by biophysical signals. In this context, mechanotransduction is defined as transduction of a load-induced biophysical signal, such as fluid flow, substrate deformation, or electrokinetic effects, to a cell and ultimately throughout a cellular network. Thus, mechanotransduction would include interactions of extracellular signals with cellular membranes, generation of intracellular second messengers, and the propagation of these messengers, or signals they induce, through a cellular network. We propose that gap junctions contribute largely to the propagation of intracellular signals.

The disclosed NOV7 nucleic acid of the invention encoding a Gap Junction Beta-5 Protein-like protein includes the nucleic acid whose sequence is provided in Table 7A or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 7A while still encoding a protein that maintains its Gap Junction Beta-5 Protein-like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 16% percent of the bases may be so changed.

The disclosed NOV7 protein of the invention includes the Gap Junction Beta-5 Protein-like protein whose sequence is provided in Table 7B. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 7B while still encoding a protein that maintains its Gap Junction Beta-

5 Protein-like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 33% percent of the residues may be so changed.

The protein similarity information, expression pattern, and map location for the Gap Junction Beta-5 Protein-like protein and nucleic acid (NOV7) disclosed herein suggest that NOV7 may have important structural and/or physiological functions characteristic of the Gap Junction Beta-5 Protein-like family. Therefore, the NOV7 nucleic acids and proteins of the invention are useful in potential diagnostic and therapeutic applications. These include serving as a specific or selective nucleic acid or protein diagnostic and/or prognostic marker, wherein the presence or amount of the nucleic acid or the protein are to be assessed, as well as potential therapeutic applications such as the following: (i) a protein therapeutic, (ii) a small molecule drug target, (iii) an antibody target (therapeutic, diagnostic, drug targeting/cytotoxic antibody), (iv) a nucleic acid useful in gene therapy (gene delivery/gene ablation), and (v) a composition promoting tissue regeneration in vitro and in vivo.

The NOV7 nucleic acids and proteins of the invention are useful in potential diagnostic and therapeutic applications implicated in various diseases and disorders described below and/or other pathologies. For example, the compositions of the present invention will have efficacy for treatment of patients suffering from Deafness, autosomal dominant 2; Elliptocytosis-1; Fucosidosis; Hypophosphatasia (adult, childhood, infantile); Muscle-eye-brain disease; Neuropathy, paraneoplastic sensory; Porphyria cutanea tarda; Porphyria, hepatoerythropoietic; Schwartz-Jampel syndrome; Thrombocytopenia, congenital amegakaryocytic; Charcot-Marie-Tooth neuropathy-2A; Galactose epimerase deficiency; Glucose transport defect, blood-brain barrier; Kostmann neutropenia; Muscular dystrophy, congenital, with early spine rigidity; Myopathy due to succinate dehydrogenase deficiency; SCID due to LCK deficiency; Colorectal cancer, resistance to; Bartter syndrome, type 3; Breast cancer, ductal; Corneal dystrophy, crystalline, Schnyder; Hyperprolinemia, type II; Inflammatory bowel disease 7; Malignant melanoma, cutaneous; Neuroblastoma; Prostate cancer-brain cancer susceptibility; erythrokeratoderma variabilis; palmoplantar keratoderma; diseases and disorders involving intercellular metabolic and electrical communication; diseases and disorders involving coordination, proliferation and differentiation; diseases and disorders involving maintenance of tissue homeostasis, growth control, development, and synchronized response of cells to stimuli; diseases and disorders involving the immune system; diseases and disorders involving regulation of bone cell differentiation, and/or other pathologies/disorders. The NOV7 nucleic acid, or fragments thereof, may further be useful in

diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

NOV7 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immunospecifically to the novel substances of the invention for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. For example the disclosed NOV7 protein have multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, contemplated NOV7 epitope is from about amino acids 40 to 70. In other embodiments, NOV7 epitope is from about amino acids 90 to 140, from about amino acids 170 to 180, or from about amino acids 220 to 255. This novel protein also has value in development of powerful assay system for functional analysis of various human disorders, which will help in understanding of pathology of the disease and development of new drug targets for various disorders.

## NOV8

A disclosed NOV8 nucleic acid of 546 nucleotides (also referred to as 56072181\_da1) encoding a novel MT-like protein is shown in Table 8A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 214-216 and ending with a TGA codon at nucleotides 397-399. A putative untranslated region upstream from the initiation codon and downstream from the termination codon is underlined in Table 8A. The start and stop codons are in bold letters.

**Table 8A. NOV8 nucleotide sequence (SEQ ID NO:21).**

CAATCTGACATGTAACTGTAAAATCACTGGGCCCAGGAGTGACTGTAATTAAGGGCTTACGTTGAAAATGT  
ATAACAGAGCTTGATTCTTAAGTTTCGAAAATCTTCACAGTGATGGTGTGAACTAAATCCACAGCCACAT  
AATAATCAAAACAGAAAAGCAAAAGAAAGCCACTTCAGGAAATACCACGGTCACACCTCAGTTGGCTTCATG  
CTACAGATTATAGAAAATATGTTGCTGCCCGGGCCACCAATCTGTTGGTTCACATTACTACGTGAGCAATGT  
AAGTGTGCAAGAAGCCATCCACTATCTAAATATCAGATATAGATGCTCCAAAGCAGCTACGTCAGTGATG  
AGAACAGAGAAAATACGTAGCAACATTTCAATTAAGTTGAATTCTAATACTTAAAGGCTCCTTTTAGTACTG  
ACATTCTGGATTTTAAAAGTTATGTTGACCGCATGTTCTCACTCACAAGTGGGAGTTGAACAATGAGAACAC  
ACGGACACGGGGAAGGGAACATCACACACCAGGGCCTGTCAG

The NOV8 nucleic acid sequence is located on the q13 region of chromosome 16.

The disclosed NOV8 polypeptide (SEQ ID NO:22) encoded by SEQ ID NO:21 has 61 amino acid residues and is presented in Table 8B using the one-letter amino acid code. Signal P, Psort and/or Hydropathy results predict that NOV8 has no signal peptide and is likely to be localized in the cytoplasm with a certainty of 0.6500. In other embodiments, NOV8 may also



be localized to the mitochondrial matrix space with a certainty of 0.3100, or the lysosome (lumen) with a certainty of 0.1000.

**Table 8B. Encoded NOV8 protein sequence (SEQ ID NO:22).**

MLQIIENMLLPGPPICWFTLLREQCKCLQEAIHYLNIRYRCSKAATSVMRTEKIRSNISLS

5 A search of sequence databases reveals that the NOV8 amino acid sequence has 16 of 41 amino acid residues (39%) identical to, and 25 of 41 amino acid residues (60%) similar to, the 48 amino acid residue Metallothionein protein from *Rhizomucor racemosus* (*Mucor circinelloides f. lusitanicus*) (SPTREMBL-ACC: Q9Y762) (E = 0.049).

SNP data for NOV8 can be found below in Example 3.

10 Masters et al. (Proc. Nat. Acad. Sci. 91: 584-588, 1994) described metallothioneins (MTs) as a family of low molecular weight, heavy metal-binding proteins characterized by a high cysteine content and lack of aromatic amino acids. MTs bind 7 to 12 heavy metal atoms per molecule of protein. They are ubiquitous in the animal and plant kingdoms and are found in prokaryotes. In mammals, the cysteine residues are absolutely conserved and serve to  
15 coordinate heavy metal atoms such as zinc, cadmium, and copper via mercaptide linkages. In human liver, MTs occur in 2 major forms, MT-I and MT-II (156360). In HeLa cells, MT synthesis is induced by either ionized zinc or ionized cadmium and by glucocorticoid hormones. In man, metallothioneins are encoded by at least 10 to 12 genes separated into 2 groups designated MT-I and MT-II. Masters et al. (1994) noted that, unlike MT-I and MT-II,  
20 which are expressed in most organs, MT-III (139255) expression appears to be restricted to the brain, and MT-IV is only expressed in certain stratified squamous epithelia.

MTs have been postulated to detoxify metals; to play a role in zinc and copper homeostasis during development; to regulate synthesis, assembly, or activity of zinc metalloproteins; and to protect against reactive oxygen species. MTs may also protect against  
25 copper toxicity in the Menkes disease (309400) and murine 'Mottled' phenotypes (X-linked diseases resulting in copper deficiency) as well as in Wilson disease (277900); see also the Animal Models section.

Karin and Richards (Nature 299: 797-802, 1982) described the molecular cloning and sequence analysis of human metallothionein transcripts. Karin et al. (1984) characterized DNA  
30 sequences that are involved in the induction of MT gene expression by cadmium and glucocorticoids.

## MAPPING

Karin et al. (Proc. Nat. Acad. Sci. 81: 5494-5498, 1984) used several different hybridization probes derived from cloned and functional human MT1 and MT2 genes to map the genes in somatic cell hybridization studies. They concluded that most of the human genes are clustered on chromosome 16. Analysis of RNA from somatic cell hybrids indicated that all hybrids that contain human chromosome 16 express both MT1 and MT2 mRNA and that expression is regulated by both heavy metal ions and glucocorticoid hormones.

In the mouse, the metallothionein genes are on chromosome 8, which has other homology to human chromosome 16; by somatic cell hybridization, Cox and Palmiter (Hum. Genet. 64: 61-64, 1983) assigned the MT-I structural gene to mouse chromosome 8, which also carries glutathione reductase in the mouse. (By chance the human 8 also carries glutathione reductase.)

Schmidt et al. (Science 224: 1104-1106, 1984) concluded that MT1 is located between PGP (172280) and DIA4 (125860) and is probably on the long arm 16cen-16q21 because APRT (102600), a 16q marker, and MT1 are both on mouse chromosome 8, whereas HB alpha (141800), a 16p marker, is on mouse chromosome 11. They stated that analysis of the involvement of the MT genes in Wilson disease (277900) and in acrodermatitis enteropathica (201100) would be of great interest.

By gel transfer hybridization analysis of the DNA from human-rodent cell hybrids, Schmidt et al. (J. Biol. Chem. 260: 7731-7737, 1985) showed that chromosome 16 contains a cluster of metallothionein sequences, including 2 functional metallothionein I genes (156351 and 156352) and a functional metallothionein II gene. The remaining sequences, including a processed pseudogene, are dispersed to at least 4 other autosomes. The absence of metallothionein sequences from the X chromosome indicates that the Menkes disease mutation affects metallothionein expression by a 'trans-acting' mechanism. The processed pseudogene is on chromosome 4 and shows allelic variation (Karin and Richards, Nucleic Acids Res. 10: 3165-3173, 1982). Two MT genes are on chromosome 1 but not close together: one is on the distal two-thirds of the short arm and the second probably on the long arm. One metallothionein gene is on chromosome 20 and another is on chromosome 18.

By in situ hybridization, Le Beau et al. (Nature 313: 709-711, 1985) assigned the metallothionein gene cluster to 16q22. This band is a breakpoint in 2 specific rearrangements, inv(16)(p13q22) and t(16;16)(p13;q22), found in a subgroup of patients with acute myelomonocytic leukemia. Hybridization of an MT probe to malignant cells from patients

with one or the other of these rearrangements showed that the breakpoint at 16q22 splits the MT gene cluster. The findings were interpreted as indicating that the MT genes or their regulatory regions may function as an 'activating' sequence for an as yet unidentified cellular gene located at 16p13. Band 16p22 carries 2 fragile sites: the rare FRA16B and the common FRA16C. Simmers et al. (Science 236: 92-94, 1987) showed that the specific leukemic break that is situated in the metallothionein gene cluster lies proximal to both fragile sites; therefore, neither of these fragile sites could have played a role in the breakage.

Using high-resolution in situ hybridization, Sutherland et al. (Cytogenet. Cell Genet. 51: 1087, 1989, Genomics 6: 144-148, 1990) corrected the mapping of the human metallothionein gene complex to 16q13. They found, furthermore, that the complex is not disrupted by the rearrangement breakpoint on 16q in the patients with myelomonocytic leukemia with abnormal eosinophils, as had previously been reported. They showed that the order is cen--MT--FRA16B--D16S4--inversion breakpoint--HB--qter.

Foster et al. (J. Biol. Chem. 263: 11528-11535, 1988) indicated that 4 functional MT1 genes had been identified and mapped to 16q: MT1A, MT1B (156349), MT1E (156351), and MT1F (156352). They also characterized a fifth MT gene, MT1G (156353). West et al. (Genomics 8: 513-518, 1990) mapped the cluster of MT genes in an 82.1-kb region of 16q13. Of the 14 tightly linked genes, 6 had not previously been described. The mapped genes included the single MT2 gene, MT2A, and at least 2 pseudogenes, MT1C and MT1D. The genes were flanked by the single MT2A gene at one end and a gene labeled MT1X (156359) at the other. The order of genes, beginning at the MT2A end, was 1L--1E--1K--1J--1A--1D--1C--1B--1F--1G--1H--1I. This was also the 5-prime to 3-prime direction of transcription for all the genes except MT1G, which had a tail-to-tail, head-to-head orientation to MT1F and MT1H, respectively.

## ANIMAL MODEL

To test the proposed detoxification and homeostasis functions of mammalian MTs in vivo, Masters et al. (1994) inactivated both alleles of the Mt1 and Mt2 genes in embryonic stem cells and generated mice homozygous for these mutant alleles. These mice were viable and reproduced normally when reared under normal laboratory conditions. They were, however, more susceptible to hepatic poisoning by cadmium. This suggested to Masters et al. (1994) that these widely expressed MTs are not essential for development but do protect against cadmium toxicity.

Human Menkes disease (309400) and the murine 'Mottled' phenotype are X-linked diseases that result from copper deficiency due to mutations in ATP7A, a copper-effluxing ATPase (300011). Male mice with the Mottled-Brindled allele accumulate copper in the intestine, fail to export copper to peripheral organs, and die a few weeks after birth. Much of the intestinal copper is bound by metallothionein. To determine the function of MT in the presence of Atp7a deficiency, Kelly and Palmiter (Nature Genet. 13: 219-222, 1996) crossed Mottled-Brindled females with males that bear a targeted disruption of the Mt1 and Mt2 genes. On the metallothionein-deficient background most Mottled males as well as heterozygous Mottled females died before embryonic day 11. The authors explained the lethality in females by preferential inactivation of the paternal X chromosome in extra embryonic tissues and resultant copper toxicity in the absence of MT.

In support of this hypothesis, Kelly and Palmiter (1996) found that cell lines derived from metallothionein deficient, Mottled embryos were very sensitive to copper toxicity. They concluded that MT is essential to protect against copper toxicity in embryonic placenta, providing a second line of defense when copper effluxers are defective. They also stated that MT probably protects against hepatic copper toxicity in Wilson disease and the LEC rat model in which a similar copper effluxer, ATP7B (277900), is defective, because MT accumulates to high levels in the liver in those diseases.

Beattie et al. (Proc. Nat. Acad. Sci. 95: 358-363, 1998) noted that mice with targeted disruption of the metallothionein-I and metallothionein-II genes were more sensitive to toxic metal and oxidative stress. In addition they were larger than most strains of mice, becoming significantly heavier at age 5 to 6 weeks. At age 14 weeks, the body weight and food intake of MT-null mice was 16 and 30% higher, respectively, compared with control mice. Most 22- to 39-week-old male MT-null mice were obese. Seven-week-old MT-null also had significantly higher levels of plasma leptin (601694) and elevated expression of OB (164160), lipoprotein lipase (238600), and CCAAT enhancer binding protein alpha (189965) genes as compared with age-matched control mice. Abnormal accretion of body fat and adipocyte maturation was initiated at 5 to 7 weeks of age, possibly coincident with sexual maturation. Beattie et al. (1998) concluded that a link between MT and the regulation of energy balance is implied by these observations. They noted the possibility that obesity and the associated biochemical changes in the MT-null mice may be caused by factors other than lack of MT. For example, disruption of MT genes by homologous recombination with DNA containing various modifications may have affected other genes around this locus or may have had downstream effects on gene expression.

The disclosed NOV8 nucleic acid of the invention encoding a MT-like protein includes the nucleic acid whose sequence is provided in Table 8A, or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 8A while still encoding a protein that maintains its MT-like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 10% percent of the bases may be so changed.

The disclosed NOV8 protein of the invention includes the MT-like protein whose sequence is provided in Table 8B. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 2 while still encoding a protein that maintains its MT-like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 61% percent of the residues may be so changed.

The invention further encompasses antibodies and antibody fragments, such as  $F_{ab}$  or  $(F_{ab})_2$ , that bind immunospecifically to any of the proteins of the invention.

The above defined information for this invention suggests that this MT-like protein (NOV8) may function as a member of a "MT family". Therefore, the NOV8 nucleic acids and proteins identified here may be useful in potential therapeutic applications implicated in (but not limited to) various pathologies and disorders as indicated below. The potential therapeutic applications for this invention include, but are not limited to: protein therapeutic, small molecule drug target, antibody target (therapeutic, diagnostic, drug targeting/cytotoxic antibody), diagnostic and/or prognostic marker, gene therapy (gene delivery/gene ablation), research tools, tissue regeneration *in vivo* and *in vitro* of all tissues and cell types composing (but not limited to) those defined here.

The NOV8 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in cancer including but not limited to Inflammation,

Autoimmune disorders, Aging and Cancer. For example, a cDNA encoding the MT-like protein (NOV8) may be useful in gene therapy, and the MT-like protein (NOV8) may be useful when administered to a subject in need thereof. By way of nonlimiting example, the compositions of the present invention will have efficacy for treatment of patients suffering from Gitelman syndrome, Menkes disease, Wilson's disease, acrodermatitis enteropathica, myelomonocytic leukemia, eosinophil disorders, hepatic disorders such as hepatic copper toxicity and other such conditions.. The NOV8 nucleic acid encoding MT-like protein, and the MT-like protein of the invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

NOV8 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immuno-specifically to the novel NOV8 substances for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. The disclosed NOV8 protein has multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated NOV8 epitope is from about amino acids 25 to 40. In another embodiment, a NOV8 epitope is from about amino acids 45 to 55. These novel proteins can be used in assay systems for functional analysis of various human disorders, which will help in understanding of pathology of the disease and development of new drug targets for various disorders.

### NOV9

A disclosed NOV9 nucleic acid of 2309 nucleotides (also referred to as 2855519\_0\_19\_da1) encoding a novel CIP4-like protein is shown in Table 9A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 78-80 and ending with a TGA codon at nucleotides 1719-1721. A putative untranslated region upstream from the initiation codon and downstream from the termination codon is underlined in Table 9A. The start and stop codons are in bold letters.

**Table 9A. NOV9 nucleotide sequence (SEQ ID NO:23).**

GAGTGAGAGGTCGGACAGACTGTGGAGCCGACAGACTGAAGGACAGCGGCACCGCCAGACGGCCAGAAAGTT  
 CCGCCATGAGCTGGGGCACGGAGCTGTGGGATCAGTTTCGACAGCTTAGACAGCATACAAATGGGGAATTG  
 ACTTCTTGGAAAGATATGCCAAATTTGTTAAAGAGAGGATAGAAATGAACAGAACTATGCGAAACAATTGA  
 GAAATCTGGTTAAGAAGTACTGCCCCAAACGTTTCATCCAAAGATGAAGAGCCACGGTTTACCTCGTGTGTAG  
 CCTTTTTTAATATCCTTAATGAGTTAAATGACTATGCAGGACAGCGAGAAGTTGTAGCAGAAGAAATGGCGC  
 ACAGAGTGTATGGTGAATTAATGAGATATGCTCATGATCTGAAACTGAAAGAAAAATGCATCTGCAAGAAG  
 GACGAAAGCTCAACAATATCTTGACATGTGCTGGAAACAGATGGGTAATAGTAAAAAGAAGTTTGAAAGAG  
 AATGTAGAGAGGCAGAAAAGGCACACAGAGTTATGAAAGATTGGATAATGATACTAATGCAACCAAGGCAG  
 ATGTTGAAAATGCCAAACAGCAGTTGAATCTGCGTACGCATATGGCCGATGAAAATAAAAATGCATATGCTG  
 CACAATTACAAAACCTTTAATGGAGAACCAACATAAACATTTTATGTAGTGATTCTCAGATTACAGCAAC  
 TACAAGAAATGGACGAACGAAGGACTATTAACCTCAGTGAGTGTACAGAGGATTTGCTGACTCAGAACGCA  
 AAGTTATTTCCCATCATTTCAAATGTTTGGAAAGGAATGATTCTTGACGAAAATCAGTTGATGAAAGAAGAG  
 ACTCTCAAATGGTGGTAGACTCCTTCAAATCTGGTTTTGAACCTCCAGGAGACTTTCCATTGGAAGATTACA  
 CTCACATATATATAGAACCATTTCTGATGGGACTATCAGTGCATCCAAACAGGAGATGGGAAGATGGATG  
 CCAAAACCCAGTAGGAAAGGCCAAGGGCAAATTTGTGGCTCTTTGGAAAGAGCCAAAGGGCCAGCAGCTAG  
 AAGATTTAGTGCATCTGCCACCAGAACAGAGACGTAAAAAACTACAGCAGCGCATTGTAGAACTTAACAGAG  
 AACTACAGAAAGAAATCAGACCAAAAAGATGCACTCAACAAAATGAAAGATGTATATGAGAAGGATCCACAAA  
 TGGGGGATCCAGGGAGTTTGCAGCCTAAATTAGCAGAGACCATGAATAACATTGACCGCTACGAATGGAAA  
 TCCATAAGAATGAGGCTTGGCTCTCTGAAGTCGAAGGCCAAAACAGGTGGGAGAGGAGACAGAAGACATAGCA  
 GTGACATAAAATCATCTTGTAAACACAGGGACGAGAAAGTCTGAGGGAAGTTACACTGATGATGCAAAACCAGG  
 AAGTCCGTGGGCCACCCAGCAGCATGGTCACCACAATGAGTTTGTATGATGAATTTGAGGATGATGATCCCT  
 TGCCTGCTATTGGACACTGCAAAGCTATCTACCTTTTGTATGGACATAATGAAGGTACTCTAGCAATGAAAG  
 AAGGTGAAGTTCTCTACATTATAGAGGAGGACAAAGGTGACGGATGGACAAGAGCTCGGAGACAGAACGGTG  
 AAGAAGGCTACGTTCCACGTCATACATAGATGTAACCTAGAGAAAAACAGTAAAGGTTCTGAAAGAGGGT  
 TTCTGAGGAAATGGGCAAGATGTTGAAGGAGGTTACATGCAGCTGCTTTTGGGGGAGGGTATTAGAGTTGTC  
 AGGCTCAAAGAGAGTGAGAGAAGCAAGTTGCATGAGTGCATGCAGACATGATTTTTTTTACTAATTCAT  
 TAGCATTTCCATACATTGTTTTTAAAAATCATAATACCAACCCCTTAAGTTCCTAGTTCACAGTTATTTCCAC  
 AAAAGAAAAGCCAAATAGTGTACCATTTTCTATTTTATTGCTGTCTAATCAATAAAGAATGCA  
 GAGCTGTGCAAAAATGTGTCTTACATTAGCTGTCCCAACAGGATTGTCTTCCCTCCAGCTCTGGTTTTAA  
 TTGGCTTTTAGACCCACTATCTGTGATCCTTGCCATCTGTGAGTGTCTGCTGCGCCACCTCCGTGCTTG  
 CCTAACATCCTGTGTCATGTCTAGCGTGATTGAGCNAGATTTTCAGGCATGTCTTTAGAATCCCTGGTINCT  
 GTCAAAGCCTGGTTTGGTTTACATTGGTNGTGAATCNCTTTGTCAACATCTCCAGCACTATNGTTCCNTCT  
 TAGGT

The disclosed NOV9 nucleic acid sequence, localized to the p21.2-22.2 region of chromosome 1, has 916 of 1460 bases (62%) identical to a 2001 bp cdc42-interacting protein 4 (CIP4 mRNA from Homo sapiens (GENBANK-ID: HSCIP4|acc:AJ000414) ( $E = 1.3e^{-97}$ ).

The disclosed NOV9 polypeptide (SEQ ID NO:24) encoded by SEQ ID NO:23 has 547 amino acid residues and is presented in Table 9B using the one-letter amino acid code. Signal P, Psort and/or Hydropathy results predict that NOV9 has no signal peptide and is likely to be localized in the nucleus with a certainty of 0.7000. In other embodiments, NOV9 may also be localized to the microbody (peroxisome) with a certainty of 0.3000, the mitochondrial matrix space with a certainty of 0.1000, or the lysosome (lumen) with a certainty of 0.1000.

**Table 9B. Encoded NOV9 protein sequence (SEQ ID NO:24).**

MSWGTSLWDQFDSLKDHTQWIDFLERYAKFVKERIEIEQNYAKQLRNLVKKYCPKRSSKDEEP  
 RFTSCVAFFNINLNDYAGQREVVAEEMAHRVYELMRYAHD LKTERKMHLEGRKAQQYLDM  
 CWKQMGNSKKKFERECREAEKAQQSYERLDNDTNATKADVENAKQQLNLRTHMADENKNYAAQ  
 LQNFNGEQHKHFYVVI PQIYKQLQEMDERRTIKLSERYGFADSERKVIPIISKCLEGMILAAK  
 SVDERRDSQMVVDSFKSGFEPFGDFPFEDYSQHIYRTISDGTISASKQESGKMDAKTPVGKAKG

KLWLFGKKPKGPALEDFSHLPPEQRRKKLQQRIDELNRELQKESDQKDALNKMKDVEKDPQMG  
 DPGSLQPKLAETMNNIDRLRMEIHKNEAWLSEVEGKTGGRGDRRHSSDINHLVTQGRESPEGSY  
 TDDANQEVRGPPQQHGHNEFDDEFEDDDPLPAIGHCKAIYPFDGHNEGTLMKEGEVLYIIEE  
 DKGDGWTRARRQNGEEGYVPTSIDVTLEKNSKGS

A search of sequence databases reveals that the NOV9 amino acid sequence has 303 of 544 amino acid residues (55%) identical to, and 403 of 544 amino acid residues (74%) similar to, the 545 amino acid residue CDC42-Interacting Protein 4 protein from *Homo sapiens* (SPTREMBL-ACC:O15184) ( $E = 8.0e^{-160}$ ).

TaqMan data for NOV9 can be found below in Example 2. The disclosed NOV9 polypeptide has homology to the amino acid sequences shown in the BLASTP data listed in Table 9C.

Table 9C. BLAST results for NOV9					
Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 13591536 emb CAC36351.1  (AL109613)	dJ1033H22.1 (KIAA0554 protein) [Homo sapiens]	434	373/430 (86%)	375/430 (86%)	0.0
gi 8923249 ref NP_060207.1	hypothetical protein FLJ20275 [Homo sapiens]	330	328/330 (99%)	329/330 (99%)	e-175
gi 10435680 dbj BAB14638.1  (AK023681)	unnamed protein product [Homo sapiens]	592	319/595 (53%)	432/595 (71%)	e-160
gi 13936547 gb AAK49824.1 AF265550_1 (AF265550)	formin-binding protein 17 [Homo sapiens]	679	307/624 (49%)	422/624 (67%)	e-148
gi 3043632 dbj BAA25480.1  (AB011126)	KIAA0554 protein [Homo sapiens]	674	307/624 (49%)	422/624 (67%)	e-148

The homology between these and other sequences is shown graphically in the ClustalW analysis shown in Table 9D. In the ClustalW alignment of the NOV9 protein, as well as all other ClustalW analyses herein, the black outlined amino acid residues indicate regions of conserved sequence (*i.e.*, regions that may be required to preserve structural or functional properties), whereas non-highlighted amino acid residues are less conserved and can potentially be altered to a much broader extent without altering protein structure or function.

Table 9D. ClustalW Analysis of NOV9

1) Novel NOV9 (SEQ ID NO:24)  
 3) gi|13591536|emb|CAC36351.1| (AL109613) dJ1033H22.1 (KIAA0554 protein) [Homo sapiens] (SEQ ID NO:66)  
 4) gi|8923249|ref|NP\_060207.1| hypothetical protein FLJ20275 [Homo sapiens] (SEQ ID NO:67)



- 5) gi|10435680|dbj|BAB14638.1| (AK023681) unnamed protein product [Homo sapiens] (SEQ ID NO:68)  
 6) gi|13936547|gb|AAK49824.1|AF265550\_1 (AF265550) formin-binding protein 17 [Homo sapiens] (SEQ ID NO:69)  
 5 7) gi|3043632|dbj|BAA25480.1| (AB011126) KIAA0554 protein [Homo sapiens] (SEQ ID NO:70)

```

      10      20      30      40
10  NOV9      . . . . .
    gi|13591536|-----
    gi|8923249|-----
    gi|10435680|-----
    gi|13936547|LWNGGEEEEPPRRPRARSCEPEEAARTPGFPPSRGSRGAKG
15  gi|3043632|-----EEEPRRPRARSCEPEEAARTPGFPPSRGSRGAKG

      50      60      70      80
20  NOV9      . . . . .MSWGTTELWDQFDL L KHT
    gi|13591536|-----
    gi|8923249|-----
    gi|10435680|-----MSWGTTELWDQFDL L KHT
    gi|13936547|SPGRGTREPRPPRGAPLRVPCTMSWGTTELWDQFDL L KHT
25  gi|3043632|SPGRGTREPRPPRGAPLRVPCTMSWGTTELWDQFDL L KHT

      90     100     110     120
30  NOV9      QWGIDILEYIKFVKERTEIELSYAKQLRNLSKKYQPKIN
    gi|13591536|-----
    gi|8923249|-----
    gi|10435680|QWGIDILEYIKFVKERTEIELSYAKQLRNLSKKYQPKIN
    gi|13936547|QWGIDILEYIKFVKERTEIELSYAKQLRNLSKKYQPKIN
35  gi|3043632|QWGIDILEYIKFVKERTEIELSYAKQLRNLSKKYQPKIN

      130     140     150     160
40  NOV9      SKDEE-PRITSCVAFFN LNE NDYAGQHEVISENMASQI
    gi|13591536|-----
    gi|8923249|-----
    gi|10435680|SKDEEYKYTSCKAFISNLNE NDYAGQHEVISENMASQI
    gi|13936547|SKDEEYKYTSCKAFISNLNE NDYAGQHEVISENMASQI
45  gi|3043632|SKDEEYKYTSCKAFISNLNE NDYAGQHEVISENMASQI

      170     180     190     200
50  NOV9      YGELMRVAHDLKTERKMELOGRKAQQHITCCKWQNGESK
    gi|13591536|-----
    gi|8923249|-----
    gi|10435680|IVDLARYVQLKQERKSFPHDGRKAQQHITCCKWQLESSEK
    gi|13936547|IVDLARYVQLKQERKSFPHDGRKAQQHITCCKWQLESSEK
55  gi|3043632|IVDLARYVQLKQERKSFPHDGRKAQQHITCCKWQLESSEK

      210     220     230     240
60  NOV9      RFEERDC EADRAQQYTEK DADINVTKADVEKARQQAQI
    gi|13591536|-----
    gi|8923249|-----
    gi|10435680|RFEERDC EADRAQQYTEK DADINVTKADVEKARQQAQI
    gi|13936547|RFEERDC EADRAQQYTEK DADINVTKADVEKARQQAQI
65  gi|3043632|RFEERDC EADRAQQYTEK DADINVTKADVEKARQQAQI

      250     260     270     280
70  NOV9      RTHMADEKNAYAAQLQNFCEQHKHYVVIPQITKQIQE
    gi|13591536|-----
    gi|8923249|-----
    gi|10435680|RTHMADEKNAYAAQLQNFCEQHKHYVVIPQITKQIQE
    gi|13936547|RTHMADEKNAYAAQLQNFCEQHKHYVVIPQITKQIQE
    gi|3043632|RTHMADEKNAYAAQLQNFCEQHKHYVVIPQITKQIQE

```

290 300 310 320

NOV9  
gi|13591536|  
gi|8923249|  
gi|10435680|  
gi|13936547|  
gi|3043632|

5

330 340 350 360

NOV9  
gi|13591536|  
gi|8923249|  
gi|10435680|  
gi|13936547|  
gi|3043632|

10

15

370 380 390 400

NOV9  
gi|13591536|  
gi|8923249|  
gi|10435680|  
gi|13936547|  
gi|3043632|

20

25

410 420 430 440

NOV9  
gi|13591536|  
gi|8923249|  
gi|10435680|  
gi|13936547|  
gi|3043632|

30

35

450 460 470 480

NOV9  
gi|13591536|  
gi|8923249|  
gi|10435680|  
gi|13936547|  
gi|3043632|

40

45

490 500 510 520

NOV9  
gi|13591536|  
gi|8923249|  
gi|10435680|  
gi|13936547|  
gi|3043632|

50

530 540 550 560

NOV9  
gi|13591536|  
gi|8923249|  
gi|10435680|  
gi|13936547|  
gi|3043632|

55

60

570 580 590 600

NOV9  
gi|13591536|  
gi|8923249|  
gi|10435680|  
gi|13936547|

65

70

```

gi|3043632| SGLYDSQNPPTVNCADRESPGSGYTFQESQES----EM
          610      620      630      640
5  NOV9      HGHHNFDDEFDDPLPAIGHCKAYPFQGHNEGTAAK
   gi|13591536| HGHHNFDDEFDDPLPAIGHCKAYPFQGHNEGTAAK
   gi|8923249| HGHHNFDDEFDDPLPAIGHCKAYPFQGHNEGTAAK
   gi|10435680| KVLATFDDEFDDPLPAIGHCKAYTFQGHNEGTAAV
10 gi|13936547| KVLATFDDEFDDPLPAIGHCKAYTFQGHNEGTAAV
   gi|3043632| KVLATFDDEFDDPLPAIGHCKAYTFQGHNEGTAAV
          650      660      670      680
15 NOV9      EGEVLYIEEDKGDGWTRARRQNGEEGVPTSYYVTLAK
   gi|13591536| EGEVLYIEEDKGDGWTRARRQNGEEGVPTSYYVTLAK
   gi|8923249| EGEVLYIEEDKGDGWTRARRQNGEEGVPTSYYVTLAK
   gi|10435680| EGETLYIEEDKGDGWTRIRRNDEEGYVPTSYYVCLAK
   gi|13936547| EGETLYIEEDKGDGWTRIRRNDEEGYVPTSYYVCLAK
20 gi|3043632| EGETLYIEEDKGDGWTRIRRNDEEGYVPTSYYVCLAK
          .....
25 NOV9      NKGSG----
   gi|13591536| NKGIVTYI
   gi|8923249| NKGSG----
   gi|10435680| NKGAKTYI
   gi|13936547| NKGDS----
   gi|3043632| NKGDS----
          .....
30

```

Table 9E-H lists the domain description from DOMAIN analysis results against NOV9. This indicates that the NOV9 sequence has properties similar to those of other proteins known to contain this domain.

**Table 9E. Domain Analysis of NOV9**

gnl|Smart|smart00326, SH3, Src homology 3 domains; Src homology 3 (SH3) domains bind to target proteins through sequences containing proline and hydrophobic amino acids. Pro-containing polypeptides may bind to SH3 domains in 2 different binding orientations. (SEQ ID NO:92)  
 CD-Length = 59 residues, 88.1% aligned  
 Score = 64.7 bits (156), Expect = 1e-11

```

Query:  484  HCKAIYPFDGHNEGTAMKEGEVLYIIIEDKGDGWTRARRQNGEEGVPTSYYI  536
          +|+| +  +  |+|+|+ +|+  ||| + |  |+|  |++|+
Sbjct:  4    QVRALYDYTAQDPDELSFKKGDIIITVLEKS-DDGWWKGR LGTGKEGLFPSNYV  55

```

### Table 9F. Domain Analysis of NOV9

gnl|Pfam|pfam00018, SH3, SH3 domain. SH3 (Src homology 3) domains are often indicative of a protein involved in signal transduction related to cytoskeletal organization. First described in the Src cytoplasmic tyrosine kinase. The structure is a partly opened beta barrel. (SEQ ID NO:93)  
CD-Length = 57 residues, 91.2% aligned  
Score = 63.5 bits (153), Expect = 3e-11

Query: 486 KAIYPFDGHNEGTLMKEGEVLYIIIEEDKGDGWTRARRQNGEEGYVPTSID 537  
A+Y + L+ K+G+++ ++E+ GW + R + +EG +P++Y++  
Sbjct: 4 VALYDYQARESEDELSFKKGDIIIVLEKSDDGGWWKGRLLKGTKEGLIPSNYVE 55

### Table 9G. Domain Analysis of NOV9

gnl|Smart|smart00055, FCH, Fes/CIP4 homology domain; Alignment extended from original report. Highly alpha-helical. Also known as the RAEYL motif or the S. pombe Cdc15 N-terminal domain. (SEQ ID NO:94)  
CD-Length = 91 residues, 97.8% aligned  
Score = 58.2 bits (139), Expect = 1e-09

Query: 1 MSWGTELWDQFDSLDKHTQWGIDFLERYAKFVKERIEIEQNYAKQLRNLVKKYCPKRSSK 60  
| + +|| | ++| + | || ||++|| +||+ |||+| + || +  
Sbjct: 1 MGFWSELDDGFEALLSRLKNGRLLEDLKKFMRERAKIEEYAKKLQKLSKK--LRAVRD 58  
Query: 61 DEEPRFTSCVAFFNILNELNDYAGQREVVAE 91  
| + | + | + | + | + | ++|  
Sbjct: 59 TESELGSLRKAWEVLLSETDALAKQHLQLSE 89

### Table 9H. Domain Analysis of NOV9

gnl|Pfam|pfam00611, FCH, Fes/CIP4 homology domain. Alignment extended from. Highly alpha-helical. (SEQ ID NO:95)  
CD-Length = 94 residues, 97.9% aligned  
Score = 40.0 bits (92), Expect = 3e-04

Query: 1 MSWGTELWDQ-FDSLDKHTQWGIDFLERYAKFVKERIEIEQNYAKQLRNLVKKYCPKRSS 59  
| +|+|| + +| + || ||+ || +||+ || +|++| + ++  
Sbjct: 1 MGFGSELCPEGHKALLSRQDNELRLLEEMKKFMAERAKIEKEYAGKLOHLSAQVGKGPAT 60  
Query: 60 KDEEPRFTSCVAFFNILNELNDYAGQREVVAE 91  
+ | +| ++ ||+| + ++|  
Sbjct: 61 AGEDELSSLKSWAVILSETEQQSKIHLQISE 92

The thyroid hormone receptors (TRs) are hormone-dependent transcription factors that regulate expression of a variety of specific target genes. They must specifically interact with a number of proteins as they progress from their initial translation and nuclear translocation to heterodimerization with retinoid X receptors (RXRs), functional interactions with other

transcription factors and the basic transcriptional apparatus, and eventually, degradation. To help elucidate the mechanisms that underlie the transcriptional effects and other potential functions of TRs, Lee et al. (Molec. Endocr. 9: 243-254, 1995) used the yeast interaction trap, a version of the yeast 2-hybrid system, to identify proteins that specifically interact with the  
5 ligand-binding domain of rat TR-beta (THRB; 190160). They isolated HeLa cell cDNAs encoding several different TR-interacting proteins (TRIPs), including TRIP10. TRIP10 interacted with rat Thrb only in the presence of thyroid hormone. It also showed a ligand-dependent interaction with RXR-alpha (RXRA; 180245), but did not interact with the glucocorticoid receptor (NR3C1; 138040) under any condition. TRIP10 contains a Src  
10 homology-3 (SH3) domain and shows sequence similarity to a Xenopus fyn homolog (see 137025) and chicken src (see 190090). Northern blot analysis detected a 2.6-kb TRIP10 transcript in several tissues, with highest expression in skeletal muscle.

Using the yeast 2-hybrid system to identify proteins that bind to a constitutively active mutant of CDC42 (116952), Aspenstrom (1997) isolated a human B-cell cDNA encoding  
15 CDC42-interacting protein-4 (CIP4), which is identical to TRIP10. The predicted 545-amino acid CIP4 protein contains an N-terminal domain that bears resemblance to the nonkinase domain of the FER (176942) and Fes/Fps family of tyrosine kinases, and a C-terminal SH3 domain. In addition, CIP4 shares sequence similarity with a number of proteins that have roles in regulating the actin cytoskeleton. Aspenstrom (Curr. Biol. 7: 479-487, 1997) demonstrated  
20 that CIP4 can bind to activated CDC42 in vitro and in vivo. Overexpression of CIP4 in fibroblasts reduced the amount of stress fibers in these cells. Recombinant CIP4 protein accumulated at the cell periphery, particularly in areas that exhibited membrane ruffling. Coexpression of activated CDC42 and CIP4 led to clustering of CIP4 to a large number of foci at the dorsal side of the cells. Northern blot analysis showed a major 2.2-kb CIP4 transcript  
25 that was abundant in skeletal muscle, heart, and placenta, present at lower levels in pancreas, lung, liver, and kidney, and barely detectable in brain. Minor transcripts of 3.5 and 5 kb were also detected.

Aspenstrom (1997) concluded that CIP4 is a downstream target of activated GTP-bound CDC42, and is similar in sequence to proteins involved in signaling and cytoskeletal  
30 control. The author suggested that CIP4 may act as a link between CDC42 signaling and regulation of the actin cytoskeleton.

The Wiskott-Aldrich syndrome is an inherited X-linked immunodeficiency characterized by thrombocytopenia, eczema, and a tendency toward lymphoid malignancy (Tian L, et.al.; J Biol Chem 2000 Mar 17;275(11):7854-61). Lymphocytes from affected

individuals have cytoskeletal abnormalities, and monocytes show impaired motility. The Wiskott-Aldrich syndrome protein (WASP) is a multi-domain protein involved in cytoskeletal organization. In a two-hybrid screen, we identified the protein Cdc42-interacting protein 4 (CIP4) as a WASP interactor. CIP4, like WASP, is a Cdc42 effector protein involved in cytoskeletal organization. We found that the WASP-CIP4 interaction is mediated by the binding of the Src homology 3 domain of CIP4 to the proline-rich segment of WASP. Cdc42 was not required for this interaction. Co-expression of CIP4 and green fluorescent protein-WASP in COS-7 cells led to the association of WASP with microtubules. In vitro experiments showed that CIP4 binds to microtubules via its NH(2) terminus. The region of CIP4 responsible for binding to active Cdc42 was localized to amino acids 383-417, and the mutation I398S abrogated binding. Deletion of the Cdc42-binding domain of CIP4 did not affect the colocalization of WASP with microtubules in vivo. We conclude that CIP4 can mediate the association of WASP with microtubules. This may facilitate transport of WASP to sites of substrate adhesion in hematopoietic cells.

The disclosed NOV9 nucleic acid of the invention encoding a CIP4-like protein includes the nucleic acid whose sequence is provided in Table 9A, or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 9A while still encoding a protein that maintains its CIP4-like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 38% percent of the bases may be so changed.

The disclosed NOV9 protein of the invention includes the CIP4-like protein whose sequence is provided in Table 9B. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 2 while still encoding a protein that maintains its CIP4-like activities and physiological functions, or a

functional fragment thereof. In the mutant or variant protein, up to about 51% percent of the residues may be so changed.

The invention further encompasses antibodies and antibody fragments, such as  $F_{ab}$  or  $(F_{ab})_2$ , that bind immunospecifically to any of the proteins of the invention.

5       The above defined information for this invention suggests that this CIP4-like protein (NOV9) may function as a member of a "MT family". Therefore, the NOV9 nucleic acids and proteins identified here may be useful in potential therapeutic applications implicated in (but not limited to) various pathologies and disorders as indicated below. The potential therapeutic applications for this invention include, but are not limited to: protein therapeutic, small  
10   molecule drug target, antibody target (therapeutic, diagnostic, drug targeting/cytotoxic antibody), diagnostic and/or prognostic marker, gene therapy (gene delivery/gene ablation), research tools, tissue regeneration *in vivo* and *in vitro* of all tissues and cell types composing (but not limited to) those defined here.

      The NOV9 nucleic acids and proteins of the invention are useful in potential  
15   therapeutic applications implicated in cancer including but not limited to Inflammation, Autoimmune disorders, Aging and Cancer. For example, a cDNA encoding the CIP4-like protein (NOV9) may be useful in gene therapy, and the CIP4-like protein (NOV9) may be useful when administered to a subject in need thereof. By way of nonlimiting example, the compositions of the present invention will have efficacy for treatment of patients suffering  
20   from Wiskott-Aldrich syndrome, immunodeficiency, thrombocytopenia, eczema, lymphoid malignancy cytoskeletal abnormalities, impaired monocyte motility, Muscular dystrophy, Lesch-Nyhan syndrome, Myasthenia gravis, Von Hippel-Lindau (VHL) syndrome, Alzheimer's disease, Stroke, Tuberous sclerosis, hypercalceimia, Parkinson's disease, Huntington's disease, Cerebral palsy, Epilepsy, Lesch-Nyhan syndrome, Ataxia-telangiectasia,  
25   Leukodystrophies, Behavioral disorders, Addiction, Anxiety, Pain, Neuroprotection, Fertility, Cardiomyopathy, Atherosclerosis, Hypertension, Congenital heart defects, Aortic stenosis , Atrial septal defect (ASD), Atrioventricular (A-V) canal defect, Ductus arteriosus, Pulmonary stenosis, Subaortic stenosis, Ventricular septal defect (VSD), valve diseases, Tuberous sclerosis, Scleroderma, Obesity, Transplantation, Diabetes, Pancreatitis, Obesity, Systemic  
30   lupus erythematosus , Autoimmune disease, Asthma, Emphysema, Scleroderma, allergy, ARDS, Cirrhosis, Transplantation, Diabetes, Autoimmune disease, Renal artery stenosis, Interstitial nephritis, Glomerulonephritis, Polycystic kidney disease, Renal tubular acidosis, IgA nephropathy, Hypercalceimia, or other pathologies or conditions. The NOV9 nucleic acid encoding CIP4-like protein, and the CIP4-like protein of the invention, or fragments thereof,

may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

NOV9 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immuno-specifically to the novel NOV9 substances for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. The disclosed NOV9 protein has multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated NOV9 epitope is from about amino acids 25 to 40. In another embodiment, a NOV9 epitope is from about amino acids 45 to 55. These novel proteins can be used in assay systems for functional analysis of various human disorders, which will help in understanding of pathology of the disease and development of new drug targets for various disorders.

#### **NOV10**

NOV10 includes three novel hepsin/plasma transmembrane serine protease-like proteins disclosed below. The disclosed sequences have been named NOV10a and NOV10b.

#### **NOV10a**

A disclosed NOV10a nucleic acid of 1787 nucleotides (also referred to as 129297354\_EXT) encoding a novel hepsin/plasma transmembrane serine protease-like protein is shown in Table 10A. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 54-56 and ending with a TAA codon at nucleotides 1470-1472. A putative untranslated region upstream from the initiation codon and downstream from the termination codon is underlined in Table 10A. The start and stop codons are in bold letters.



**Table 10A. NOV10a nucleotide sequence (SEQ ID NO:25).**

GCGGAACATTGCCTAGTAGACCCTGAGGCTTTACAACAGTGCCACTGACCCCTATGAGCCTGATGCTGGATG  
 ACCAACCCCTATGGAGGCCAGTATGCAGAGGAGGGCCAGGACCTGGGATCTTCAGAGCAGAGCCTGGAG  
 ACCAGCAGCATCCCATTTCTCAGGCGGTGTGCTGGCGTTCCATGCGACGTGGCTGTGCAGTGTGGGAGCCC  
 TGGGGCTGCTGGCCGGTGCAGGTGTTGGCTCATGGCTCCTAGTGTGTATCTGTGTCTGCTGCCTCTCAGC  
 CCATTTCCGGGACCTTGACAGATGAGGAGATAACTTTGAGCTGCTCAGAGGCCAGCGTGAGGAAGCTCTGC  
 TCCCTGCACTTCCCAAAACAGTATCTTTCAGAATAAACAGCGAAGACTTCTTGCTGGAAGCGCAAGTGAGGG  
 ATCAGCCACGCTGGCTCCTGGTCTGCCATGAGGGCTGGAGCCCCGCCCTGGGGCTGCAGATCTGTGGAGCC  
 TTGGGCATCTCAGACTCACTCACCACAAGGGAGTAAACCTCACTGACATCAAACCTCAACAGTTCCAGGAGT  
 TTGCTCAGCTCTCTCCTAGACTGGGAGGCTTCTGGAGGAGGCGTGGCAGCCAGTAGGACTACTGAGGCTG  
 TTAGGAACAACATGCACCTTCTGGTCAAGTTGTTTCCCTCAGATGCTCTGAGTGTGGAGCGAGGCCCTGGCTT  
 CCCGGATAGTTGGTGGGAGTCTGTGGCTCCTGGGCGCTGGCCGTGGCAGGCCAGCGTGGCCCTGGGCTTCC  
 GGCACACGTGTGGGGCTCTGTGCTAGCGCCACGCTGGGTGGTACTGCTGCACATTGTATGCACAGTTTCA  
 GGCTGGCCCGCTGTCCAGCTGGCGGGTTCATGCGGGGTGGTCAAGCCACAGTGCCGTGAGGCCCCACCAAG  
 GGGCTCTGGTGGAGAGGATTATCCACACCCCTCTACAGTGCCAGAAATCATGACTACGACGTGCGCCCTCC  
 TGAGGCTCCAGACCGCTCTCAACTTCTCAGACACTGTGGGCGCTGTGTGCTGCGGCCAAGGAACAGCATT  
 TTCCGAAGGGCTCGCGGTGTGGGTGTCTGGCTGGGGCCACACCCACCCTAGCCATACTTACAGCTCGGATA  
 TGCTCCAGGACACGGTGGTGCCTGCTCAGCACTCAGCTCTGCAACAGCTCTGCGTGTACAGCGGAGCCC  
 TCACCCCCCGCATGCTTTGCGCTGGCTACCTGGACGGAAGGGCTGATGCATGCCAGGGAGATAGCGGGGCC  
 CCCTAGTGTGCCAGATGGGACACATGGCGCTAGTGGGGGTGGTCACTGGGGGGTGGTGGTGGTGGTGGTGGT  
 CCAATCACCCAGGTGTCTACGCCAAGGTAGCTGAGTTTCTGACTGGATCCATGACACTGCTCAGGTGAGTG  
 TGGGGGAGGAGTAGGGCAGGGAGATTCTAAAGGACCTGCCCTCGAATGCAAGGAACCTTACCCTTAGGC  
 CCGGGCCCTGCTGGGACTGGGAGGGTGTAGGACATATTCACAGAGTGAAGTGGAGGAAGAGTGAAGCT  
 TAAACATGGAATCCATTGGATTCTATCAGTTTAAAGGATGAAGTGGGTAAGAGTATGCCTGAGTTGTATCC  
 CAGATCTACCATTTCTGTGTCGACCTTTGGCAAAATTTCAACTTTGTTAAACCTTAAATTTCTGATAATAA  
 CCATGATGGCTACTTATATGCTATTGTTATATGCTATTAAATAAGACCCGTACAATGCC

The reverse complement of NOV10a is shown in Table 10B.

**Table 10B. NOV10a reverse complement sequence (SEQ ID NO:26).**

GGCATTGTACGGGTCTTATTTAATAGCATATAACAATAGCATATAAGTAGCCATCATGGTTATTATCAGGAA  
 ATTAAGGTTTAACAAAGTTAGAAATTTGCCAAAGGTCGACACAGGAAATGGTAGATCTGGGATACAACTCA  
 GGCATACTCTTACCAGTTTCATCTTAACTGATAGAAATCCAATGGATTCCATGTTTAACTTCACTTCTT  
 CCTCCACTCACTCTGGGGAATATGTCTTAGCACCTCCCCAGTCCCCAGCAGGGCCCCGGGCTAAGGGGTAA  
 GGTTCTTGCATTGAGGGCAGGTCTTTAGAAATCTCCCTGCCCTACTCCTGCCCCCACTCACCTGAGC  
 AGTGTCACTGGATCCAGTCCAGAACTCAGTACCTTGGCGTAGACACCTGGGTGATTGGGCTCTGCGCAGCC  
 ACGCCCCAGCTGACCAACCCCACTAGGCGCCATGTGTCCCATCTGGGCACACTAGGGGGCCCCCGCTATC  
 TCCCTGGCATGCATCAGCCCTTCCGTCCAGGTAGCCAGCGCAAAGCATGCGGGGGGTGAGGGCTCCGCTGTA  
 CACGCAAGAGCTGTTGCAGAGCTGAGTGTGAGCAGGGGCACCACCGTGTCTGGAGCATATCCGAGCTGTA  
 AGTATGGCTAGGGTGGGTGCGCCCCAGCCAGACCCAGCACCAGCGAGCCCTTCGGAAAATGCTGTTCTCTT  
 GGCCGGCAGGCACACAGCGCCCCACAGTGTCTGAGAAATTGAGAGCGGTCTGGAGCCTCAGGAGGGCGACGTC  
 GTAGTCATGATTCTGGGCACTGTAGAGGGGTGTGGGATAATCCTCTCCACCAGAGCCCTTGGTGGGGCT  
 GACGGCACTGTGGCTGACCAACCCCGCATGAACCCGCCAGCTGGACAGGCGGGCCAGCCTGAAACTGTGCAT  
 ACAATGTGCAGCAGTCAACACCCAGCGTGGCGCTAGCACAGAGCCCCACACGTGTGCCGGAAGCCAGGGC  
 CACGCTGGCCTGCCACGGCCAGCGCCAGGAGCCACAGACTGCCCACTATCCGGGAAGCCAGGGGCCCT  
 CGCTCCACACTCAGAGCATCTGAGGGAACAACCTGACCAGAAGTGCAGTTGTTCTAACAGCCTCAGTAGT  
 CCTACTGGGCTGCCACGCCTCTCCAGGAAGCCTCCAGTCTAGGAGAGAGCTGAGCAAACTCCTGGGAAC  
 GTTGAGTTGATGTCACTGAGGTTTACTCCCTTGTGGTGTGAGTGTGAGATGCCCAAGGCTCCAGCAGAT  
 CTGCAGCCCCAGGGCGGGCTCCAGCCCTCATGGCAGACCAGGAGCCAGCGTGGCTGATCCCTCACTTGGC  
 TTCCAGCAAGAAGTCTTCGCTGTTTATTCTGAAAGATACTGTTTGGGAAGTGCAGGGAGCAGAGCTTCTC  
 AGCGCTGGCCTCTGAGCAGCTCAAAGTTATCTCTCATCTGCAAGGTCCCGAAATGGGCTGAGAGGCAGC  
 AGGACACAGATACAGCACTAGGAGCCATGAGCCAACACCTGCACCGGCCAGCAGCCCCAGGGCTCCAGCAC  
 TGCACAGCCACGTGCGATGGAACGCCAGCACACCGCCTGAGAAATGGGATGCTGCTGGTCTCCAGGCTCTGC  
 TCTGAAGATCCAGGTCCTGGGCCCTCTCTGCATAGTGGCCTCATAGGGGTGGTCACTCCAGCATCAG  
 GCTCATAGGGGTCACTGGCACTGTTGTAAAGCCTCAGGCTACTAGGCAATGTTCCGC

In a search of public sequence databases, the NOV10a nucleic acid sequence has 424  
 5 of 699 bases (60%) identical to a gb:GENBANK-ID:AF243500|acc:AF243500 transmembrane

protein X mRNA, complete cds, 1735 bp. mRNA from mouse/ *Mus musculus* ( $E = 4.9e^{-19}$ ).

Public nucleotide databases include all GenBank databases and the GeneSeq patent database.

The disclosed NOV10a polypeptide (SEQ ID NO:27) encoded by SEQ ID NO:25 has 472 amino acid residues and is presented in Table 10B using the one-letter amino acid code.

5 Signal P, Psort and/or Hydropathy results predict that NOV10a has a signal peptide and is likely to be localized in the plasma membrane with a certainty of 0.7900. In other embodiments, NOV10a may also be localized to the microbody (peroxisome) with a certainty of 0.5425, the Golgi body with a certainty of 0.3000, or in the endoplasmic reticulum (membrane) with a certainty of 0.2000. The most likely cleavage site for a NOV10a peptide is  
10 between amino acids 13 and 14, at: MEA-QY.

**Table 10B. Encoded NOV10a protein sequence (SEQ ID NO:27).**

MSLMLDDQPPMEAQYAEEGPGPGIFRAEPGDQQHPISQAVCWRSMRRCVAVLGALGLLAGAGVGSWLLVLYL CPAASQPISGTLQDEEITLSCSEASAEALLPALPKTVSFRINSEDFLEAQVRDQPRWLLVCHEGWSPALG LQICWSLGHRLRLTHHKGVNLTDIKLNSSQEFQALSPRLGGFLEEAWQPSRTTEAVRNNCTSGQVSLRCSEC GARPLASRIVGGQSVAPGRWPQASVALGFRHTCGGSVLAPRWVVTAAHCMHSFRLARLSSWRVHAGLVSHS AVRPHQGALVERIIPHPLYSAQNHDYDALLRLQTALNFSDTVGA VCLPAKEQHFPKGSRCWVSGWGHTHPS HTYSSDMLQD TVVPLLSTQLCNSSCVYSGALT PRMLCAGYLDGRADACQGD SGGPLVCPDGD TWRLVG VVSW GRGCAEPNHG VYAKVAEFLDWIHD TAQVSVGAGVGQGDF
---

A search of sequence databases reveals that the NOV10a amino acid sequence has 130 of 346 amino acid residues (37%) identical to, and 190 of 346 amino acid residues (54%) similar to, the 417 amino acid residue Serine Protease Hepsin (EC 3.4.21.-) (Transmembrane Protease, Serine 1) protein from *Homo sapiens* (ptnr:SWISSPROT-ACC:P05981 ( $E = 3.5e^{-56}$ )).  
15 Public amino acid databases include the GenBank databases, SwissProt, PDB and PIR.

NOV10a is expressed in at least the adrenal gland.

## 20 NOV10b

A disclosed NOV10b nucleic acid of 2148 nucleotides (also referred to as CG106783-02) encoding a novel Spinesin-like protein is shown in Table 10C. An open reading frame was identified beginning with an ATG initiation codon at nucleotides 156-158 and ending with a TGA codon at nucleotides 1410-1412. A putative untranslated region upstream from the  
25 initiation codon is underlined in Table 10C. The start and stop codons are in bold letters.

**Table 10C. NOV10b nucleotide sequence (SEQ ID NO:28).**

ACGCGGGATACAGGGAGGGGCCATGTGCGAACCAGGGAGACCTCATCTTCCAACCAAGCTTGCTGGGCTTGC  
 ATTTAATCAATGCATGGCCAGAGAACAGGAGCGGAACATTGCCTAGTAGACCCTGAGGCTTTACAACAGTGC  
 TACTGACCCCTATGAGCCTGATGCTGGATGACCAACCCCTATGGAGGCCAGTATGCAGAGGAGGGCCAG  
 GACCTGGGATCTTCAGAGCAGAGCCTGGAGACCAGCAGCATCCCATTCTCAGGCGGTGTGCTGGCGTTCCA  
 TGCGACGTGGCTGTGCAGTGTGGAGCCCTGGGGCTGCTGGCCGGTGCAGGTGTTGGCTCATGGCTCCTAG  
 TGCTGTATCTGTGCTGCTGCTCCTCAGCCCATTTCCGGGACCTTGCCAGGATGAGGAGATAACTTTGAGCT  
 GCTCAGAGGCCAGCGCTGAGGAAGCTGTGCTCCTGCACTCCCCAAAACAGTATCTTTCAGAATAAACAGCG  
 AAGACTTCTTGCTGGAAGCGCAAGTGAAGGATCAGCCACGCTGGCTCCTGGTCTGCCATGAGGGCTGGAGCC  
 CCGCCCTGGGGCTGCAGATCTGCTGGAGCCTTGGGCATCTCAGACTCACTCACCACAAGGGAGTAAACCTCA  
 CTGACATCAAACTCAACAGTTCCAGGAGTTTGGCTCAGCTCTCTCTAGACTGGGAGGCTTCTGGAGGAGG  
 CGTGGCAGCCAGGAACAACAGTCTGCTGCTCAAGTTGTTTCCCTCAGATGCTCTGAGTGTGGAGCGAGGC  
 CCCTGGCTTCCCGGATAGTTGGTGGGAGTCTGTGGCTCCTGGGCGCTGGCCGTGGCAGGCCAGCGTGGCCC  
 TGGGCTTCCGGCACACGTGTGGGGCTCTGTGCTAGCGCCACGCTGGGTGGTGAAGTCTGCACATTGTATGC  
 ACAGTGGCCAGAAATCATGACTACGAGCTCGCCCTCCTGAGGCTCCAGACCGCTCTCAACTTCTCAGACACTG  
 TGGGCGCTGTGTGCTGCGGCCAAGGAACAGCATTTTCCGAAGGGCTCGCGGTGCTGGGTGTGCGGCTGGT  
 GCCACACCCACCCTAGCCATACTTACAGCTCGGATATGCTCCAGGACACGGTGGTGGCCCTGCTCAGCACTC  
 AGCTCTGCAACAGCTCTTGCGTGTACAGCGGAGCCCTCACCCCCCGCATGCTTTGCGCTGGCTACCTGGACG  
 GAAGGGCTGATGCATGCCAGGGAGATAGCGGGGGCCCCCTAGTGTGCCAGATGGGGACACATGGCGCCTAG  
 TGGGGGTGGTCACTGCGGGGCGTGGCTGCGCAGAGCCCAATCACCAGGTGTCTACGCCAAGGTAGCTGAGT  
 TTCTGGACTGGATCCATGACACTGCTCAGGACTCCCTCCTCTGAGTCTCTGCTGTTTCTCCAGTCTCACTGC  
 ACACCACTGCCTCATGCTTCTGGGGCTCCAGCAGCTCCACTAATGGAGGAGAGGCAGTAGCCTCCGACAC  
 AGAACGATGGACCTCCTACTACTGTGTGTGAGGAACAGTCACTACCCACTGGCCAGCCACCCAGCCAACAG  
 GTCTCTCTCTTGGGCCCCGATTTCAGAGTCTCTTTCTCACTAGAGACTCAATGACAGAAGAGAGGCTGGG  
 ACTTGGTTGGGCATGCTGTGTTGCTGAGGATGAGGGGAGGAGAGAGGTAGGAGCTGGAGATGAAGAGAG  
 TGCTAGAAGCAGCAGGAAGCCTGCCCTTCTGCCCTCTCCCTCCCTGCCCTGTGTGAGTCTTTTAGGGAGG  
 GTGACTGGGAGGTGCCCCCGTCCCACCTTTTCTGTGCTCTAGGTGGGTAAGTGCCTCCCTAGAGGACT  
 CCATGGCTGAGAGGCTCCTGGGCAGATGGGGTCAAGGCTGGGCCAGTCCCAGATGAAGCCTATGGGAGTCAG  
 GACCTCTCCACTCTCCCTCTCCACTCCCTTCTGTTCTACCTGGCTGTGGCTGGCCCTGTGTGGGGTGG  
 GTACACTGGAACAAGAAGGTGGAGTTGGTCTAGGACATTGGTTTAAATGACAGTTCTGTGAACCTGGTC  
 CAAGGAGGTTCTGTTATTAAGTGATATATGGTCTTGAACAAAAAAAAAAAAAAAAAAAAA

In a search of public sequence databases, the NOV10b nucleic acid sequence, located on chromosome 7 has 1343 of 1446 bases (92%) identical to a gb:GENBANK-  
 ID:AB028140|acc:AB028140.1 mRNA from *Homo sapiens* (*Homo sapiens* mRNA for  
 5 spinesin, complete cds) ( $E = 3.2e^{-268}$ ). Public nucleotide databases include all GenBank  
 databases and the GeneSeq patent database.

The disclosed NOV10b polypeptide (SEQ ID NO:29) encoded by SEQ ID NO:28 has  
 418 amino acid residues and is presented in Table 10D using the one-letter amino acid code.  
 Signal P, Psort and/or Hydropathy results predict that NOV10a has a signal peptide and is  
 10 likely to be localized in the plasma membrane with a certainty of 0.7900. In other  
 embodiments, NOV10a may also be localized to the Golgi body with a certainty of 0.3000, the  
 microbody (peroxisome) with a certainty of 0.2036 or in the endoplasmic reticulum  
 (membrane) with a certainty of 0.2000. The most likely cleavage site for a NOV10a peptide is  
 between amino acids 13 and 14, at: MEA-QY.

**Table 10D. Encoded NOV10a protein sequence (SEQ ID NO:29).**

MSLMLDDQPMEAQYAE EGP GPGIFRAEPGDQQHPISQAVCWRS MRRC AVL GALGLLAGAGVGSWLLVLYL  
 CPAASQPISGTLQDEEITLSCSEASAE EALLPALPKTVSFRINSEDFLLEAQVRDQPRWLLVCHEGWSPALG  
 LQICWSLGLRLTHHKGVNLTDIKLNSSQEFALSPRLGGFLEAWQPRNNCTSGQVVSRLRCSECGARPLAS

RIVGGQSVAPGRWPWQASVALGFRHTCGGSLVAPRWVVTAAHCMHSAQNHDYDVALLRLQTALNFSDTVGA  
CLPAKEQHFPKGSRCWVSGWCHTHPSHTYSSDMLQDTVVPLLSTQLCNSSCVYSGALTFRMLCAGYLDGRAD  
ACQGDSSGGLVCPDGDTRWLVGVVSWGRGCAEPNHPGVYAKVAEFLDWIHDTAQDSL

A search of sequence databases reveals that the NOV10b amino acid sequence has 262 of 262 amino acid residues (100%) identical to, and 262 of 262 amino acid residues (100%) similar to, the 457 amino acid residue ptnr:SWISSNEW-ACC:Q9H3S3 protein from *Homo sapiens* (Human) (Transmembrane protease, serine 5 (EC 3.4.21.-) (Spinesin) ( $E = 8.7e^{-233}$ ). Public amino acid databases include the GenBank databases, SwissProt, PDB and PIR.

NOV10b is expressed in at least the following tissues: Colon, Brain, Placenta, Testis, Adrenal Gland/Suprarenal gland, Retina. Expression information was derived from the tissue sources of the sequences that were included in the derivation of the sequence of NOV10b. The sequence is predicted to be expressed in the following tissues because of the expression pattern of (GENBANK-ID: gb:GENBANK-ID:AB028140|acc:AB028140.1) a closely related *Homo sapiens* mRNA for spinesin, complete cds homolog in species *Homo sapiens* : brain.

TaqMan data for NOV10a can be found below in Example 2. The proteins encoded by the NOV10a and 10b nucleotides are very closely homologous as is shown in the alignment in Table 10E.

**Table 10E Alignment of NOV10a and 10b.**

		10	20	30	40	50
20	NOV10a	MSLMLDDQPPMEAQYAE	EGPGPGIFRAEPGDQ	QHPISQAVCWRSMR	RGCA	
	NOV10b	MSLMLDDQPPMEAQYAE	EGPGPGIFRAEPGDQ	QHPISQAVCWRSMR	RGCA	
		60	70	80	90	100
25	NOV10a	VLGALGLLAGAGVGS	WLLVLYLCPAASQPI	SGTLQDEEITLSC	SEASAE	
	NOV10b	VLGALGLLAGAGVGS	WLLVLYLCPAASQPI	SGTLQDEEITLSC	SEASAE	
		110	120	130	140	150
30	NOV10a	ALLPALPKTVSFRIN	SEDFLL	EAQVRDQPRWLL	VCHEGWSPALGL	QICWS
	NOV10b	ALLPALPKTVSFRIN	SEDFLL	EAQVRDQPRWLL	VCHEGWSPALGL	QICWS
		160	170	180	190	200
35	NOV10a	LGHLRLTHHKG'NL	TDIKLNSSQ	EFAQLSPRLGGF	LEEAWQH	SRTTEAVF
	NOV10b	LGHLRLTHHKG'NL	TDIKLNSSQ	EFAQLSPRLGGF	LEEAWQH	-----R
		210	220	230	240	250
40	NOV10a	NNCTSGQVVS	LRCECGARPLAS	RIVGGQSVAPGR	WPWQASVALG	FRHTC
	NOV10b	NNCTSGQVVS	LRCECGARPLAS	RIVGGQSVAPGR	WPWQASVALG	FRHTC
		260	270	280	290	300
45	NOV10a	GGSVLAPRWVVTAA	HCMHSFRLARLSS	WRVHAGLVSHSA	VRPHQGALVER	
	NOV10b	GGSVLAPRWVVTAA	HCMH	-----	-----	-----
		310	320	330	340	350
	NOV10a	IIPHPLY	SAQNHDYDVALL	RLQTALNFS	DTVGA	CLPAKEQHFPKGSRC

NOV10b -----SADNHDYDVALRLQTALNFSDTVGAVCLPAEQHFFFGSRCT

360 370 380 390 400

5 NOV10a VSGWGHTHPSHTYSSDMLQDTVVPLLSTQLCNSSCVYSGALTTPRMLCAGT  
 NOV10b VSGWGHTHPSHTYSSDMLQDTVVPLLSTQLCNSSCVYSGALTTPRMLCAGT

410 420 430 440 450

10 NOV10a LDGRADACQGDSSGGLVCPDGLTWRLVGVVSWGRGCAEPNHPG/YAKVAF  
 NOV10b LDGRADACQGDSSGGLVCPDGLTWRLVGVVSWGRGCAEPNHPG/YAKVAF

460 470

15 NOV10a FLDWIHDTACVSGAGVGQGDF (SEQ ID NO:27)  
 NOV10b FLDWIHDTACDGL----- (SEQ ID NO:29)

Homologies to any of the above NOV10 proteins will be shared by the other NOV10 protein insofar as they are homologous to each other as shown above. Any reference to NOV10 is assumed to refer to both of the NOV10 proteins in general, unless otherwise noted.

The disclosed NOV10a polypeptide has homology to the amino acid sequences shown in the BLASTP data listed in Table 10F.

Table 10F. BLAST results for NOV10a					
Gene Index/ Identifier	Protein/ Organism	Length (aa)	Identity (%)	Positives (%)	Expect
gi 13540535 ref NP_110397.1	transmembrane protease, serine 5; spinesin [Homo sapiens]	457	452/463 (97%)	453/463 (97%)	0.0
gi 13878822 sp Q9ER04 TMS5_MOUSE	TRANSMEMBRANE PROTEASE, SERINE 5 (SPINESIN)	455	360/461 (78%)	394/461 (85%)	0.0
gi 13507652 ref NP_109634.1	transmembrane protease, serine 5 (spinesin) [Mus musculus]	445	354/451 (78%)	388/451 (85%)	0.0
gi 14770563 ref XP_041427.1	transmembrane protease, serine 5 [Homo sapiens]	398	354/362 (97%)	355/362 (97%)	0.0
gi 12248777 dbj BAB20276.1  (AB016229)	type 2 spinesin [Mus musculus]	311	260/317 (82%)	281/317 (88%)	e-146

The homology between these and other sequences is shown graphically in the ClustalW analysis shown in Table 10G. In the ClustalW alignment of the NOV10 protein, as well as all other ClustalW analyses herein, the black outlined amino acid residues indicate regions of conserved sequence (*i.e.*, regions that may be required to preserve structural or functional properties), whereas non-highlighted amino acid residues are less conserved and can potentially be altered to a much broader extent without altering protein structure or function.

Table 10G. ClustalW Analysis of NOV10

- 1) Novel NOV10a (SEQ ID NO:27)
- 2) Novel NOV10b (SEQ ID NO:29)
- 5 4) gi|13540535|ref|NP\_110397.1| transmembrane protease, serine 5; spinesin [Homo sapiens] (SEQ ID NO:71)
- 5) gi|13878822|sp|Q9ER04|TMS5\_MOUSE TRANSMEMBRANE PROTEASE, SERINE 5 (SPINESIN) (SEQ ID NO:72)
- 10 6) gi|13507652|ref|NP\_109634.1| transmembrane protease, serine 5 (spinesin) [Mus musculus] (SEQ ID NO:73)
- 7) gi|14770563|ref|XP\_041427.1| transmembrane protease, serine 5 [Homo sapiens] (SEQ ID NO:74)
- 8) gi|12248777|dbj|BAB20276.1| (AB016229) type 2 spinesin [Mus musculus] (SEQ ID NO:75)

15

		10	20	30	40
NOV10a		MSLM	DDQP	PMEA	QYAE
NOV10b		MSLM	DDQP	PMEA	QYAE
gi 13540535		MSLM	DDQP	PMEA	QYAE
gi 13878822		MSPT	LDLQ	SPME	IRCT
gi 13507652		-----	MEIR	CTEE	CGPG
gi 14770563		MSLM	DDQP	PMEA	QYAE
gi 12248777		MSLM	DDQP	PMEA	QYAE

25

		50	60	70	80
NOV10a		CWRS	MRGC	AVLG	ALGL
NOV10b		CWRS	MRGC	AVLG	ALGL
gi 13540535		CWRS	MRGC	AVLG	ALGL
gi 13878822		RWCCL	RGCV	ILGL	VLLG
gi 13507652		RWCCL	RGCV	ILGL	VLLG
gi 14770563		CWRS	MRGC	AVLG	ALGL
gi 12248777		CWRS	MRGC	AVLG	ALGL

35

		90	100	110	120
NOV10a		SGTL	QEE	ETL	SCSE
NOV10b		SGTL	QEE	ETL	SCSE
gi 13540535		SGTL	QEE	ETL	SCSE
gi 13878822		SGTL	QEE	ETL	SCSE
gi 13507652		SGTL	QEE	ETL	SCSE
gi 14770563		SGTL	QEE	ETL	SCSE
gi 12248777		SGTL	QEE	ETL	SCSE

45

		130	140	150	160
NOV10a		LAQV	RD	CP	RWLL
NOV10b		LAQV	RD	CP	RWLL
gi 13540535		LAQV	RD	CP	RWLL
gi 13878822		LVQV	RA	PD	WLLV
gi 13507652		LVQV	RA	PD	WLLV
gi 14770563		LAQV	RD	CP	RWLL
gi 12248777		LAQV	RD	CP	RWLL

55

		170	180	190	200
NOV10a		GVLN	LDIK	LNSS	QEF
NOV10b		GVLN	LDIK	LNSS	QEF
gi 13540535		GVLN	LDIK	LNSS	QEF
gi 13878822		AVNL	SDIK	LNRS	QEF
gi 13507652		AVNL	SDIK	LNRS	QEF
gi 14770563		GVLN	LDIK	LNSS	QEF
gi 12248777		AVNL	SDIK	LNRS	QEF

65

210 220 230 240

NOV10a  
 NOV10b  
 gi | 13540535  
 gi | 13878822  
 gi | 13507652  
 gi | 14770563  
 gi | 12248777

250 260 270 280  
 NOV10a  
 NOV10b  
 gi | 13540535  
 gi | 13878822  
 gi | 13507652  
 gi | 14770563  
 gi | 12248777

290 300 310 320  
 NOV10a  
 NOV10b  
 gi | 13540535  
 gi | 13878822  
 gi | 13507652  
 gi | 14770563  
 gi | 12248777

330 340 350 360  
 NOV10a  
 NOV10b  
 gi | 13540535  
 gi | 13878822  
 gi | 13507652  
 gi | 14770563  
 gi | 12248777

370 380 390 400  
 NOV10a  
 NOV10b  
 gi | 13540535  
 gi | 13878822  
 gi | 13507652  
 gi | 14770563  
 gi | 12248777

410 420 430 440  
 NOV10a  
 NOV10b  
 gi | 13540535  
 gi | 13878822  
 gi | 13507652  
 gi | 14770563  
 gi | 12248777

450 460 470  
 NOV10a  
 NOV10b  
 gi | 13540535  
 gi | 13878822  
 gi | 13507652  
 gi | 14770563  
 gi | 12248777

Tables 1H-1I lists the domain description from DOMAIN analysis results against NOV10. This indicates that the NOV10 sequence has properties similar to those of other proteins known to contain this domain.

5

**Table 10H Domain Analysis of NOV10**

gnl|Smart|smart00020, Tryp\_SpC, Trypsin-like serine protease; Many of these are synthesised as inactive precursor zymogens that are cleaved during limited proteolysis to generate their active forms. A few, however, are active as single chain molecules, and others are inactive due to substitutions of the catalytic triad residues. (SEQ ID NO:96)  
CD-Length = 230 residues, 100.0% aligned  
Score = 266 bits (681), Expect = 2e-72

10 Query: 224 RIVGGQSVAPGRWPWQASVAL-GFRHTCGGSVLAPRWVVTAAHCMHSFRLARLSSWRVHA 282  
| | | | | + | | | + | | | | | + + + | | | + | | | | + + | | |  
Sbjct: 1 RIVGGSEANIGSFPPWQVSLQYRGGRHFCGGSLSIPRWVLTAAHCVYGSAP---SSIRVRL 57

15 Query: 283 GLVSHSAVRPHQGALVERIIPHPLYSAQNHDYDVALLRLQTALNFSDTVGA VCLPAKEQH 342  
| | + | | + + | | + + | + | | + | + | | + | | | + | | | +  
Sbjct: 58 GSHDLSSGEETQTVKVS KVI VHPNYPSTYDNDIAL LKLSEPVTLSDTVRPICLPSSGYN 117

20 Query: 343 FPKGSRWCWVGWGH THPSHTYSSDMLQDTVVP LLSTQLCNSSCVYSGALT PRLCAGYLD 402  
| | + | | | | | | | | | | + | | + | | + | | + | | | | +  
Sbjct: 118 VPAGTTCTVSGWGRTSESSGSLPDTLQEVNVP I VSNATCRRAYS GGPAITDNMLCAGGLE 177

Query: 403 GRADACQGD SGGPLVCPDGD TWR LVGVVSWGR -GCAEPNHPGVYAKVAEFLDWI 455  
| | | | | | | | | | + | | | + | | | | | | | | | | | + + + | | |  
Sbjct: 178 GGKDACQGD SGGPLVC -NDPRWVLVGI VSWGSYG CARPNKPGVYTRVSSYLDWI 230

**Table 10I. Domain Analysis of NOV10**

gnl|Pfam|pfam00089, trypsin, Trypsin. Proteins recognized include all proteins in families S1, S2A, S2B, S2C, and S5 in the classification of peptidases. Also included are proteins that are clearly members, but that lack peptidase activity, such as haptoglobin and protein Z (PRTZ\*). (SEQ ID NO:97)  
CD-Length = 217 residues, 100.0% aligned  
core = 211 bits (538), Expect = 6e-56

25 Query: 225 IVGGQSVAPGRWPWQASVALGFRHTCGGSVLAPRWVVTAAHCMHSFRLARLSSWRVHAGL 284  
| | | | + | + | | + + | | | | + + | | | | + | | | |  
Sbjct: 1 IVGGREAAQAGSFPPWQVSLQVSSGHFCGGSLSIENWVLTAAHCVSG-----ASSVRVVLGE 55

30 Query: 285 VSHSAVRPH-QGALVERIIPHPLYSAQNHDYDVALLRLQTALNFSDTVGA VCLPAKEQHF 343  
+ | | + + | | | + + | + | | + | + + + | | + | | +  
Sbjct: 56 HNLGTTEGTEQKFDVKKIIVHPNYPNPD - -NDIAL LKLKSPVTLGDTV RPICLP SASSDL 113

35 Query: 344 PKGSRWCWVGWGH THPSHTYSSDMLQDTVVP LLSTQLCNSSCVYSGALT PRLCAGYLDG 403  
| | + | | | | | + + | | + | | + + | + | + | | + | | + | |  
Sbjct: 114 PVGTTCSVSGWGR TK - -NLGTS DTLQEVVVP I VSR ETCRSA - -YGGTVTD TMICAGALGG 169

Query: 404 RADACQGD SGGPLVCPDGD TWR LVGVVSWGRGCAEPNHPGVYAKVAEFLDWI 455  
+ | | | | | | | | | | | | | | + | | | | | | + | | | + + + | | |  
Sbjct: 170 K-DACQGD SGGPLVCS DG - -ELVGIVSWGYGCAVGNYPGVYTRVSRYLDWI 217



Failure to thrive, nutritional edema, and hypoproteinemia with normal sweat electrolytes were features of 2 affected male infants reported by Townes (1965) and Townes et al. (1967). A protein hydrolysate diet was beneficial. A male sib of the first patient reported by Townes (1965) had died, apparently of the same condition. Morris and Fisher (1967) reported an affected female who also had imperforate anus. The clinical picture in enterokinase deficiency (226200) is closely similar; however, the defect is not in the synthesis of trypsinogen but in the synthesis of the enterokinase which activates proteolytic enzymes produced by the pancreas. Oral pancreatin represents a therapeutically successful form of enzyme replacement (Townes, 1972). Trypsin (EC 3.4.21.4), like elastase (130120), is a member of the pancreatic family of serine proteases. MacDonald et al. (1982) reported nucleotide sequences of cDNAs representing 2 pancreatic rat trypsinogens. Using a rat cDNA probe, Honey et al. (1984, 1984) found that a 3.8-kb DNA fragment containing human trypsin-1 gene sequences cosegregated with chromosome 7, and assigned the gene further to 7q22-7qter by study of hybrids with a deletion of this segment. The trypsin gene is on mouse chromosome 6 (Honey et al., 1984). Carboxypeptidase A (114850) and trypsin are a syntenic pair conserved in mouse and man. Emi et al. (1986) isolated cDNA clones for 2 major human trypsinogen isozymes from a pancreatic cDNA library. The deduced amino acid sequences had 89% homology and the same number of amino acids (247), including a 15-amino acid signal peptide and an 8-amino acid activation peptide. Southern blot analysis of human genomic DNA with the cloned cDNA as a probe showed that the human trypsinogen genes constitute a family of more than 10, some of which may be pseudogenes or may be expressed in other stages of development.

Rowen et al. (1996) found that there are 8 trypsinogen genes embedded in the beta T-cell receptor locus or cluster of genes (TCRB; 186930) mapping to 7q35. In the 685-kb DNA segment that they sequenced they found 5 tandemly arrayed 10-kb locus-specific repeats (homology units) at the 3-prime end of the locus. These repeats exhibited 90 to 91% overall nucleotide similarity, and embedded within each is a trypsinogen gene. Alignment of pancreatic trypsinogen cDNAs with the germline sequences showed that these trypsinogen genes contain 5 exons that span approximately 3.6 kb. Further analyses revealed 2 trypsinogen pseudogenes and 1 relic trypsinogen gene at the 5-prime end of the sequence, all in inverted transcriptional orientation. They denoted 8 trypsinogen genes T1 through T8 from 5-prime to 3-prime.

Rowen et al. (1996) found that only 2 of 3 pancreatically expressed trypsinogen cDNAs correspond to trypsinogen genes in the TCRB locus; T4 was denoted trypsinogen 1

and T8 was denoted trypsinogen 2 (601564). The third pancreatic cDNA, identified independently as trypsinogen 3 (Tani et al., 1990) and 4 (Wiegand et al., 1993), is distinct from the third apparently functional trypsinogen gene (T6) in the TCRB locus but related to the other pancreatic trypsinogens. Rowen et al. (1996) stated that the T6 gene is deleted in a common insertion-deletion polymorphism; if it is functional, its function is apparently not essential. Some of the trypsinogen genes are expressed in nonpancreatic tissues where their function is unknown. Rowen et al. (1996) noted that the intercalation of the trypsinogen genes in the TCRB locus is conserved in mouse and chicken, suggesting shared functional or regulatory constraints, as has been postulated for genes in the major histocompatibility complex (such as class I, II, and III genes) that share similar long-term organizational relationships.

Rowen et al. (1996) mapped the gene corresponding to the third pancreatic trypsinogen cDNA by fluorescence in situ hybridization. They used a cosmid clone containing 3 trypsinogen genes. Strong hybridization to chromosome 7 and weaker hybridization to chromosome 9 were observed. They isolated and partially sequenced 4 cosmid clones from the chromosome 9 region. They found that the region represents a duplication and translocation of a DNA segment from the 3-prime end of the TCRB locus that includes at least 7 V(beta) elements and a functional trypsinogen gene denoted T9. The assignment of the PRSS1 gene to 7q35 is established by the demonstration of its sequence within the sequence of the 'locus' for the T-cell receptor beta-chain (Rowen et al., 1996). Since hereditary pancreatitis (167800) has been mapped rather precisely to 7q35 and since a defect in the trypsinogen gene has been identified in hereditary pancreatitis, the assignment of the trypsinogen gene can be refined from 7q32-qter to 7q35.

Whitcomb et al. (1996) stated that the high degree of DNA sequence homology (more than 91%) present among this cluster of 5 trypsinogen genes identified by Rowen et al. (1996) demanded that highly specific sequence analysis strategies be developed for mutational screening in families with hereditary pancreatitis. This was necessary to ensure that each sequencing run contained only the 2 alleles corresponding to a single gene, thereby permitting detection of heterozygotes in this autosomal dominant disorder, and not a dozen or more alleles from multiple related trypsinogen-like genes, which would make detection of heterozygotes nearly impossible. In a family with hereditary pancreatitis, Whitcomb et al. (1996) found that affected individuals had a single G-to-A transition mutation in the third exon of cationic trypsinogen (276000.0001). This mutation was predicted to result in an arg105-to-his substitution in the trypsin gene (residue number 117 in the more common chymotrypsin

number system). Subsequently, the same mutation was found in a total of 5 different hereditary pancreatitis kindreds (4 from the U.S. and 1 from Italy) containing a total of 20 affected individuals and 6 obligate carriers. The mutation was found in none of the obligate unaffected members (individuals who married into the family). Subsequent haplotyping  
5 revealed that all 4 of the American families displayed the same high risk haplotype over a 4-cM region encompassing 7 STR markers, confirming the likelihood that these kindreds shared a common ancestor, although no link could be found through 8 generations. A fifth family from Italy displayed a unique haplotype indicating that the same mutation had occurred on at least 2 occasions. The G-to-A mutation at codon 117 created a novel enzyme recognition site  
10 for AflIII which provided a facile means to screen for the mutation. As with the obligate unaffected members of the pancreatitis kindreds, none of 140 controls possessed the G-to-A mutation as assayed by the lack of AflIII digestion of the amplified exonic DNA.

Ferec et al. (1999) studied 14 families with hereditary pancreatitis and found mutations in the PRSS1 gene in 8 families. In 4 of these families, the mutation (R117H; 276000.0001)  
15 had been described by Whitcomb et al. (1996). Three mutations were described in 4 other families (276000.0002, 276000.0003, 276000.0005).

Sahin-Toth et al. (1999) studied the roles of the 2 most frequent PRSS1 mutations in hereditary pancreatitis, R117H and N21I (276000.0002). They stated that the R117H mutation is believed to cause pancreatitis by eliminating an essential autolytic cleavage site in trypsin,  
20 thereby rendering the protease resistant to inactivation through autolysis. Sahin-Toth et al. (1999) demonstrated that the R117H mutation also significantly inhibited autocatalytic trypsinogen breakdown under Ca(2+)-free conditions and stabilized the zymogen form of rat trypsin. Taken together with findings demonstrating that the N21I mutation stabilized rat trypsinogen against

25 autoactivation and consequent autocatalytic degradation, the observations suggested a unifying molecular pathomechanism for hereditary pancreatitis in which zymogen stabilization plays a central role.

Sahin-Toth and Toth (2000) demonstrated that the R117H and N21I mutations significantly enhance autoactivation of human cationic trypsinogen in vitro, in a manner that  
30 correlates with the severity of clinical symptoms in hereditary pancreatitis. In addition, the R117H mutation inhibited autocatalytic inactivation of trypsin, while the N21I mutation had no such effect. Thus, increased trypsinogen activation in the pancreas is presumably the common initiating step in both forms of hereditary pancreatitis, whereas trypsin stabilization may also contribute to hereditary pancreatitis associated with the R117H mutation.

The disclosed NOV10 nucleic acid of the invention encoding a spinesin-like protein includes the nucleic acid whose sequence is provided in Table 10A and 10D or a fragment thereof. The invention also includes a mutant or variant nucleic acid any of whose bases may be changed from the corresponding base shown in Table 10A and 10D while still encoding a protein that maintains its spinesin-like activities and physiological functions, or a fragment of such a nucleic acid. The invention further includes nucleic acids whose sequences are complementary to those just described, including nucleic acid fragments that are complementary to any of the nucleic acids just described. The invention additionally includes nucleic acids or nucleic acid fragments, or complements thereto, whose structures include chemical modifications. Such modifications include, by way of nonlimiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in therapeutic applications in a subject. In the mutant or variant nucleic acids, and their complements, up to about 40% percent of the bases may be so changed.

The disclosed NOV10 protein of the invention includes the spinesin-like protein whose sequence is provided in Table 10B or 10E. The invention also includes a mutant or variant protein any of whose residues may be changed from the corresponding residue shown in Table 10B or 10E while still encoding a protein that maintains its spinesin-like activities and physiological functions, or a functional fragment thereof. In the mutant or variant protein, up to about 22% percent of the residues may be so changed.

The invention further encompasses antibodies and antibody fragments, such as  $F_{ab}$  or  $(F_{ab})_2$ , that bind immunospecifically to any of the proteins of the invention.

The above defined information for this invention suggests that this spinesin-like protein (NOV10) may function as a member of a "spinesin family". Therefore, the NOV10 nucleic acids and proteins identified here may be useful in potential therapeutic applications implicated in (but not limited to) various pathologies and disorders as indicated below. The potential therapeutic applications for this invention include, but are not limited to: protein therapeutic, small molecule drug target, antibody target (therapeutic, diagnostic, drug targeting/cytotoxic antibody), diagnostic and/or prognostic marker, gene therapy (gene delivery/gene ablation), research tools, tissue regeneration *in vivo* and *in vitro* of all tissues and cell types composing (but not limited to) those defined here.

The NOV10 nucleic acids and proteins of the invention are useful in potential therapeutic applications implicated in cancer including but not limited to various pathologies

and disorders as indicated below. For example, a cDNA encoding the spinesin-like protein (NOV10) may be useful in gene therapy, and the spinesin-like protein (NOV10) may be useful when administered to a subject in need thereof. By way of nonlimiting example, the compositions of the present invention will have efficacy for treatment of patients suffering

5 from cancer, trauma, tissue regeneration (in vitro and in vivo), viral/bacterial/parasitic infections, immunological disease, respiratory disease, gastro-intestinal diseases, reproductive health, neurological and neurodegenerative diseases, bone marrow transplantation, metabolic and endocrine diseases, allergy and inflammation, nephrological disorders, cardiovascular diseases, muscle, bone, joint and skeletal disorders, hematopoietic disorders, urinary system

10 disorders, Tissue and organ transplantation, Cardiomyopathy, Atherosclerosis, Hypertension, Congenital heart defects, Aortic stenosis, Atrial septal defect (ASD), Atrioventricular (A-V) canal defect, Ductus arteriosus, Pulmonary stenosis, Subaortic stenosis, Ventricular septal defect (VSD), valve diseases, Scleroderma, Obesity, Hypertension, Fibromuscular dysplasia, Stroke, Aneurysm, Myocardial infarction, Embolism, Bypass surgery, Anemia, Bleeding

15 disorders, Adrenoleukodystrophy, Congenital Adrenal Hyperplasia, Diabetes, Von Hippel-Lindau (VHL) syndrome, Pancreatitis, Hyperparathyroidism, Hypoparathyroidism, Hyperthyroidism and Hypothyroidism, AIDS, Endometriosis, infertility, Xerostomia, Hypercalcaemia, Ulcers, Cirrhosis, Inflammatory bowel disease, Diverticular disease, Hirschsprung's disease, Crohn's Disease, Appendicitis, Hemophilia, hypercoagulation,

20 autoimmune disease, allergies, immunodeficiencies, transplantation, Graft versus host disease (GVHD), Ataxia-telangiectasia, Autoimmune disease, Hemophilia, Hypercoagulation, Idiopathic thrombocytopenic purpura, Immunodeficiencies, Lymphedema, Allergies, Hemophilia, hypercoagulation, idiopathic thrombocytopenic purpura, Lymphoedema, Tonsillitis, Osteoporosis, Hypercalcaemia, Arthritis, Ankylosing spondylitis, Scoliosis,

25 Tendinitis, Muscular dystrophy, Lesch-Nyhan syndrome, Myasthenia gravis, Dental disease and infection, Alzheimer's disease, Stroke, Tuberous sclerosis, hypercalcaemia, Parkinson's disease, Huntington's disease, Cerebral palsy, Epilepsy, Lesch-Nyhan syndrome, Multiple sclerosis, Ataxia-telangiectasia, Behavioral disorders, Addiction, Anxiety, Pain, Neuroprotection, Endocrine dysfunctions, Growth and reproductive disorders, Myasthenia

30 gravis, Leukodystrophies, Pain, Neuroprotection, Systemic lupus erythematosus, Autoimmune disease, Emphysema, Scleroderma, ARDS, Pharyngitis, Laryngitis, Asthma, Hearing loss, Tinnitus, Psoriasis, Actinic keratosis, Tuberous sclerosis, Acne, Hair growth, alopecia, pigmentation disorders, endocrine disorders, cystitis, incontinence, Autoimmune disease, Renal artery stenosis, Interstitial nephritis, Glomerulonephritis, Polycystic kidney disease,

Systemic lupus erythematosus, Renal tubular acidosis, IgA nephropathy, Hypercalcaemia, Lesch-Nyhan syndrome, Vesicoureteral reflux, and other pathologies and conditions.. The NOV10 nucleic acid encoding the spinesin-like protein of the invention, or fragments thereof, may further be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed.

NOV10 nucleic acids and polypeptides are further useful in the generation of antibodies that bind immuno-specifically to the novel NOV10 substances for use in therapeutic or diagnostic methods. These antibodies may be generated according to methods known in the art, using prediction from hydrophobicity charts, as described in the "Anti-NOVX Antibodies" section below. The disclosed NOV10 protein has multiple hydrophilic regions, each of which can be used as an immunogen. In one embodiment, a contemplated NOV10 epitope is from about amino acids 5 to 50. In another embodiment, a NOV10 epitope is from about amino acids 60 to 70. In additional embodiments, NOV10 epitope is from about amino acids 100 to 130, from about amino acids 140 to 210, from about amino acid 270 to 320, from about amino acid 340 to 360, from about amino acid 390 to 410, and from about amino acids 430 to 450. These novel proteins can be used in assay systems for functional analysis of various human disorders, which will help in understanding of pathology of the disease and development of new drug targets for various disorders.

## **NOVX Nucleic Acids and Polypeptides**

One aspect of the invention pertains to isolated nucleic acid molecules that encode NOVX polypeptides or biologically active portions thereof. Also included in the invention are nucleic acid fragments sufficient for use as hybridization probes to identify NOVX-encoding nucleic acids (*e.g.*, NOVX mRNAs) and fragments for use as PCR primers for the amplification and/or mutation of NOVX nucleic acid molecules. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA), RNA molecules (*e.g.*, mRNA), analogs of the DNA or RNA generated using nucleotide analogs, and derivatives, fragments and homologs thereof. The nucleic acid molecule may be single-stranded or double-stranded, but preferably is comprised double-stranded DNA.

An NOVX nucleic acid can encode a mature NOVX polypeptide. As used herein, a "mature" form of a polypeptide or protein disclosed in the present invention is the product of a naturally occurring polypeptide or precursor form or proprotein. The naturally occurring

polypeptide, precursor or proprotein includes, by way of nonlimiting example, the full-length gene product, encoded by the corresponding gene. Alternatively, it may be defined as the polypeptide, precursor or proprotein encoded by an ORF described herein. The product "mature" form arises, again by way of nonlimiting example, as a result of one or more naturally occurring processing steps as they may take place within the cell, or host cell, in which the gene product arises. Examples of such processing steps leading to a "mature" form of a polypeptide or protein include the cleavage of the N-terminal methionine residue encoded by the initiation codon of an ORF, or the proteolytic cleavage of a signal peptide or leader sequence. Thus a mature form arising from a precursor polypeptide or protein that has residues 1 to N, where residue 1 is the N-terminal methionine, would have residues 2 through N remaining after removal of the N-terminal methionine. Alternatively, a mature form arising from a precursor polypeptide or protein having residues 1 to N, in which an N-terminal signal sequence from residue 1 to residue M is cleaved, would have the residues from residue M+1 to residue N remaining. Further as used herein, a "mature" form of a polypeptide or protein may arise from a step of post-translational modification other than a proteolytic cleavage event. Such additional processes include, by way of non-limiting example, glycosylation, myristoylation or phosphorylation. In general, a mature polypeptide or protein may result from the operation of only one of these processes, or a combination of any of them.

The term "probes", as utilized herein, refers to nucleic acid sequences of variable length, preferably between at least about 10 nucleotides (nt), 100 nt, or as many as approximately, *e.g.*, 6,000 nt, depending upon the specific use. Probes are used in the detection of identical, similar, or complementary nucleic acid sequences. Longer length probes are generally obtained from a natural or recombinant source, are highly specific, and much slower to hybridize than shorter-length oligomer probes. Probes may be single- or double-stranded and designed to have specificity in PCR, membrane-based hybridization technologies, or ELISA-like technologies.

The term "isolated" nucleic acid molecule, as utilized herein, is one, which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*, sequences located at the 5'- and 3'-termini of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated NOVX nucleic acid molecules can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell/tissue from which the nucleic acid is derived (*e.g.*, brain, heart, liver,

spleen, etc.). Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material or culture medium when produced by recombinant techniques, or of chemical precursors or other chemicals when chemically synthesized.

5           A nucleic acid molecule of the invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, or a complement of this aforementioned nucleotide sequence, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or a portion of the nucleic acid sequence of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23,  
10       25, and 28 as a hybridization probe, NOVX molecules can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook, *et al.*, (eds.), MOLECULAR CLONING: A LABORATORY MANUAL 2<sup>nd</sup> Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989; and Ausubel, *et al.*, (eds.), CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993.)

15           A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to NOVX nucleotide sequences can be prepared by standard  
20       synthetic techniques, *e.g.*, using an automated DNA synthesizer.

          As used herein, the term "oligonucleotide" refers to a series of linked nucleotide residues, which oligonucleotide has a sufficient number of nucleotide bases to be used in a PCR reaction. A short oligonucleotide sequence may be based on, or designed from, a genomic or cDNA sequence and is used to amplify, confirm, or reveal the presence of an  
25       identical, similar or complementary DNA or RNA in a particular cell or tissue. Oligonucleotides comprise portions of a nucleic acid sequence having about 10 nt, 50 nt, or 100 nt in length, preferably about 15 nt to 30 nt in length. In one embodiment of the invention, an oligonucleotide comprising a nucleic acid molecule less than 100 nt in length would further comprise at least 6 contiguous nucleotides SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15,  
30       17, 19, 21, 23, 25, and 28, or a complement thereof. Oligonucleotides may be chemically synthesized and may also be used as probes.

          In another embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule that is a complement of the nucleotide sequence shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, or a portion of this nucleotide



sequence (*e.g.*, a fragment that can be used as a probe or primer or a fragment encoding a biologically-active portion of an NOVX polypeptide). A nucleic acid molecule that is complementary to the nucleotide sequence shown SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, or 28 is one that is sufficiently complementary to the nucleotide sequence shown SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, or 28 that it can hydrogen bond with little or no mismatches to the nucleotide sequence shown SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, thereby forming a stable duplex.

As used herein, the term "complementary" refers to Watson-Crick or Hoogsteen base pairing between nucleotides units of a nucleic acid molecule, and the term "binding" means the physical or chemical interaction between two polypeptides or compounds or associated polypeptides or compounds or combinations thereof. Binding includes ionic, non-ionic, van der Waals, hydrophobic interactions, and the like. A physical interaction can be either direct or indirect. Indirect interactions may be through or due to the effects of another polypeptide or compound. Direct binding refers to interactions that do not take place through, or due to, the effect of another polypeptide or compound, but instead are without other substantial chemical intermediates.

Fragments provided herein are defined as sequences of at least 6 (contiguous) nucleic acids or at least 4 (contiguous) amino acids, a length sufficient to allow for specific hybridization in the case of nucleic acids or for specific recognition of an epitope in the case of amino acids, respectively, and are at most some portion less than a full length sequence. Fragments may be derived from any contiguous portion of a nucleic acid or amino acid sequence of choice. Derivatives are nucleic acid sequences or amino acid sequences formed from the native compounds either directly or by modification or partial substitution. Analogs are nucleic acid sequences or amino acid sequences that have a structure similar to, but not identical to, the native compound but differs from it in respect to certain components or side chains. Analogs may be synthetic or from a different evolutionary origin and may have a similar or opposite metabolic activity compared to wild type. Homologs are nucleic acid sequences or amino acid sequences of a particular gene that are derived from different species.

Derivatives and analogs may be full length or other than full length, if the derivative or analog contains a modified nucleic acid or amino acid, as described below. Derivatives or analogs of the nucleic acids or proteins of the invention include, but are not limited to, molecules comprising regions that are substantially homologous to the nucleic acids or proteins of the invention, in various embodiments, by at least about 70%, 80%, or 95% identity (with a preferred identity of 80-95%) over a nucleic acid or amino acid sequence of

identical size or when compared to an aligned sequence in which the alignment is done by a computer homology program known in the art, or whose encoding nucleic acid is capable of hybridizing to the complement of a sequence encoding the aforementioned proteins under stringent, moderately stringent, or low stringent conditions. See e.g. Ausubel, *et al.*, CURRENT  
5 PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, NY, 1993, and below.

A "homologous nucleic acid sequence" or "homologous amino acid sequence," or variations thereof, refer to sequences characterized by a homology at the nucleotide level or amino acid level as discussed above. Homologous nucleotide sequences encode those sequences coding for isoforms of NOVX polypeptides. Isoforms can be expressed in different  
10 tissues of the same organism as a result of, for example, alternative splicing of RNA. Alternatively, isoforms can be encoded by different genes. In the invention, homologous nucleotide sequences include nucleotide sequences encoding for an NOVX polypeptide of species other than humans, including, but not limited to: vertebrates, and thus can include, e.g., frog, mouse, rat, rabbit, dog, cat, cow, horse, and other organisms. Homologous nucleotide  
15 sequences also include, but are not limited to, naturally occurring allelic variations and mutations of the nucleotide sequences set forth herein. A homologous nucleotide sequence does not, however, include the exact nucleotide sequence encoding human NOVX protein. Homologous nucleic acid sequences include those nucleic acid sequences that encode conservative amino acid substitutions (see below) in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17,  
20 19, 21, 23, 25, and 28, as well as a polypeptide possessing NOVX biological activity. Various biological activities of the NOVX proteins are described below.

An NOVX polypeptide is encoded by the open reading frame ("ORF") of an NOVX nucleic acid. An ORF corresponds to a nucleotide sequence that could potentially be translated into a polypeptide. A stretch of nucleic acids comprising an ORF is uninterrupted by a stop  
25 codon. An ORF that represents the coding sequence for a full protein begins with an ATG "start" codon and terminates with one of the three "stop" codons, namely, TAA, TAG, or TGA. For the purposes of this invention, an ORF may be any part of a coding sequence, with or without a start codon, a stop codon, or both. For an ORF to be considered as a good candidate for coding for a *bona fide* cellular protein, a minimum size requirement is often set,  
30 e.g., a stretch of DNA that would encode a protein of 50 amino acids or more.

The nucleotide sequences determined from the cloning of the human NOVX genes allows for the generation of probes and primers designed for use in identifying and/or cloning NOVX homologues in other cell types, e.g. from other tissues, as well as NOVX homologues from other vertebrates. The probe/primer typically comprises substantially purified

oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 25, 50, 100, 150, 200, 250, 300, 350 or 400 consecutive sense strand nucleotide sequence SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, or 28; or an anti-sense strand nucleotide sequence of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, or 28; or of a naturally occurring mutant of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28.

Probes based on the human NOVX nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In various embodiments, the probe further comprises a label group attached thereto, *e.g.* the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissues which mis-express an NOVX protein, such as by measuring a level of an NOVX-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting NOVX mRNA levels or determining whether a genomic NOVX gene has been mutated or deleted.

"A polypeptide having a biologically-active portion of an NOVX polypeptide" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. A nucleic acid fragment encoding a "biologically-active portion of NOVX" can be prepared by isolating a portion SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, or 28, that encodes a polypeptide having an NOVX biological activity (the biological activities of the NOVX proteins are described below), expressing the encoded portion of NOVX protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of NOVX.

#### **NOVX Nucleic Acid and Polypeptide Variants**

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequences shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28 due to degeneracy of the genetic code and thus encode the same NOVX proteins as that encoded by the nucleotide sequences shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29.

In addition to the human NOVX nucleotide sequences shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, it will be appreciated by those skilled in the art that

DNA sequence polymorphisms that lead to changes in the amino acid sequences of the NOVX polypeptides may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the NOVX genes may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame (ORF) encoding an NOVX protein, preferably a vertebrate NOVX protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of the NOVX genes. Any and all such nucleotide variations and resulting amino acid polymorphisms in the NOVX polypeptides, which are the result of natural allelic variation and that do not alter the functional activity of the NOVX polypeptides, are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding NOVX proteins from other species, and thus that have a nucleotide sequence that differs from the human SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28 are intended to be within the scope of the invention. Nucleic acid molecules corresponding to natural allelic variants and homologues of the NOVX cDNAs of the invention can be isolated based on their homology to the human NOVX nucleic acids disclosed herein using the human cDNAs, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 6 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28. In another embodiment, the nucleic acid is at least 10, 25, 50, 100, 250, 500, 750, 1000, 1500, or 2000 or more nucleotides in length. In yet another embodiment, an isolated nucleic acid molecule of the invention hybridizes to the coding region. As used herein, the term "hybridizes under stringent conditions" is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each other.

Homologs (*i.e.*, nucleic acids encoding NOVX proteins derived from species other than human) or other related sequences (*e.g.*, paralogs) can be obtained by low, moderate or high stringency hybridization with all or a portion of the particular human sequence as a probe using methods well known in the art for nucleic acid hybridization and cloning.

As used herein, the phrase "stringent hybridization conditions" refers to conditions under which a probe, primer or oligonucleotide will hybridize to its target sequence, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures than

shorter sequences. Generally, stringent conditions are selected to be about 5 °C lower than the thermal melting point (T<sub>m</sub>) for the specific sequence at a defined ionic strength and pH. The T<sub>m</sub> is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium. Since the target sequences are generally present at excess, at T<sub>m</sub>, 50% of the probes are occupied at equilibrium. Typically, stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes, primers or oligonucleotides (*e.g.*, 10 nt to 50 nt) and at least about 60°C for longer probes, primers and oligonucleotides. Stringent conditions may also be achieved with the addition of destabilizing agents, such as formamide.

Stringent conditions are known to those skilled in the art and can be found in Ausubel, *et al.*, (eds.), CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. Preferably, the conditions are such that sequences at least about 65%, 70%, 75%, 85%, 90%, 95%, 98%, or 99% homologous to each other typically remain hybridized to each other. A non-limiting example of stringent hybridization conditions are hybridization in a high salt buffer comprising 6X SSC, 50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.02% BSA, and 500 mg/ml denatured salmon sperm DNA at 65°C, followed by one or more washes in 0.2X SSC, 0.01% BSA at 50°C. An isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequences SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In a second embodiment, a nucleic acid sequence that is hybridizable to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, or fragments, analogs or derivatives thereof, under conditions of moderate stringency is provided. A non-limiting example of moderate stringency hybridization conditions are hybridization in 6X SSC, 5X Denhardt's solution, 0.5% SDS and 100 mg/ml denatured salmon sperm DNA at 55°C, followed by one or more washes in 1X SSC, 0.1% SDS at 37°C. Other conditions of moderate stringency that may be used are well-known within the art. *See, e.g.*, Ausubel, *et al.* (eds.), 1993, CURRENT PROTOCOLS IN

MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990; GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY.

In a third embodiment, a nucleic acid that is hybridizable to the nucleic acid molecule comprising the nucleotide sequences SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, or fragments, analogs or derivatives thereof, under conditions of low stringency, is provided. A non-limiting example of low stringency hybridization conditions are hybridization in 35% formamide, 5X SSC, 50 mM Tris-HCl (pH 7.5), 5 mM EDTA, 0.02% PVP, 0.02% Ficoll, 0.2% BSA, 100 mg/ml denatured salmon sperm DNA, 10% (wt/vol) dextran sulfate at 40°C, followed by one or more washes in 2X SSC, 25 mM Tris-HCl (pH 7.4), 5 mM EDTA, and 0.1% SDS at 50°C. Other conditions of low stringency that may be used are well known in the art (*e.g.*, as employed for cross-species hybridizations). *See, e.g.*, Ausubel, *et al.* (eds.), 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, NY, and Kriegler, 1990, GENE TRANSFER AND EXPRESSION, A LABORATORY MANUAL, Stockton Press, NY; Shilo and Weinberg, 1981. *Proc Natl Acad Sci USA* 78: 6789-6792.

### Conservative Mutations

In addition to naturally-occurring allelic variants of NOVX sequences that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequences SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, thereby leading to changes in the amino acid sequences of the encoded NOVX proteins, without altering the functional ability of said NOVX proteins. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequences of the NOVX proteins without altering their biological activity, whereas an "essential" amino acid residue is required for such biological activity. For example, amino acid residues that are conserved among the NOVX proteins of the invention are predicted to be particularly non-amenable to alteration. Amino acids for which conservative substitutions can be made are well-known within the art.

Another aspect of the invention pertains to nucleic acid molecules encoding NOVX proteins that contain changes in amino acid residues that are not essential for activity. Such NOVX proteins differ in amino acid sequence from SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28 yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein

comprises an amino acid sequence at least about 45% homologous to the amino acid sequences SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, and 29. Preferably, the protein encoded by the nucleic acid molecule is at least about 60% homologous to SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, and 29; more preferably at least about 70% homologous SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29; still more preferably at least about 80% homologous to SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29; even more preferably at least about 90% homologous to SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29; and most preferably at least about 95% homologous to SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29.

An isolated nucleic acid molecule encoding an NOVX protein homologous to the protein of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein.

Mutations can be introduced into SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28 by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted, non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined within the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted non-essential amino acid residue in the NOVX protein is replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of an NOVX coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for NOVX biological activity to identify mutants that retain activity. Following mutagenesis SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, the encoded protein can be expressed by any recombinant technology known in the art and the activity of the protein can be determined.

The relatedness of amino acid families may also be determined based on side chain interactions. Substituted amino acids may be fully conserved "strong" residues or fully conserved "weak" residues. The "strong" group of conserved amino acid residues may be any one of the following groups: STA, NEQK, NHQK, NDEQ, QHRK, MILV, MILF, HY, FYW, wherein the single letter amino acid codes are grouped by those amino acids that may be substituted for each other. Likewise, the "weak" group of conserved residues may be any one of the following: CSA, ATV, SAG, STNK, STPA, SGND, SNDEQK, NDEQHK, NEQHRK, VLIM, HFY, wherein the letters within each group represent the single letter amino acid code.

In one embodiment, a mutant NOVX protein can be assayed for (i) the ability to form protein:protein interactions with other NOVX proteins, other cell-surface proteins, or biologically-active portions thereof, (ii) complex formation between a mutant NOVX protein and an NOVX ligand; or (iii) the ability of a mutant NOVX protein to bind to an intracellular target protein or biologically-active portion thereof; (e.g. avidin proteins).

In yet another embodiment, a mutant NOVX protein can be assayed for the ability to regulate a specific biological function (e.g., regulation of insulin release).

#### **Antisense Nucleic Acids**

Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein (e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence). In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire NOVX coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of an NOVX protein of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29, or antisense nucleic acids complementary to an NOVX nucleic acid sequence of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, are additionally provided.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding an NOVX protein. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues. In another embodiment, the antisense nucleic acid



molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding the NOVX protein. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

5           Given the coding strand sequences encoding the NOVX protein disclosed herein, antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of NOVX mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of NOVX mRNA. For example, 10 the antisense oligonucleotide can be complementary to the region surrounding the translation start site of NOVX mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) 15 can be chemically synthesized using naturally-occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids (*e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used).

Examples of modified nucleotides that can be used to generate the antisense nucleic 20 acid include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 25 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 30 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding an NOVX protein to thereby inhibit expression of the protein (e.g., by inhibiting transcription and/or translation). The hybridization can be by conventional  
5 nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered  
10 systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface (e.g., by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens). The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient nucleic acid molecules, vector  
15 constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an  $\alpha$ -anomeric nucleic acid molecule. An  $\alpha$ -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual  $\beta$ -units, the  
20 strands run parallel to each other. See, e.g., Gaultier, *et al.*, 1987. *Nucl. Acids Res.* **15**: 6625-6641. The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (See, e.g., Inoue, *et al.* 1987. *Nucl. Acids Res.* **15**: 6131-6148) or a chimeric RNA-DNA analogue (See, e.g., Inoue, *et al.*, 1987. *FEBS Lett.* **215**: 327-330.

## 25 **Ribozymes and PNA Moieties**

Nucleic acid modifications include, by way of non-limiting example, modified bases, and nucleic acids whose sugar phosphate backbones are modified or derivatized. These modifications are carried out at least in part to enhance the chemical stability of the modified nucleic acid, such that they may be used, for example, as antisense binding nucleic acids in  
30 therapeutic applications in a subject.

In one embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes as described in

Haselhoff and Gerlach 1988. *Nature* 334: 585-591) can be used to catalytically cleave NOVX mRNA transcripts to thereby inhibit translation of NOVX mRNA. A ribozyme having specificity for an NOVX-encoding nucleic acid can be designed based upon the nucleotide sequence of an NOVX cDNA disclosed herein (*i.e.*, SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28). For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in an NOVX-encoding mRNA. *See, e.g.*, U.S. Patent 4,987,071 to Cech, *et al.* and U.S. Patent 5,116,742 to Cech, *et al.* NOVX mRNA can also be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. *See, e.g.*, Bartel *et al.*, (1993) *Science* 261:1411-1418.

Alternatively, NOVX gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the NOVX nucleic acid (*e.g.*, the NOVX promoter and/or enhancers) to form triple helical structures that prevent transcription of the NOVX gene in target cells. *See, e.g.*, Helene, 1991. *Anticancer Drug Des.* 6: 569-84; Helene, *et al.* 1992. *Ann. N.Y. Acad. Sci.* 660: 27-36; Maher, 1992. *Bioassays* 14: 807-15.

In various embodiments, the NOVX nucleic acids can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids. *See, e.g.*, Hyrup, *et al.*, 1996. *Bioorg Med Chem* 4: 5-23. As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics (*e.g.*, DNA mimics) in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup, *et al.*, 1996. *supra*; Perry-O'Keefe, *et al.*, 1996. *Proc. Natl. Acad. Sci. USA* 93: 14670-14675.

PNAs of NOVX can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of NOVX can also be used, for example, in the analysis of single base pair mutations in a gene (*e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S<sub>1</sub> nucleases (*See*, Hyrup, *et al.*, 1996. *supra*); or as probes or primers for DNA sequence and hybridization (*See*, Hyrup, *et al.*, 1996, *supra*; Perry-O'Keefe, *et al.*, 1996. *supra*).

In another embodiment, PNAs of NOVX can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras of NOVX can be generated that may combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes (*e.g.*, RNase H and DNA polymerases) to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (*see*, Hyrup, *et al.*, 1996. *supra*). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup, *et al.*, 1996. *supra* and Finn, *et al.*, 1996. *Nucl Acids Res* 24: 3357-3363. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry, and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA. *See, e.g.*, Mag, *et al.*, 1989. *Nucl Acid Res* 17: 5973-5988. PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment. *See, e.g.*, Finn, *et al.*, 1996. *supra*. Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment. *See, e.g.*, Petersen, *et al.*, 1975. *Bioorg. Med. Chem. Lett.* 5: 1119-11124.

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (*see, e.g.*, Letsinger, *et al.*, 1989. *Proc. Natl. Acad. Sci. U.S.A.* 86: 6553-6556; Lemaitre, *et al.*, 1987. *Proc. Natl. Acad. Sci.* 84: 648-652; PCT Publication No. WO88/09810) or the blood-brain barrier (*see, e.g.*, PCT Publication No. WO 89/10134). In addition, oligonucleotides can be modified with hybridization triggered cleavage agents (*see, e.g.*, Krol, *et al.*, 1988. *BioTechniques* 6:958-976) or intercalating agents (*see, e.g.*, Zon, 1988. *Pharm. Res.* 5: 539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, and the like.

### **NOVX Polypeptides**

A polypeptide according to the invention includes a polypeptide including the amino acid sequence of NOVX polypeptides whose sequences are provided in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29. The invention also includes a mutant or variant

protein any of whose residues may be changed from the corresponding residues shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29 while still encoding a protein that maintains its NOVX activities and physiological functions, or a functional fragment thereof.

5 In general, an NOVX variant that preserves NOVX-like function includes any variant in which residues at a particular position in the sequence have been substituted by other amino acids, and further include the possibility of inserting an additional residue or residues between two residues of the parent protein as well as the possibility of deleting one or more residues from the parent sequence. Any amino acid substitution, insertion, or deletion is encompassed by the invention. In favorable circumstances, the substitution is a conservative substitution as  
10 defined above.

One aspect of the invention pertains to isolated NOVX proteins, and biologically-active portions thereof, or derivatives, fragments, analogs or homologs thereof. Also provided are polypeptide fragments suitable for use as immunogens to raise anti-NOVX antibodies. In one embodiment, native NOVX proteins can be isolated from cells or tissue sources by an  
15 appropriate purification scheme using standard protein purification techniques. In another embodiment, NOVX proteins are produced by recombinant DNA techniques. Alternative to recombinant expression, an NOVX protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" polypeptide or protein or biologically-active portion thereof  
20 is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the NOVX protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of NOVX proteins in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly-produced. In  
25 one embodiment, the language "substantially free of cellular material" includes preparations of NOVX proteins having less than about 30% (by dry weight) of non-NOVX proteins (also referred to herein as a "contaminating protein"), more preferably less than about 20% of non-NOVX proteins, still more preferably less than about 10% of non-NOVX proteins, and most preferably less than about 5% of non-NOVX proteins. When the NOVX protein or  
30 biologically-active portion thereof is recombinantly-produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the NOVX protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of NOVX proteins in which the protein is separated from chemical precursors or other chemicals that are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of NOVX proteins having less than about 30% (by dry weight) of chemical precursors or non-NOVX chemicals, more preferably less than about 20% chemical precursors or non-NOVX chemicals, still more preferably less than about 10% chemical precursors or non-NOVX chemicals, and most preferably less than about 5% chemical precursors or non-NOVX chemicals.

Biologically-active portions of NOVX proteins include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequences of the NOVX proteins (*e.g.*, the amino acid sequence shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29) that include fewer amino acids than the full-length NOVX proteins, and exhibit at least one activity of an NOVX protein. Typically, biologically-active portions comprise a domain or motif with at least one activity of the NOVX protein. A biologically-active portion of an NOVX protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acid residues in length.

Moreover, other biologically-active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one or more of the functional activities of a native NOVX protein.

In an embodiment, the NOVX protein has an amino acid sequence shown SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29. In other embodiments, the NOVX protein is substantially homologous to SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29, and retains the functional activity of the protein of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail, below. Accordingly, in another embodiment, the NOVX protein is a protein that comprises an amino acid sequence at least about 45% homologous to the amino acid sequence SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29, and retains the functional activity of the NOVX proteins of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29.

#### **Determining Homology Between Two or More Sequences**

To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced

in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity").

The nucleic acid sequence homology may be determined as the degree of identity between two sequences. The homology may be determined using computer programs known in the art, such as GAP software provided in the GCG program package. *See*, Needleman and Wunsch, 1970. *J Mol Biol* 48: 443-453. Using GCG GAP software with the following settings for nucleic acid sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3, the coding region of the analogous nucleic acid sequences referred to above exhibits a degree of identity preferably of at least 70%, 75%, 80%, 85%, 90%, 95%, 98%, or 99%, with the CDS (encoding) part of the DNA sequence shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28.

The term "sequence identity" refers to the degree to which two polynucleotide or polypeptide sequences are identical on a residue-by-residue basis over a particular region of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over that region of comparison, determining the number of positions at which the identical nucleic acid base (*e.g.*, A, T, C, G, U, or I, in the case of nucleic acids) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the region of comparison (*i.e.*, the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The term "substantial identity" as used herein denotes a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that has at least 80 percent sequence identity, preferably at least 85 percent identity and often 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison region.

### **Chimeric and Fusion Proteins**

The invention also provides NOVX chimeric or fusion proteins. As used herein, an NOVX "chimeric protein" or "fusion protein" comprises an NOVX polypeptide operatively-linked to a non-NOVX polypeptide. An "NOVX polypeptide" refers to a polypeptide having

an amino acid sequence corresponding to an NOVX protein SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 27, or 29, whereas a "non-NOVX polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein that is not substantially homologous to the NOVX protein, *e.g.*, a protein that is different from the NOVX protein and that is  
5 derived from the same or a different organism. Within an NOVX fusion protein the NOVX polypeptide can correspond to all or a portion of an NOVX protein. In one embodiment, an NOVX fusion protein comprises at least one biologically-active portion of an NOVX protein. In another embodiment, an NOVX fusion protein comprises at least two biologically-active portions of an NOVX protein. In yet another embodiment, an NOVX fusion protein  
10 comprises at least three biologically-active portions of an NOVX protein. Within the fusion protein, the term "operatively-linked" is intended to indicate that the NOVX polypeptide and the non-NOVX polypeptide are fused in-frame with one another. The non-NOVX polypeptide can be fused to the N-terminus or C-terminus of the NOVX polypeptide.

In one embodiment, the fusion protein is a GST-NOVX fusion protein in which the  
15 NOVX sequences are fused to the C-terminus of the GST (glutathione S-transferase) sequences. Such fusion proteins can facilitate the purification of recombinant NOVX polypeptides.

In another embodiment, the fusion protein is an NOVX protein containing a heterologous signal sequence at its N-terminus. In certain host cells (*e.g.*, mammalian host  
20 cells), expression and/or secretion of NOVX can be increased through use of a heterologous signal sequence.

In yet another embodiment, the fusion protein is an NOVX-immunoglobulin fusion protein in which the NOVX sequences are fused to sequences derived from a member of the immunoglobulin protein family. The NOVX-immunoglobulin fusion proteins of the invention  
25 can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between an NOVX ligand and an NOVX protein on the surface of a cell, to thereby suppress NOVX-mediated signal transduction *in vivo*. The NOVX-immunoglobulin fusion proteins can be used to affect the bioavailability of an NOVX cognate ligand. Inhibition of the NOVX ligand/NOVX interaction may be useful therapeutically for both the  
30 treatment of proliferative and differentiative disorders, as well as modulating (*e.g.* promoting or inhibiting) cell survival. Moreover, the NOVX-immunoglobulin fusion proteins of the invention can be used as immunogens to produce anti-NOVX antibodies in a subject, to purify NOVX ligands, and in screening assays to identify molecules that inhibit the interaction of NOVX with an NOVX ligand.



An NOVX chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (*see, e.g.*, Ausubel, *et al.* (eds.) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). An NOVX-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the NOVX protein.

#### **NOVX Agonists and Antagonists**

The invention also pertains to variants of the NOVX proteins that function as either NOVX agonists (*i.e.*, mimetics) or as NOVX antagonists. Variants of the NOVX protein can be generated by mutagenesis (*e.g.*, discrete point mutation or truncation of the NOVX protein). An agonist of the NOVX protein can retain substantially the same, or a subset of, the biological activities of the naturally occurring form of the NOVX protein. An antagonist of the NOVX protein can inhibit one or more of the activities of the naturally occurring form of the NOVX protein by, for example, competitively binding to a downstream or upstream member of a cellular signaling cascade which includes the NOVX protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the NOVX proteins.

Variants of the NOVX proteins that function as either NOVX agonists (*i.e.*, mimetics) or as NOVX antagonists can be identified by screening combinatorial libraries of mutants (*e.g.*, truncation mutants) of the NOVX proteins for NOVX protein agonist or antagonist activity. In one embodiment, a variegated library of NOVX variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene

library. A variegated library of NOVX variants can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential NOVX sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of NOVX sequences therein. There are a variety of methods which can be used to produce libraries of potential NOVX variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential NOVX sequences. Methods for synthesizing degenerate oligonucleotides are well-known within the art. *See, e.g.*, Narang, 1983. *Tetrahedron* 39: 3; Itakura, *et al.*, 1984. *Annu. Rev. Biochem.* 53: 323; Itakura, *et al.*, 1984. *Science* 198: 1056; Ike, *et al.*, 1983. *Nucl. Acids Res.* 11: 477.

## **Polypeptide Libraries**

In addition, libraries of fragments of the NOVX protein coding sequences can be used to generate a variegated population of NOVX fragments for screening and subsequent selection of variants of an NOVX protein. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of an NOVX coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double-stranded DNA that can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with  $S_1$  nuclease, and ligating the resulting fragment library into an expression vector. By this method, expression libraries can be derived which encodes N-terminal and internal fragments of various sizes of the NOVX proteins.

Various techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of NOVX proteins. The most widely used techniques, which are amenable to high throughput analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates

isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique that enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify NOVX variants.

See, e.g., Arkin and Yourvan, 1992. *Proc. Natl. Acad. Sci. USA* 89: 7811-7815; Delgrave, *et*

5 *al.*, 1993. *Protein Engineering* 6:327-331.

### Anti-NOVX Antibodies

Also included in the invention are antibodies to NOVX proteins, or fragments of NOVX proteins. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, i.e., molecules that

10 contain an antigen binding site that specifically binds (immunoreacts with) an antigen. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain,  $F_{ab}$ ,  $F_{ab}'$  and  $F_{(ab)2}$  fragments, and an  $F_{ab}$  expression library. In general, an antibody molecule obtained from humans relates to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes

15 have subclasses as well, such as IgG<sub>1</sub>, IgG<sub>2</sub>, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain. Reference herein to antibodies includes a reference to all such classes, subclasses and types of human antibody species.

An isolated NOVX-related protein of the invention may be intended to serve as an antigen, or a portion or fragment thereof, and additionally can be used as an immunogen to

20 generate antibodies that immunospecifically bind the antigen, using standard techniques for polyclonal and monoclonal antibody preparation. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments of the antigen for use as immunogens. An antigenic peptide fragment comprises at least 6 amino acid residues of the amino acid sequence of the full length protein and encompasses an epitope thereof such that an

25 antibody raised against the peptide forms a specific immune complex with the full length protein or with any fragment that contains the epitope. Preferably, the antigenic peptide comprises at least 10 amino acid residues, or at least 15 amino acid residues, or at least 20 amino acid residues, or at least 30 amino acid residues. Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on its surface; commonly these

30 are hydrophilic regions.

In certain embodiments of the invention, at least one epitope encompassed by the antigenic peptide is a region of NOVX-related protein that is located on the surface of the protein, e.g., a hydrophilic region. A hydrophobicity analysis of the human NOVX-related

protein sequence will indicate which regions of a NOVX-related protein are particularly hydrophilic and, therefore, are likely to encode surface residues useful for targeting antibody production. As a means for targeting antibody production, hydropathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for example, the Kyte Doolittle or the Hopp Woods methods, either with or without Fourier transformation. See, e.g., Hopp and Woods, 1981, *Proc. Nat. Acad. Sci. USA* 78: 3824-3828; Kyte and Doolittle 1982, *J. Mol. Biol.* 157: 105-142, each of which is incorporated herein by reference in its entirety. Antibodies that are specific for one or more domains within an antigenic protein, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

A protein of the invention, or a derivative, fragment, analog, homolog or ortholog thereof, may be utilized as an immunogen in the generation of antibodies that immunospecifically bind these protein components.

Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies directed against a protein of the invention, or against derivatives, fragments, analogs homologs or orthologs thereof (see, for example, *Antibodies: A Laboratory Manual*, Harlow and Lane, 1988, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference). Some of these antibodies are discussed below.

## **20 Polyclonal Antibodies**

For the production of polyclonal antibodies, various suitable host animals (e.g., rabbit, goat, mouse or other mammal) may be immunized by one or more injections with the native protein, a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic preparation can contain, for example, the naturally occurring immunogenic protein, a chemically synthesized polypeptide representing the immunogenic protein, or a recombinantly expressed immunogenic protein. Furthermore, the protein may be conjugated to a second protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. The preparation can further include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to, Freund's (complete and incomplete), mineral gels (e.g., aluminum hydroxide), surface active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), adjuvants usable in humans such as Bacille Calmette-Guerin and *Corynebacterium parvum*, or similar immunostimulatory agents.

Additional examples of adjuvants which can be employed include MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate).

5 The polyclonal antibody molecules directed against the immunogenic protein can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as affinity chromatography using protein A or protein G, which provide primarily the IgG fraction of immune serum. Subsequently, or alternatively, the specific antigen which is the target of the immunoglobulin sought, or an epitope thereof, may be immobilized on a column to purify the immune specific antibody by immunoaffinity chromatography. Purification of immunoglobulins is discussed, for example, by D. Wilkinson  
10 (The Scientist, published by The Scientist, Inc., Philadelphia PA, Vol. 14, No. 8 (April 17, 2000), pp. 25-28).

### **Monoclonal Antibodies**

15 The term "monoclonal antibody" (MAb) or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one molecular species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs thus contain an antigen binding site capable of immunoreacting with a particular epitope of the  
20 antigen characterized by a unique binding affinity for it.

Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically  
25 bind to the immunizing agent. Alternatively, the lymphocytes can be immunized in vitro.

The immunizing agent will typically include the protein antigen, a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell  
30 line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, MONOCLONAL ANTIBODIES: PRINCIPLES AND PRACTICE, Academic Press, (1986) pp. 59-103). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture medium that

preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur et al., MONOCLONAL ANTIBODY PRODUCTION TECHNIQUES AND APPLICATIONS, Marcel Dekker, Inc., New York, (1987) pp. 51-63).

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, *Anal. Biochem.*, 107:220 (1980). Preferably, antibodies having a high degree of specificity and a high binding affinity for the target antigen are isolated.

After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells can be grown in vivo as ascites in a mammal.

The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by

using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, *Nature* 368, 812-13 (1994)) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

#### **Humanized Antibodies**

The antibodies directed against the protein antigens of the invention can further comprise humanized antibodies or human antibodies. These antibodies are suitable for administration to humans without engendering an immune response by the human against the administered immunoglobulin. Humanized forms of antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding subsequences of antibodies) that are principally comprised of the sequence of a human immunoglobulin, and contain minimal sequence derived from a non-human immunoglobulin. Humanization can be performed following the method of Winter and co-workers (Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeyen et al., *Science*, 239:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. (See also U.S. Patent No. 5,225,539.) In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies can also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that

of a human immunoglobulin (Jones et al., 1986; Riechmann et al., 1988; and Presta, *Curr. Op. Struct. Biol.*, 2:593-596 (1992)).

### Human Antibodies

5 Fully human antibodies relate to antibody molecules in which essentially the entire sequences of both the light chain and the heavy chain, including the CDRs, arise from human genes. Such antibodies are termed "human antibodies", or "fully human antibodies" herein. Human monoclonal antibodies can be prepared by the trioma technique; the human B-cell hybridoma technique (see Kozbor, et al., 1983 *Immunol Today* 4: 72) and the EBV hybridoma  
10 technique to produce human monoclonal antibodies (see Cole, et al., 1985 In: MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be produced by using human hybridomas (see Cote, et al., 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus in vitro (see Cole, et al., 1985 In:  
15 MONOCLONAL ANTIBODIES AND CANCER THERAPY, Alan R. Liss, Inc., pp. 77-96).

In addition, human antibodies can also be produced using additional techniques, including phage display libraries (Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the  
20 endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in Marks et al. (*Bio/Technology* 10, 779-783 (1992));  
25 Lonberg et al. (*Nature* 368 856-859 (1994)); Morrison ( *Nature* 368, 812-13 (1994)); Fishwild et al, (*Nature Biotechnology* 14, 845-51 (1996)); Neuberger (*Nature Biotechnology* 14, 826 (1996)); and Lonberg and Huszar (*Intern. Rev. Immunol.* 13 65-93 (1995)).

Human antibodies may additionally be produced using transgenic nonhuman animals which are modified so as to produce fully human antibodies rather than the animal's  
30 endogenous antibodies in response to challenge by an antigen. (See PCT publication WO94/02602). The endogenous genes encoding the heavy and light immunoglobulin chains in the nonhuman host have been incapacitated, and active loci encoding human heavy and light chain immunoglobulins are inserted into the host's genome. The human genes are incorporated, for example, using yeast artificial chromosomes containing the requisite human



DNA segments. An animal which provides all the desired modifications is then obtained as progeny by crossbreeding intermediate transgenic animals containing fewer than the full complement of the modifications. The preferred embodiment of such a nonhuman animal is a mouse, and is termed the Xenomouse<sup>TM</sup> as disclosed in PCT publications WO 96/33735 and  
5 WO 96/34096. This animal produces B cells which secrete fully human immunoglobulins. The antibodies can be obtained directly from the animal after immunization with an immunogen of interest, as, for example, a preparation of a polyclonal antibody, or alternatively from immortalized B cells derived from the animal, such as hybridomas producing monoclonal antibodies. Additionally, the genes encoding the immunoglobulins with human  
10 variable regions can be recovered and expressed to obtain the antibodies directly, or can be further modified to obtain analogs of antibodies such as, for example, single chain Fv molecules.

An example of a method of producing a nonhuman host, exemplified as a mouse, lacking expression of an endogenous immunoglobulin heavy chain is disclosed in U.S. Patent  
15 No. 5,939,598. It can be obtained by a method including deleting the J segment genes from at least one endogenous heavy chain locus in an embryonic stem cell to prevent rearrangement of the locus and to prevent formation of a transcript of a rearranged immunoglobulin heavy chain locus, the deletion being effected by a targeting vector containing a gene encoding a selectable marker; and producing from the embryonic stem cell a transgenic mouse whose somatic and  
20 germ cells contain the gene encoding the selectable marker.

A method for producing an antibody of interest, such as a human antibody, is disclosed in U.S. Patent No. 5,916,771. It includes introducing an expression vector that contains a nucleotide sequence encoding a heavy chain into one mammalian host cell in culture, introducing an expression vector containing a nucleotide sequence encoding a light chain into  
25 another mammalian host cell, and fusing the two cells to form a hybrid cell. The hybrid cell expresses an antibody containing the heavy chain and the light chain.

In a further improvement on this procedure, a method for identifying a clinically relevant epitope on an immunogen, and a correlative method for selecting an antibody that binds immunospecifically to the relevant epitope with high affinity, are disclosed in PCT  
30 publication WO 99/53049.

#### **F<sub>ab</sub> Fragments and Single Chain Antibodies**

According to the invention, techniques can be adapted for the production of single-chain antibodies specific to an antigenic protein of the invention (see e.g., U.S. Patent

No. 4,946,778). In addition, methods can be adapted for the construction of F<sub>ab</sub> expression libraries (see e.g., Huse, et al., 1989 Science 246: 1275-1281) to allow rapid and effective identification of monoclonal F<sub>ab</sub> fragments with the desired specificity for a protein or derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the  
5 idiotype to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F<sub>(ab')<sub>2</sub></sub> fragment produced by pepsin digestion of an antibody molecule; (ii) an F<sub>ab</sub> fragment generated by reducing the disulfide bridges of an F<sub>(ab')<sub>2</sub></sub> fragment; (iii) an F<sub>ab</sub> fragment generated by the treatment of the antibody molecule with papain and a reducing agent and (iv) F<sub>v</sub> fragments.

### **Bispecific Antibodies**

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for an antigenic protein of the invention. The second binding target is  
15 any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, *Nature*, 305:537-539 (1983)). Because of the random  
20 assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, 1991 *EMBO J.*, 10:3655-3659.

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the  
30 fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g.  $F(ab')_2$  bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate  $F(ab')_2$  fragments. These fragments are reduced in the presence of the dithiol compound sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The  $Fab'$  fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the  $Fab'$ -TNB derivatives is then reconverted to the  $Fab'$ -thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other  $Fab'$ -TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Additionally,  $Fab'$  fragments can be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217-225 (1992) describe the production of a fully humanized bispecific antibody  $F(ab')_2$  molecule. Each  $Fab'$  fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the  $Fab'$  portions of two different antibodies by gene fusion. The antibody homodimers were reduced

at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments  
5 comprise a heavy-chain variable domain ( $V_H$ ) connected to a light-chain variable domain ( $V_L$ ) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the  $V_H$  and  $V_L$  domains of one fragment are forced to pair with the complementary  $V_L$  and  $V_H$  domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv  
10 (sFv) dimers has also been reported. See, Gruber et al., *J. Immunol.* 152:5368 (1994).

Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

Exemplary bispecific antibodies can bind to two different epitopes, at least one of which originates in the protein antigen of the invention. Alternatively, an anti-antigenic arm  
15 of an immunoglobulin molecule can be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (Fc $\gamma$ R), such as Fc $\gamma$ RI (CD64), Fc $\gamma$ RII (CD32) and Fc $\gamma$ RIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular antigen. Bispecific antibodies can also be used to direct cytotoxic agents to cells which express a particular  
20 antigen. These antibodies possess an antigen-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the protein antigen described herein and further binds tissue factor (TF).

## 25        **Heteroconjugate Antibodies**

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO  
30 92/200373; EP 03089). It is contemplated that the antibodies can be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate

and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

### **Effector Function Engineering**

5 It can be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and  
10 antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp Med.*, 176: 1191-1195 (1992) and Shopes, *J. Immunol.*, 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff et al. *Cancer Research*, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and can thereby have enhanced complement lysis  
15 and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*, 3: 219-230 (1989).

### **Immunoconjugates**

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active  
20 toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain  
25 (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcumin, croton, saponaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples  
30 include  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ .

Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as

glutareldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in  
5 Vitetta et al., Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody can be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is  
10 administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is in turn conjugated to a cytotoxic agent.

In one embodiment, methods for the screening of antibodies that possess the desired specificity include, but are not limited to, enzyme-linked immunosorbent assay (ELISA) and  
15 other immunologically-mediated techniques known within the art. In a specific embodiment, selection of antibodies that are specific to a particular domain of an NOVX protein is facilitated by generation of hybridomas that bind to the fragment of an NOVX protein possessing such a domain. Thus, antibodies that are specific for a desired domain within an NOVX protein, or derivatives, fragments, analogs or homologs thereof, are also provided  
20 herein.

Anti-NOVX antibodies may be used in methods known within the art relating to the localization and/or quantitation of an NOVX protein (e.g., for use in measuring levels of the NOVX protein within appropriate physiological samples, for use in diagnostic methods, for use in imaging the protein, and the like). In a given embodiment, antibodies for NOVX  
25 proteins, or derivatives, fragments, analogs or homologs thereof, that contain the antibody derived binding domain, are utilized as pharmacologically-active compounds (hereinafter "Therapeutics").

An anti-NOVX antibody (e.g., monoclonal antibody) can be used to isolate an NOVX polypeptide by standard techniques, such as affinity chromatography or immunoprecipitation.  
30 An anti-NOVX antibody can facilitate the purification of natural NOVX polypeptide from cells and of recombinantly-produced NOVX polypeptide expressed in host cells. Moreover, an anti-NOVX antibody can be used to detect NOVX protein (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the NOVX protein. Anti-NOVX antibodies can be used diagnostically to monitor protein levels in tissue

as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{35}\text{S}$  or  $^3\text{H}$ .

#### **NOVX Recombinant Expression Vectors and Host Cells**

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding an NOVX protein, or derivatives, fragments, analogs or homologs thereof. As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively-linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that

the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, that is operatively-linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably-linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner that allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell).

The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Regulatory sequences include those that direct constitutive expression of a nucleotide sequence in many types of host cell and those that direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (*e.g.*, NOVX proteins, mutant forms of NOVX proteins, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for expression of NOVX proteins in prokaryotic or eukaryotic cells. For example, NOVX proteins can be expressed in bacterial cells such as *Escherichia coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *Escherichia coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: (*i*) to increase expression of recombinant protein; (*ii*) to increase the solubility of the recombinant protein; and (*iii*) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the



recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith and Johnson, 1988. *Gene* 67: 31-40), pMAL (New England Biolabs, Beverly, Mass.) and pRIT5 (Pharmacia, Piscataway, N.J.) that fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amrann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990) 60-89).

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein. See, e.g., Gottesman, GENE EXPRESSION TECHNOLOGY: METHODS IN ENZYMOLOGY 185, Academic Press, San Diego, Calif. (1990) 119-128. Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (see, e.g., Wada, *et al.*, 1992. *Nucl. Acids Res.* 20: 2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the NOVX expression vector is a yeast expression vector. Examples of vectors for expression in yeast *Saccharomyces cerevisiae* include pYepSecI (Baldari, *et al.*, 1987. *EMBO J.* 6: 229-234), pMFa (Kurjan and Herskowitz, 1982. *Cell* 30: 933-943), pJRY88 (Schultz *et al.*, 1987. *Gene* 54: 113-123), pYES2 (Invitrogen Corporation, San Diego, Calif.), and picZ (Invitrogen Corp, San Diego, Calif.).

Alternatively, NOVX can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., SF9 cells) include the pAc series (Smith, *et al.*, 1983. *Mol. Cell. Biol.* 3: 2156-2165) and the pVL series (Lucklow and Summers, 1989. *Virology* 170: 31-39).

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, 1987. *Nature* 329: 840) and pMT2PC (Kaufman, *et al.*, 1987. *EMBO J.* 6: 187-195). When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, adenovirus 2, cytomegalovirus, and simian virus 40. For other suitable

expression systems for both prokaryotic and eukaryotic cells see, *e.g.*, Chapters 16 and 17 of Sambrook, *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (*e.g.*, tissue-specific regulatory elements are used to express the nucleic acid). Tissue-specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert, *et al.*, 1987. *Genes Dev.* 1: 268-277), lymphoid-specific promoters (Calame and Eaton, 1988. *Adv. Immunol.* 43: 235-275), in particular promoters of T cell receptors (Winoto and Baltimore, 1989. *EMBO J.* 8: 729-733) and immunoglobulins (Banerji, *et al.*, 1983. *Cell* 33: 729-740; Queen and Baltimore, 1983. *Cell* 33: 741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle, 1989. *Proc. Natl. Acad. Sci. USA* 86: 5473-5477), pancreas-specific promoters (Edlund, *et al.*, 1985. *Science* 230: 912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Pat. No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are also encompassed, *e.g.*, the murine hox promoters (Kessel and Gruss, 1990. *Science* 249: 374-379) and the  $\alpha$ -fetoprotein promoter (Campes and Tilghman, 1989. *Genes Dev.* 3: 537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively-linked to a regulatory sequence in a manner that allows for expression (by transcription of the DNA molecule) of an RNA molecule that is antisense to NOVX mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen that direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen that direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see, *e.g.*, Weintraub, *et al.*, "Antisense RNA as a molecular tool for genetic analysis," *Reviews-Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and

"recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, NOVX protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (MOLECULAR CLONING: A LABORATORY MANUAL. 2nd ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Various selectable markers include those that confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be introduced into a host cell on the same vector as that encoding NOVX or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (*i.e.*, express) NOVX protein. Accordingly, the invention further provides methods for producing NOVX protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding NOVX protein has been introduced) in a suitable

medium such that NOVX protein is produced. In another embodiment, the method further comprises isolating NOVX protein from the medium or the host cell.

### **Transgenic NOVX Animals**

The host cells of the invention can also be used to produce non-human transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which NOVX protein-coding sequences have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous NOVX sequences have been introduced into their genome or homologous recombinant animals in which endogenous NOVX sequences have been altered. Such animals are useful for studying the function and/or activity of NOVX protein and for identifying and/or evaluating modulators of NOVX protein activity. As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene is exogenous DNA that is integrated into the genome of a cell from which a transgenic animal develops and that remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous NOVX gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing NOVX-encoding nucleic acid into the male pronuclei of a fertilized oocyte (*e.g.*, by microinjection, retroviral infection) and allowing the oocyte to develop in a pseudopregnant female foster animal. The human NOVX cDNA sequences SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28 can be introduced as a transgene into the genome of a non-human animal. Alternatively, a non-human homologue of the human NOVX gene, such as a mouse NOVX gene, can be isolated based on hybridization to the human NOVX cDNA (described further *supra*) and used as a transgene. Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific regulatory sequence(s) can be operably-linked to the NOVX transgene to direct expression of NOVX protein to particular cells. Methods for generating transgenic animals via embryo

manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866; 4,870,009; and 4,873,191; and Hogan, 1986. In: MANIPULATING THE MOUSE EMBRYO, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of the NOVX transgene in its genome and/or expression of NOVX mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a transgene-encoding NOVX protein can further be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of an NOVX gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the NOVX gene. The NOVX gene can be a human gene (*e.g.*, the cDNA of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28), but more preferably, is a non-human homologue of a human NOVX gene. For example, a mouse homologue of human NOVX gene of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28 can be used to construct a homologous recombination vector suitable for altering an endogenous NOVX gene in the mouse genome. In one embodiment, the vector is designed such that, upon homologous recombination, the endogenous NOVX gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector).

Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous NOVX gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous NOVX protein). In the homologous recombination vector, the altered portion of the NOVX gene is flanked at its 5'- and 3'-termini by additional nucleic acid of the NOVX gene to allow for homologous recombination to occur between the exogenous NOVX gene carried by the vector and an endogenous NOVX gene in an embryonic stem cell. The additional flanking NOVX nucleic acid is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5'- and 3'-termini) are included in the vector. *See, e.g.*, Thomas, *et al.*, 1987. *Cell* 51: 503 for a description of homologous recombination vectors. The vector is then introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced NOVX

gene has homologously-recombined with the endogenous NOVX gene are selected. *See, e.g., Li, et al., 1992. Cell 69: 915.*

The selected cells are then injected into a blastocyst of an animal (*e.g., a mouse*) to form aggregation chimeras. *See, e.g., Bradley, 1987. In: TERATOCARCINOMAS AND*  
5 *EMBRYONIC STEM CELLS: A PRACTICAL APPROACH, Robertson, ed. IRL, Oxford, pp. 113-152.*  
A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the homologously-recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously-recombined DNA by germline transmission of the transgene. Methods for  
10 constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, 1991. *Curr. Opin. Biotechnol. 2: 823-829*; PCT International Publication Nos.: WO 90/11354; WO 91/01140; WO 92/0968; and WO 93/04169.

In another embodiment, transgenic non-humans animals can be produced that contain selected systems that allow for regulated expression of the transgene. One example of such a  
15 system is the cre/loxP recombinase system of bacteriophage P1. For a description of the cre/loxP recombinase system, *See, e.g., Lakso, et al., 1992. Proc. Natl. Acad. Sci. USA 89: 6232-6236.* Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae*. *See, O'Gorman, et al., 1991. Science 251:1351-1355.* If a cre/loxP recombinase system is used to regulate expression of the transgene, animals containing  
20 transgenes encoding both the Cre recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g., by* mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

Clones of the non-human transgenic animals described herein can also be produced  
25 according to the methods described in Wilmut, *et al., 1997. Nature 385: 810-813.* In brief, a cell (*e.g., a somatic cell*) from the transgenic animal can be isolated and induced to exit the growth cycle and enter G<sub>0</sub> phase. The quiescent cell can then be fused, *e.g., through the use of* electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to  
30 morula or blastocyte and then transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell (*e.g., the somatic cell*) is isolated.

## Pharmaceutical Compositions

The NOVX nucleic acid molecules, NOVX proteins, and anti-NOVX antibodies (also referred to herein as "active compounds") of the invention, and derivatives, fragments, analogs and homologs thereof, can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are not limited to, water, saline, finger's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (*i.e.*, topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF,

Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or  
5 dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various  
10 antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum  
15 monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, an NOVX protein or anti-NOVX antibody) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a  
20 sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, methods of preparation are vacuum drying and freeze-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and  
30 swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a



lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be

achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (*see, e.g.*, U.S. Patent No. 5,328,470) or by stereotactic injection (*see, e.g.*, Chen, *et al.*, 1994. *Proc. Natl. Acad. Sci. USA* 91: 3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells that produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

#### **Screening and Detection Methods**

The isolated nucleic acid molecules of the invention can be used to express NOVX protein (*e.g.*, via a recombinant expression vector in a host cell in gene therapy applications), to detect NOVX mRNA (*e.g.*, in a biological sample) or a genetic lesion in an NOVX gene, and to modulate NOVX activity, as described further, below. In addition, the NOVX proteins can be used to screen drugs or compounds that modulate the NOVX protein activity or expression as well as to treat disorders characterized by insufficient or excessive production of NOVX protein or production of NOVX protein forms that have decreased or aberrant activity compared to NOVX wild-type protein (*e.g.*; diabetes (regulates insulin release); obesity (binds and transport lipids); metabolic disturbances associated with obesity, the metabolic syndrome X as well as anorexia and wasting disorders associated with chronic diseases and various cancers, and infectious disease (possesses anti-microbial activity) and the various dyslipidemias. In addition, the anti-NOVX antibodies of the invention can be used to detect and isolate NOVX proteins and modulate NOVX activity. In yet a further aspect, the invention can be used in methods to influence appetite, absorption of nutrients and the disposition of metabolic substrates in both a positive and negative fashion.

The invention further pertains to novel agents identified by the screening assays described herein and uses thereof for treatments as described, *supra*.

### Screening Assays

The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) that bind to NOVX proteins or have a stimulatory or inhibitory effect on, *e.g.*, NOVX protein expression or NOVX protein activity. The invention also includes compounds identified in the screening assays described herein.

In one embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity of the membrane-bound form of an NOVX protein or polypeptide or biologically-active portion thereof. The test compounds of the invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the "one-bead one-compound" library method; and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds. *See, e.g.*, Lam, 1997. *Anticancer Drug Design* 12: 145.

A "small molecule" as used herein, is meant to refer to a composition that has a molecular weight of less than about 5 kD and most preferably less than about 4 kD. Small molecules can be, *e.g.*, nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids or other organic or inorganic molecules. Libraries of chemical and/or biological mixtures, such as fungal, bacterial, or algal extracts, are known in the art and can be screened with any of the assays of the invention.

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt, *et al.*, 1993. *Proc. Natl. Acad. Sci. U.S.A.* 90: 6909; Erb, *et al.*, 1994. *Proc. Natl. Acad. Sci. U.S.A.* 91: 11422; Zuckermann, *et al.*, 1994. *J. Med. Chem.* 37: 2678; Cho, *et al.*, 1993. *Science* 261: 1303; Carrell, *et al.*, 1994. *Angew. Chem. Int. Ed. Engl.* 33: 2059; Carell, *et al.*, 1994. *Angew. Chem. Int. Ed. Engl.* 33: 2061; and Gallop, *et al.*, 1994. *J. Med. Chem.* 37: 1233.

Libraries of compounds may be presented in solution (*e.g.*, Houghten, 1992. *Biotechniques* 13: 412-421), or on beads (Lam, 1991. *Nature* 354: 82-84), on chips (Fodor, 1993. *Nature* 364: 555-556), bacteria (Ladner, U.S. Patent No. 5,223,409), spores (Ladner, U.S. Patent 5,233,409), plasmids (Cull, *et al.*, 1992. *Proc. Natl. Acad. Sci. USA* 89: 1865-1869) or on phage (Scott and Smith, 1990. *Science* 249: 386-390; Devlin, 1990. *Science*

249: 404-406; Cwirla, *et al.*, 1990. *Proc. Natl. Acad. Sci. U.S.A.* 87: 6378-6382; Felici, 1991. *J. Mol. Biol.* 222: 301-310; Ladner, U.S. Patent No. 5,233,409.).

In one embodiment, an assay is a cell-based assay in which a cell which expresses a membrane-bound form of NOVX protein, or a biologically-active portion thereof, on the cell surface is contacted with a test compound and the ability of the test compound to bind to an NOVX protein determined. The cell, for example, can be of mammalian origin or a yeast cell. Determining the ability of the test compound to bind to the NOVX protein can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the NOVX protein or biologically-active portion thereof can be determined by detecting the labeled compound in a complex. For example, test compounds can be labeled with  $^{125}\text{I}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$ , or  $^3\text{H}$ , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically-labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product. In one embodiment, the assay comprises contacting a cell which expresses a membrane-bound form of NOVX protein, or a biologically-active portion thereof, on the cell surface with a known compound which binds NOVX to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an NOVX protein, wherein determining the ability of the test compound to interact with an NOVX protein comprises determining the ability of the test compound to preferentially bind to NOVX protein or a biologically-active portion thereof as compared to the known compound.

In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a membrane-bound form of NOVX protein, or a biologically-active portion thereof, on the cell surface with a test compound and determining the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the NOVX protein or biologically-active portion thereof. Determining the ability of the test compound to modulate the activity of NOVX or a biologically-active portion thereof can be accomplished, for example, by determining the ability of the NOVX protein to bind to or interact with an NOVX target molecule. As used herein, a "target molecule" is a molecule with which an NOVX protein binds or interacts in nature, for example, a molecule on the surface of a cell which expresses an NOVX interacting protein, a molecule on the surface of a second cell, a molecule in the extracellular milieu, a molecule associated with the internal surface of a cell membrane or a cytoplasmic molecule. An NOVX target molecule can be a non-NOVX molecule or an

NOVX protein or polypeptide of the invention. In one embodiment, an NOVX target molecule is a component of a signal transduction pathway that facilitates transduction of an extracellular signal (*e.g.* a signal generated by binding of a compound to a membrane-bound NOVX molecule) through the cell membrane and into the cell. The target, for example, can be a second intercellular protein that has catalytic activity or a protein that facilitates the association of downstream signaling molecules with NOVX.

Determining the ability of the NOVX protein to bind to or interact with an NOVX target molecule can be accomplished by one of the methods described above for determining direct binding. In one embodiment, determining the ability of the NOVX protein to bind to or interact with an NOVX target molecule can be accomplished by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular  $\text{Ca}^{2+}$ , diacylglycerol,  $\text{IP}_3$ , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (comprising an NOVX-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, cell survival, cellular differentiation, or cell proliferation.

In yet another embodiment, an assay of the invention is a cell-free assay comprising contacting an NOVX protein or biologically-active portion thereof with a test compound and determining the ability of the test compound to bind to the NOVX protein or biologically-active portion thereof. Binding of the test compound to the NOVX protein can be determined either directly or indirectly as described above. In one such embodiment, the assay comprises contacting the NOVX protein or biologically-active portion thereof with a known compound which binds NOVX to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an NOVX protein, wherein determining the ability of the test compound to interact with an NOVX protein comprises determining the ability of the test compound to preferentially bind to NOVX or biologically-active portion thereof as compared to the known compound.

In still another embodiment, an assay is a cell-free assay comprising contacting NOVX protein or biologically-active portion thereof with a test compound and determining the ability of the test compound to modulate (*e.g.* stimulate or inhibit) the activity of the NOVX protein or biologically-active portion thereof. Determining the ability of the test compound to modulate the activity of NOVX can be accomplished, for example, by determining the ability of the NOVX protein to bind to an NOVX target molecule by one of the methods described

above for determining direct binding. In an alternative embodiment, determining the ability of the test compound to modulate the activity of NOVX protein can be accomplished by determining the ability of the NOVX protein further modulate an NOVX target molecule. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate  
5 can be determined as described, *supra*.

In yet another embodiment, the cell-free assay comprises contacting the NOVX protein or biologically-active portion thereof with a known compound which binds NOVX protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an NOVX protein, wherein determining the  
10 ability of the test compound to interact with an NOVX protein comprises determining the ability of the NOVX protein to preferentially bind to or modulate the activity of an NOVX target molecule.

The cell-free assays of the invention are amenable to use of both the soluble form or the membrane-bound form of NOVX protein. In the case of cell-free assays comprising the  
15 membrane-bound form of NOVX protein, it may be desirable to utilize a solubilizing agent such that the membrane-bound form of NOVX protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton<sup>®</sup> X-100, Triton<sup>®</sup> X-114, Thesit<sup>®</sup>,  
20 Isotridecypoly(ethylene glycol ether)<sub>n</sub>, N-dodecyl--N,N-dimethyl-3-ammonio-1-propane sulfonate, 3-(3-cholamidopropyl) dimethylamminiol-1-propane sulfonate (CHAPS), or 3-(3-cholamidopropyl)dimethylamminiol-2-hydroxy-1-propane sulfonate (CHAPSO).

In more than one embodiment of the above assay methods of the invention, it may be desirable to immobilize either NOVX protein or its target molecule to facilitate separation of  
25 complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test compound to NOVX protein, or interaction of NOVX protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and micro-centrifuge tubes. In one embodiment, a  
30 fusion protein can be provided that adds a domain that allows one or both of the proteins to be bound to a matrix. For example, GST-NOVX fusion proteins or GST-target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtiter plates, that are then combined with the test compound or the test compound and either the non-adsorbed target protein or NOVX protein, and the mixture is

incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described, *supra*. Alternatively, the complexes can be dissociated from the matrix, and the level of NOVX protein binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, either the NOVX protein or its target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated NOVX protein or target molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well-known within the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, Ill.), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with NOVX protein or target molecules, but which do not interfere with binding of the NOVX protein to its target molecule, can be derivatized to the wells of the plate, and unbound target or NOVX protein trapped in the wells by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the NOVX protein or target molecule, as well as enzyme-linked assays that rely on detecting an enzymatic activity associated with the NOVX protein or target molecule.

In another embodiment, modulators of NOVX protein expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of NOVX mRNA or protein in the cell is determined. The level of expression of NOVX mRNA or protein in the presence of the candidate compound is compared to the level of expression of NOVX mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of NOVX mRNA or protein expression based upon this comparison. For example, when expression of NOVX mRNA or protein is greater (*i.e.*, statistically significantly greater) in the presence of the candidate compound than in its absence, the candidate compound is identified as a stimulator of NOVX mRNA or protein expression. Alternatively, when expression of NOVX mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of NOVX mRNA or protein expression. The level of NOVX mRNA or protein expression in the cells can be determined by methods described herein for detecting NOVX mRNA or protein.

In yet another aspect of the invention, the NOVX proteins can be used as "bait proteins" in a two-hybrid assay or three hybrid assay (*see, e.g.*, U.S. Patent No. 5,283,317; Zervos, *et al.*, 1993. *Cell* 72: 223-232; Madura, *et al.*, 1993. *J. Biol. Chem.* 268: 12046-12054; Bartel, *et al.*, 1993. *Biotechniques* 14: 920-924; Iwabuchi, *et al.*, 1993. *Oncogene* 8:

5 1693-1696; and Brent WO 94/10300), to identify other proteins that bind to or interact with NOVX ("NOVX-binding proteins" or "NOVX-bp") and modulate NOVX activity. Such NOVX-binding proteins are also likely to be involved in the propagation of signals by the NOVX proteins as, for example, upstream or downstream elements of the NOVX pathway.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. In one construct, the gene that codes for NOVX is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact, *in vivo*, forming an NOVX-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) that is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene that encodes the protein which interacts with NOVX.

The invention further pertains to novel agents identified by the aforementioned screening assays and uses thereof for treatments as described herein.

### Detection Assays

25 Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. By way of example, and not of limitation, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. Some of these applications are described in the subsections, below.



## Chromosome Mapping

Once the sequence (or a portion of the sequence) of a gene has been isolated, this sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the NOVX sequences, SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, or fragments or derivatives thereof, can be used to map the location of the NOVX genes, respectively, on a chromosome. The mapping of the NOVX sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, NOVX genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the NOVX sequences. Computer analysis of the NOVX sequences can be used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the NOVX sequences will yield an amplified fragment.

Somatic cell hybrids are prepared by fusing somatic cells from different mammals (e.g., human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the mouse chromosomes. By using media in which mouse cells cannot grow, because they lack a particular enzyme, but in which human cells can, the one human chromosome that contains the gene encoding the needed enzyme will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, allowing easy mapping of individual genes to specific human chromosomes. *See, e.g., D'Eustachio, et al., 1983. Science 220: 919-924.* Somatic cell hybrids containing only fragments of human chromosomes can also be produced by using human chromosomes with translocations and deletions.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be assigned per day using a single thermal cycler. Using the NOVX sequences to design oligonucleotide primers, sub-localization can be achieved with panels of fragments from specific chromosomes.

Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one

step. Chromosome spreads can be made using cells whose division has been blocked in metaphase by a chemical like colcemid that disrupts the mitotic spindle. The chromosomes can be treated briefly with trypsin, and then stained with Giemsa. A pattern of light and dark bands develops on each chromosome, so that the chromosomes can be identified individually.

- 5 The FISH technique can be used with a DNA sequence as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases, will suffice to get good results at a reasonable amount of time. For a review of this technique, *see*, Verma, *et al.*, HUMAN CHROMOSOMES: A  
10 MANUAL OF BASIC TECHNIQUES (Pergamon Press, New York 1988).

- Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more  
15 likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

- Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, *e.g.*, in McKusick, MENDELIAN INHERITANCE IN MAN, available on-line  
20 through Johns Hopkins University Welch Medical Library). The relationship between genes and disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, *e.g.*, Egeland, *et al.*, 1987. *Nature*, 325: 783-787.

- Moreover, differences in the DNA sequences between individuals affected and  
25 unaffected with a disease associated with the NOVX gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome  
30 spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

### **Tissue Typing**

The NOVX sequences of the invention can also be used to identify individuals from minute biological samples. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. The sequences of the invention are useful as additional DNA markers for RFLP ("restriction fragment length polymorphisms," described in U.S. Patent No. 5,272,057).

Furthermore, the sequences of the invention can be used to provide an alternative technique that determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the NOVX sequences described herein can be used to prepare two PCR primers from the 5'- and 3'-termini of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the invention can be used to obtain such identification sequences from individuals and from tissue. The NOVX sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of about once per each 500 bases. Much of the allelic variation is due to single nucleotide polymorphisms (SNPs), which include restriction fragment length polymorphisms (RFLPs).

Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers that each yield a noncoding amplified sequence of 100 bases. If predicted coding sequences, such as those in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

### **Predictive Medicine**

The invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, pharmacogenomics, and monitoring clinical trials are used for

prognostic (predictive) purposes to thereby treat an individual prophylactically. Accordingly, one aspect of the invention relates to diagnostic assays for determining NOVX protein and/or nucleic acid expression as well as NOVX activity, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant NOVX expression or activity. The disorders include metabolic disorders, diabetes, obesity, infectious disease, anorexia, cancer-associated cachexia, cancer, neurodegenerative disorders, Alzheimer's Disease, Parkinson's Disorder, immune disorders, and hematopoietic disorders, and the various dyslipidemias, metabolic disturbances associated with obesity, the metabolic syndrome X and wasting disorders associated with chronic diseases and various cancers. The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with NOVX protein, nucleic acid expression or activity. For example, mutations in an NOVX gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior to the onset of a disorder characterized by or associated with NOVX protein, nucleic acid expression, or biological activity.

Another aspect of the invention provides methods for determining NOVX protein, nucleic acid expression or activity in an individual to thereby select appropriate therapeutic or prophylactic agents for that individual (referred to herein as "pharmacogenomics"). Pharmacogenomics allows for the selection of agents (*e.g.*, drugs) for therapeutic or prophylactic treatment of an individual based on the genotype of the individual (*e.g.*, the genotype of the individual examined to determine the ability of the individual to respond to a particular agent.)

Yet another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of NOVX in clinical trials.

These and other agents are described in further detail in the following sections.

### **Diagnostic Assays**

An exemplary method for detecting the presence or absence of NOVX in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent capable of detecting NOVX protein or nucleic acid (*e.g.*, mRNA, genomic DNA) that encodes NOVX protein such that the presence of NOVX is detected in the biological sample. An agent for detecting NOVX mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to NOVX mRNA or genomic DNA. The

nucleic acid probe can be, for example, a full-length NOVX nucleic acid, such as the nucleic acid of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, and 28, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to NOVX mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

An agent for detecting NOVX protein is an antibody capable of binding to NOVX protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (e.g., Fab or F(ab')<sub>2</sub>) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody by coupling (i.e., physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently-labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently-labeled streptavidin. The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect NOVX mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of NOVX mRNA include Northern hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of NOVX protein include enzyme linked immunosorbent assays (ELISAs), Western blots, immunoprecipitations, and immunofluorescence. *In vitro* techniques for detection of NOVX genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of NOVX protein include introducing into a subject a labeled anti-NOVX antibody. For example, the antibody can be labeled with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting NOVX protein, mRNA, or genomic DNA, such that the presence of

NOVX protein, mRNA or genomic DNA is detected in the biological sample, and comparing the presence of NOVX protein, mRNA or genomic DNA in the control sample with the presence of NOVX protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of NOVX in a biological sample. For example, the kit can comprise: a labeled compound or agent capable of detecting NOVX protein or mRNA in a biological sample; means for determining the amount of NOVX in the sample; and means for comparing the amount of NOVX in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect NOVX protein or nucleic acid.

### Prognostic Assays

The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant NOVX expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder associated with NOVX protein, nucleic acid expression or activity. Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a disease or disorder. Thus, the invention provides a method for identifying a disease or disorder associated with aberrant NOVX expression or activity in which a test sample is obtained from a subject and NOVX protein or nucleic acid (*e.g.*, mRNA, genomic DNA) is detected, wherein the presence of NOVX protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant NOVX expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid (*e.g.*, serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant NOVX expression or activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a disorder. Thus, the invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant NOVX expression or activity in which a test sample is obtained and NOVX protein or nucleic acid is detected (*e.g.*, wherein the presence of NOVX protein or nucleic acid is diagnostic for a subject that can be

administered the agent to treat a disorder associated with aberrant NOVX expression or activity).

The methods of the invention can also be used to detect genetic lesions in an NOVX gene, thereby determining if a subject with the lesioned gene is at risk for a disorder characterized by aberrant cell proliferation and/or differentiation. In various embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic lesion characterized by at least one of an alteration affecting the integrity of a gene encoding an NOVX-protein, or the misexpression of the NOVX gene. For example, such genetic lesions can be detected by ascertaining the existence of at least one of: (i) a deletion of one or more nucleotides from an NOVX gene; (ii) an addition of one or more nucleotides to an NOVX gene; (iii) a substitution of one or more nucleotides of an NOVX gene, (iv) a chromosomal rearrangement of an NOVX gene; (v) an alteration in the level of a messenger RNA transcript of an NOVX gene, (vi) aberrant modification of an NOVX gene, such as of the methylation pattern of the genomic DNA, (vii) the presence of a non-wild-type splicing pattern of a messenger RNA transcript of an NOVX gene, (viii) a non-wild-type level of an NOVX protein, (ix) allelic loss of an NOVX gene, and (x) inappropriate post-translational modification of an NOVX protein. As described herein, there are a large number of assay techniques known in the art which can be used for detecting lesions in an NOVX gene. A preferred biological sample is a peripheral blood leukocyte sample isolated by conventional means from a subject. However, any biological sample containing nucleated cells may be used, including, for example, buccal mucosal cells.

In certain embodiments, detection of the lesion involves the use of a probe/primer in a polymerase chain reaction (PCR) (*see, e.g.,* U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (*see, e.g.,* Landegran, *et al.*, 1988. *Science* 241: 1077-1080; and Nakazawa, *et al.*, 1994. *Proc. Natl. Acad. Sci. USA* 91: 360-364), the latter of which can be particularly useful for detecting point mutations in the NOVX-gene (*see, Abravaya, et al.*, 1995. *Nucl. Acids Res.* 23: 675-682). This method can include the steps of collecting a sample of cells from a patient, isolating nucleic acid (*e.g.,* genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers that specifically hybridize to an NOVX gene under conditions such that hybridization and amplification of the NOVX gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated

that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (*see*, Guatelli, *et al.*, 1990. *Proc. Natl. Acad. Sci. USA* 87: 1874-1878), transcriptional amplification system (*see*, Kwoh, *et al.*, 1989. *Proc. Natl. Acad. Sci. USA* 86: 1173-1177); Q $\beta$  Replicase (*see*, Lizardi, *et al.*, 1988. *BioTechnology* 6: 1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers.

10 In an alternative embodiment, mutations in an NOVX gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (*see, e.g.*, U.S. Patent  
15 No. 5,493,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In other embodiments, genetic mutations in NOVX can be identified by hybridizing a sample and control nucleic acids, *e.g.*, DNA or RNA, to high-density arrays containing  
20 hundreds or thousands of oligonucleotides probes. *See, e.g.*, Cronin, *et al.*, 1996. *Human Mutation* 7: 244-255; Kozal, *et al.*, 1996. *Nat. Med.* 2: 753-759. For example, genetic mutations in NOVX can be identified in two dimensional arrays containing light-generated DNA probes as described in Cronin, *et al.*, *supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base  
25 changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This is followed by a second hybridization array that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is composed of parallel probe sets, one complementary to the wild-type gene and the  
30 other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the NOVX gene and detect mutations by comparing the sequence of the sample NOVX with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and



Gilbert, 1977. *Proc. Natl. Acad. Sci. USA* 74: 560 or Sanger, 1977. *Proc. Natl. Acad. Sci. USA* 74: 5463. It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays (see, e.g., Naeve, *et al.*, 1995.

*Biotechniques* 19: 448), including sequencing by mass spectrometry (see, e.g., PCT

5 International Publication No. WO 94/16101; Cohen, *et al.*, 1996. *Adv. Chromatography* 36: 127-162; and Griffin, *et al.*, 1993. *Appl. Biochem. Biotechnol.* 38: 147-159).

Other methods for detecting mutations in the NOVX gene include methods in which protection from cleavage agents is used to detect mismatched bases in RNA/RNA or RNA/DNA heteroduplexes. See, e.g., Myers, *et al.*, 1985. *Science* 230: 1242. In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type NOVX sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent that cleaves single-stranded regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S<sub>1</sub> nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, e.g., Cotton, *et al.*, 1988. *Proc. Natl. Acad. Sci. USA* 85: 4397; Saleeba, *et al.*, 1992. *Methods Enzymol.* 217: 286-295. In an embodiment, the control DNA or RNA can be labeled for detection.

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in NOVX cDNAs obtained from samples of cells. For example, the mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches. See, e.g., Hsu, *et al.*, 1994. *Carcinogenesis* 15: 1657-1662. According to an exemplary embodiment, a probe based on an NOVX sequence, e.g., a wild-type NOVX sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, e.g., U.S. Patent No. 5,459,039.

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in NOVX genes. For example, single strand conformation polymorphism (SSCP)

may be used to detect differences in electrophoretic mobility between mutant and wild type nucleic acids. *See, e.g., Orita, et al., 1989. Proc. Natl. Acad. Sci. USA: 86: 2766; Cotton, 1993. Mutat. Res. 285: 125-144; Hayashi, 1992. Genet. Anal. Tech. Appl. 9: 73-79.*

Single-stranded DNA fragments of sample and control NOVX nucleic acids will be denatured  
5 and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In one embodiment,  
10 the subject method utilizes heteroduplex analysis to separate double stranded heteroduplex molecules on the basis of changes in electrophoretic mobility. *See, e.g., Keen, et al., 1991. Trends Genet. 7: 5.*

In yet another embodiment, the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient  
15 gel electrophoresis (DGGE). *See, e.g., Myers, et al., 1985. Nature 313: 495.* When DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA. *See,*  
20 *e.g., Rosenbaum and Reissner, 1987. Biophys. Chem. 265: 12753.*

Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions that permit  
25 hybridization only if a perfect match is found. *See, e.g., Saiki, et al., 1986. Nature 324: 163; Saiki, et al., 1989. Proc. Natl. Acad. Sci. USA 86: 6230.* Such allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

30 Alternatively, allele specific amplification technology that depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization; *see, e.g., Gibbs, et al., 1989. Nucl. Acids Res. 17: 2437-2448*) or at the extreme 3'-terminus of one primer where,

under appropriate conditions, mismatch can prevent, or reduce polymerase extension (*see, e.g.,* Prossner, 1993. *Tibtech.* 11: 238). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection. *See, e.g.,* Gasparini, *et al.*, 1992. *Mol. Cell Probes* 6: 1. It is anticipated that in certain embodiments  
5 amplification may also be performed using *Taq* ligase for amplification. *See, e.g.,* Barany, 1991. *Proc. Natl. Acad. Sci. USA* 88: 189. In such cases, ligation will occur only if there is a perfect match at the 3'-terminus of the 5' sequence, making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing  
10 pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.*, in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving an NOVX gene.

Furthermore, any cell type or tissue, preferably peripheral blood leukocytes, in which  
15 NOVX is expressed may be utilized in the prognostic assays described herein. However, any biological sample containing nucleated cells may be used, including, for example, buccal mucosal cells.

### **Pharmacogenomics**

20 Agents, or modulators that have a stimulatory or inhibitory effect on NOVX activity (*e.g.*, NOVX gene expression), as identified by a screening assay described herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (The disorders include metabolic disorders, diabetes, obesity, infectious disease, anorexia, cancer-associated cachexia, cancer, neurodegenerative disorders, Alzheimer's Disease, Parkinson's  
25 Disorder, immune disorders, and hematopoietic disorders, and the various dyslipidemias, metabolic disturbances associated with obesity, the metabolic syndrome X and wasting disorders associated with chronic diseases and various cancers.) In conjunction with such treatment, the pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) of the individual may  
30 be considered. Differences in metabolism of therapeutics can lead to severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, the pharmacogenomics of the individual permits the selection of effective agents (*e.g.*, drugs) for prophylactic or therapeutic treatments based on a consideration of the individual's genotype. Such pharmacogenomics can further be used to

determine appropriate dosages and therapeutic regimens. Accordingly, the activity of NOVX protein, expression of NOVX nucleic acid, or mutation content of NOVX genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual.

5            Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See *e.g.*, Eichelbaum, 1996. *Clin. Exp. Pharmacol. Physiol.*, 23: 983-985; Linder, 1997. *Clin. Chem.*, 43: 254-266. In general, two types of pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on  
10 the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare defects or as polymorphisms. For example, glucose-6-phosphate dehydrogenase (G6PD) deficiency is a common inherited enzymopathy in which the main clinical complication is hemolysis after ingestion of oxidant drugs (anti-malarials,  
15 sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why  
20 some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been  
25 identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. At the other extreme are the so called ultra-rapid  
30 metabolizers who do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Thus, the activity of NOVX protein, expression of NOVX nucleic acid, or mutation content of NOVX genes in an individual can be determined to thereby select appropriate agent(s) for therapeutic or prophylactic treatment of the individual. In addition,

pharmacogenetic studies can be used to apply genotyping of polymorphic alleles encoding drug-metabolizing enzymes to the identification of an individual's drug responsiveness phenotype. This knowledge, when applied to dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when  
5 treating a subject with an NOVX modulator, such as a modulator identified by one of the exemplary screening assays described herein.

### **Monitoring of Effects During Clinical Trials**

Monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or  
10 activity of NOVX (*e.g.*, the ability to modulate aberrant cell proliferation and/or differentiation) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase NOVX gene expression, protein levels, or upregulate NOVX activity, can be monitored in clinical trials of subjects exhibiting decreased NOVX gene expression, protein  
15 levels, or downregulated NOVX activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease NOVX gene expression, protein levels, or downregulate NOVX activity, can be monitored in clinical trials of subjects exhibiting increased NOVX gene expression, protein levels, or upregulated NOVX activity. In such clinical trials, the expression or activity of NOVX and, preferably, other genes that have been  
20 implicated in, for example, a cellular proliferation or immune disorder can be used as a "read out" or markers of the immune responsiveness of a particular cell.

By way of example, and not of limitation, genes, including NOVX, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) that modulates NOVX activity (*e.g.*, identified in a screening assay as described herein) can be identified.  
25 Thus, to study the effect of agents on cellular proliferation disorders, for example, in a clinical trial, cells can be isolated and RNA prepared and analyzed for the levels of expression of NOVX and other genes implicated in the disorder. The levels of gene expression (*i.e.*, a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods  
30 as described herein, or by measuring the levels of activity of NOVX or other genes. In this manner, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this response state may be determined before, and at various points during, treatment of the individual with the agent.

In one embodiment, the invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, protein, peptide, peptidomimetic, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of an NOVX protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the NOVX protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the NOVX protein, mRNA, or genomic DNA in the pre-administration sample with the NOVX protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of NOVX to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of NOVX to lower levels than detected, *i.e.*, to decrease the effectiveness of the agent.

#### **Methods of Treatment**

The invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant NOVX expression or activity. The disorders include cardiomyopathy, atherosclerosis, hypertension, congenital heart defects, aortic stenosis, atrial septal defect (ASD), atrioventricular (A-V) canal defect, ductus arteriosus, pulmonary stenosis, subaortic stenosis, ventricular septal defect (VSD), valve diseases, tuberous sclerosis, scleroderma, obesity, transplantation, adrenoleukodystrophy, congenital adrenal hyperplasia, prostate cancer, neoplasm; adenocarcinoma, lymphoma, uterus cancer, fertility, hemophilia, hypercoagulation, idiopathic thrombocytopenic purpura, immunodeficiencies, graft versus host disease, AIDS, bronchial asthma, Crohn's disease; multiple sclerosis, treatment of Albright Hereditary Osteodystrophy, and other diseases, disorders and conditions of the like.

These methods of treatment will be discussed more fully, below.

#### **Disease and Disorders**

Diseases and disorders that are characterized by increased (relative to a subject not suffering from the disease or disorder) levels or biological activity may be treated with

Therapeutics that antagonize (*i.e.*, reduce or inhibit) activity. Therapeutics that antagonize activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may be utilized include, but are not limited to: (i) an aforementioned peptide, or analogs, derivatives, fragments or homologs thereof; (ii) antibodies to an aforementioned peptide; (iii) nucleic acids encoding an aforementioned peptide; (iv) administration of antisense nucleic acid and nucleic acids that are "dysfunctional" (*i.e.*, due to a heterologous insertion within the coding sequences of coding sequences to an aforementioned peptide) that are utilized to "knockout" endogenous function of an aforementioned peptide by homologous recombination (see, *e.g.*, Capecchi, 1989. *Science* 244: 1288-1292); or (v) modulators (*i.e.*, inhibitors, agonists and antagonists, including additional peptide mimetic of the invention or antibodies specific to a peptide of the invention) that alter the interaction between an aforementioned peptide and its binding partner.

Diseases and disorders that are characterized by decreased (relative to a subject not suffering from the disease or disorder) levels or biological activity may be treated with Therapeutics that increase (*i.e.*, are agonists to) activity. Therapeutics that upregulate activity may be administered in a therapeutic or prophylactic manner. Therapeutics that may be utilized include, but are not limited to, an aforementioned peptide, or analogs, derivatives, fragments or homologs thereof; or an agonist that increases bioavailability.

Increased or decreased levels can be readily detected by quantifying peptide and/or RNA, by obtaining a patient tissue sample (*e.g.*, from biopsy tissue) and assaying it *in vitro* for RNA or peptide levels, structure and/or activity of the expressed peptides (or mRNAs of an aforementioned peptide). Methods that are well-known within the art include, but are not limited to, immunoassays (*e.g.*, by Western blot analysis, immunoprecipitation followed by sodium dodecyl sulfate (SDS) polyacrylamide gel electrophoresis, immunocytochemistry, etc.) and/or hybridization assays to detect expression of mRNAs (*e.g.*, Northern assays, dot blots, *in situ* hybridization, and the like).

### **Prophylactic Methods**

In one aspect, the invention provides a method for preventing, in a subject, a disease or condition associated with an aberrant NOVX expression or activity, by administering to the subject an agent that modulates NOVX expression or at least one NOVX activity. Subjects at risk for a disease that is caused or contributed to by aberrant NOVX expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation

of symptoms characteristic of the NOVX aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending upon the type of NOVX aberrancy, for example, an NOVX agonist or NOVX antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described  
5 herein. The prophylactic methods of the invention are further discussed in the following subsections.

### Therapeutic Methods

Another aspect of the invention pertains to methods of modulating NOVX expression  
10 or activity for therapeutic purposes. The modulatory method of the invention involves contacting a cell with an agent that modulates one or more of the activities of NOVX protein activity associated with the cell. An agent that modulates NOVX protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring cognate ligand of an NOVX protein, a peptide, an NOVX peptidomimetic, or other small molecule. In  
15 one embodiment, the agent stimulates one or more NOVX protein activity. Examples of such stimulatory agents include active NOVX protein and a nucleic acid molecule encoding NOVX that has been introduced into the cell. In another embodiment, the agent inhibits one or more NOVX protein activity. Examples of such inhibitory agents include antisense NOVX nucleic acid molecules and anti-NOVX antibodies. These modulatory methods can be performed *in*  
20 *vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject). As such, the invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of an NOVX protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or  
25 combination of agents that modulates (e.g., up-regulates or down-regulates) NOVX expression or activity. In another embodiment, the method involves administering an NOVX protein or nucleic acid molecule as therapy to compensate for reduced or aberrant NOVX expression or activity.

Stimulation of NOVX activity is desirable *in situations* in which NOVX is abnormally  
30 downregulated and/or in which increased NOVX activity is likely to have a beneficial effect. One example of such a situation is where a subject has a disorder characterized by aberrant cell proliferation and/or differentiation (e.g., cancer or immune associated disorders). Another example of such a situation is where the subject has a gestational disease (e.g., preeclampsia).



### Determination of the Biological Effect of the Therapeutic

In various embodiments of the invention, suitable *in vitro* or *in vivo* assays are performed to determine the effect of a specific Therapeutic and whether its administration is indicated for treatment of the affected tissue.

5 In various specific embodiments, *in vitro* assays may be performed with representative cells of the type(s) involved in the patient's disorder, to determine if a given Therapeutic exerts the desired effect upon the cell type(s). Compounds for use in therapy may be tested in suitable animal model systems including, but not limited to rats, mice, chicken, cows, monkeys, rabbits, and the like, prior to testing in human subjects. Similarly, for *in vivo* testing, any of the animal model system known in the art may be used prior to administration to human subjects.

### Prophylactic and Therapeutic Uses of the Compositions of the Invention

The NOVX nucleic acids and proteins of the invention are useful in potential prophylactic and therapeutic applications implicated in a variety of disorders including, but not limited to: metabolic disorders, diabetes, obesity, infectious disease, anorexia, cancer-associated cancer, neurodegenerative disorders, Alzheimer's Disease, Parkinson's Disorder, immune disorders, hematopoietic disorders, and the various dyslipidemias, metabolic disturbances associated with obesity, the metabolic syndrome X and wasting disorders associated with chronic diseases and various cancers.

20 As an example, a cDNA encoding the NOVX protein of the invention may be useful in gene therapy, and the protein may be useful when administered to a subject in need thereof. By way of non-limiting example, the compositions of the invention will have efficacy for treatment of patients suffering from: metabolic disorders, diabetes, obesity, infectious disease, anorexia, cancer-associated cachexia, cancer, neurodegenerative disorders, Alzheimer's Disease, Parkinson's Disorder, immune disorders, hematopoietic disorders, and the various dyslipidemias.

Both the novel nucleic acid encoding the NOVX protein, and the NOVX protein of the invention, or fragments thereof, may also be useful in diagnostic applications, wherein the presence or amount of the nucleic acid or the protein are to be assessed. A further use could be as an anti-bacterial molecule (*i.e.*, some peptides have been found to possess anti-bacterial properties). These materials are further useful in the generation of antibodies, which immunospecifically-bind to the novel substances of the invention for use in therapeutic or diagnostic methods.

The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

## Examples

### 5           **Example 1. Identification of NOVX clones**

10           The novel NOVX target sequences identified in the present invention were subjected to the exon linking process to confirm the sequence. PCR primers were designed by starting at the most upstream sequence available, for the forward primer, and at the most downstream sequence available for the reverse primer. Table 11A shows the sequences of the PCR primers used for obtaining different clones. In each case, the sequence was examined, walking inward from the respective termini toward the coding sequence, until a suitable sequence that is either unique or highly selective was encountered, or, in the case of the reverse primer, until the stop codon was reached. Such primers were designed based on in silico predictions for the full length cDNA, part (one or more exons) of the DNA or protein sequence of the target  
15           sequence, or by translated homology of the predicted exons to closely related human sequences from other species. These primers were then employed in PCR amplification based on the following pool of human cDNAs: adrenal gland, bone marrow, brain - amygdala, brain - cerebellum, brain - hippocampus, brain - substantia nigra, brain - thalamus, brain -whole, fetal brain, fetal kidney, fetal liver, fetal lung, heart, kidney, lymphoma - Raji, mammary  
20           gland, pancreas, pituitary gland, placenta, prostate, salivary gland, skeletal muscle, small intestine, spinal cord, spleen, stomach, testis, thyroid, trachea, uterus. Usually the resulting amplicons were gel purified, cloned and sequenced to high redundancy. The PCR product derived from exon linking was cloned into the pCR2.1 vector from Invitrogen. The resulting bacterial clone has an insert covering the entire open reading frame cloned into the pCR2.1  
25           vector. Table 17B shows a list of these bacterial clones. The resulting sequences from all clones were assembled with themselves, with other fragments in CuraGen Corporation's database and with public ESTs. Fragments and ESTs were included as components for an assembly when the extent of their identity with another component of the assembly was at least 95% over 50 bp. In addition, sequence traces were evaluated manually and edited for  
30           corrections if appropriate. These procedures provide the sequence reported herein.

**Table 11A. PCR Primers for Exon Linking**

NOVX Clone	Primer 1 (5' - 3')	SEQ ID NO	Primer 2 (5' - 3')	SEQ ID NO
NOV5	CTCCCACTCCTGCTGCTTCTGACT	98	AAGGCTGGGCTAACCAGTCTCAT	99

NOV7	CATGAAGTGGGCATTCTGCAGG	100	TTATCTGCTGATCTCGCAGGTTATGGA	101
NOV8	CTGACAGGCCCTGGTGTGTGAT	102	TCACACATGTTTCATGTGGGAGTTAGA	103
NOV9	GAGTGAGAGGTCGGACAGACTGTG	104	ACTCATGCAACTTGCTTCTCTCACTCT	105
NOV10b	CCTATGAGCCTGATGCTGGATGAC	106	AGGACTCAGAGGAGGGAGTCTCTGAG	107

Physical clone: Exons were predicted by homology and the intron/exon boundaries were determined using standard genetic rules. Exons were further selected and refined by means of similarity determination using multiple BLAST (for example, tBlastN, BlastX, and BlastN) searches, and, in some instances, GeneScan and Grail. Expressed sequences from both public and proprietary databases were also added when available to further define and complete the gene sequence. The DNA sequence was then manually corrected for apparent inconsistencies thereby obtaining the sequences encoding the full-length protein.

**Table 11B. Physical Clones for PCR products**

NOVX Clone	Bacterial Clone
NOV1a	Physical clone: AC010269
NOV1b	Physical clone: 137043926, 138213196, AC010269.5
NOV1d	Physical clone: 168392429, 138213193
NOV2	Physical clone: 78316254, 123164361
NOV3	Physical clone: GMAC079237.2
NOV4	Physical clone: AP002795.2, AL161453
NOV9	Bacterial clone sggc_draft_ba327p22_20000819.698237.A7
NOV10a	Physical clone: 129297354, AC068190.2, AC036188.2, AC002436.1
NOV10b	Physical clone: 206164528, 206184919, 165210772, 206764887, 162395636

### Example 2. Quantitative expression analysis of clones in various cells and tissues

The quantitative expression of various clones was assessed using microtiter plates containing RNA samples from a variety of normal and pathology-derived cells, cell lines and tissues using real time quantitative PCR (RTQ PCR). RTQ PCR was performed on a Perkin-Elmer Biosystems ABI PRISM® 7700 Sequence Detection System. Various collections of samples are assembled on the plates, and referred to as Panel 1 (containing normal tissues and cancer cell lines), Panel 2 (containing samples derived from tissues from normal and cancer sources), Panel 3 (containing cancer cell lines), Panel 4 (containing cells and cell lines from normal tissues and cells related to inflammatory conditions), Panel 5D/5I (containing human tissues and cell lines with an emphasis on metabolic diseases), Panel CNSD.01 (containing samples from normal and diseased brains) and CNS\_neurodegeneration\_panel (containing samples from normal and diseased brains).

RNA integrity from all samples is controlled for quality by visual assessment of agarose gel electropherograms using 28S and 18S ribosomal RNA staining intensity ratio as a guide (2:1 to 2.5:1 28s:18s) and the absence of low molecular weight RNAs that would be indicative of degradation products. Samples are controlled against genomic DNA

contamination by RTQ PCR reactions run in the absence of reverse transcriptase using probe and primer sets designed to amplify across the span of a single exon.

First, the RNA samples were normalized to reference nucleic acids such as constitutively expressed genes (for example,  $\beta$ -actin and GAPDH). Normalized RNA (5  $\mu$ l) was converted to cDNA and analyzed by RTQ-PCR using One Step RT-PCR Master Mix Reagents (PE Biosystems; Catalog No. 4309169) and gene-specific primers according to the manufacturer's instructions. Probes and primers were designed for each assay according to Perkin Elmer Biosystem's *Primer Express* Software package (version I for Apple Computer's Macintosh Power PC) or a similar algorithm using the target sequence as input. Default settings were used for reaction conditions and the following parameters were set before selecting primers: primer concentration = 250 nM, primer melting temperature ( $T_m$ ) range = 58°-60° C, primer optimal  $T_m$  = 59° C, maximum primer difference = 2° C, probe does not have 5' G, probe  $T_m$  must be 10° C greater than primer  $T_m$ , amplicon size 75 bp to 100 bp. The probes and primers selected (see below) were synthesized by Synthegen (Houston, TX, USA). Probes were double purified by HPLC to remove uncoupled dye and evaluated by mass spectroscopy to verify coupling of reporter and quencher dyes to the 5' and 3' ends of the probe, respectively. Their final concentrations were: forward and reverse primers, 900 nM each, and probe, 200nM.

PCR conditions: Normalized RNA from each tissue and each cell line was spotted in each well of a 96 well PCR plate (Perkin Elmer Biosystems). PCR cocktails including two probes (a probe specific for the target clone and another gene-specific probe multiplexed with the target probe) were set up using 1X TaqMan™ PCR Master Mix for the PE Biosystems 7700, with 5 mM MgCl<sub>2</sub>, dNTPs (dA, G, C, U at 1:1:1:2 ratios), 0.25 U/ml AmpliTaq Gold™ (PE Biosystems), and 0.4 U/ $\mu$ l RNase inhibitor, and 0.25 U/ $\mu$ l reverse transcriptase. Reverse transcription was performed at 48° C for 30 minutes followed by amplification/PCR cycles as follows: 95° C 10 min, then 40 cycles of 95° C for 15 seconds, 60° C for 1 minute. Results were recorded as CT values (cycle at which a given sample crosses a threshold level of fluorescence) using a log scale, with the difference in RNA concentration between a given sample and the sample with the lowest CT value being represented as 2 to the power of delta CT. The percent relative expression is then obtained by taking the reciprocal of this RNA difference and multiplying by 100.

#### **Panels 1, 1.1, 1.2, and 1.3D**

The plates for Panels 1, 1.1, 1.2 and 1.3D include 2 control wells (genomic DNA control and chemistry control) and 94 wells containing cDNA from various samples. The samples in these panels are broken into 2 classes: samples derived from cultured cell lines and samples derived from primary normal tissues. The cell lines are derived from cancers of the following types: lung cancer, breast cancer, melanoma, colon cancer, prostate cancer, CNS cancer, squamous cell carcinoma, ovarian cancer, liver cancer, renal cancer, gastric cancer and pancreatic cancer. Cell lines used in these panels are widely available through the American Type Culture Collection (ATCC), a repository for cultured cell lines, and were cultured using the conditions recommended by the ATCC. The normal tissues found on these panels are comprised of samples derived from all major organ systems from single adult individuals or fetuses. These samples are derived from the following organs: adult skeletal muscle, fetal skeletal muscle, adult heart, fetal heart, adult kidney, fetal kidney, adult liver, fetal liver, adult lung, fetal lung, various regions of the brain, the spleen, bone marrow, lymph node, pancreas, salivary gland, pituitary gland, adrenal gland, spinal cord, thymus, stomach, small intestine, colon, bladder, trachea, breast, ovary, uterus, placenta, prostate, testis and adipose.

In the results for Panels 1, 1.1, 1.2 and 1.3D, the following abbreviations are used:

ca. = carcinoma,

\* = established from metastasis,

met = metastasis,

s cell var = small cell variant,

non-s = non-sm = non-small,

squam = squamous,

pl. eff = pl effusion = pleural effusion,

glio = glioma,

astro = astrocytoma, and

neuro = neuroblastoma.

#### **GENERAL\_SCREENING\_PANEL\_V1.4**

The plates for Panel 1.4 include 2 control wells (genomic DNA control and chemistry control) and 94 wells containing cDNA from various samples. The samples in Panel 1.4 are broken into 2 classes: samples derived from cultured cell lines and samples derived from primary normal tissues. The cell lines are derived from cancers of the following types: lung cancer, breast cancer, melanoma, colon cancer, prostate cancer, CNS cancer, squamous cell

carcinoma, ovarian cancer, liver cancer, renal cancer, gastric cancer and pancreatic cancer. Cell lines used in Panel 1.4 are widely available through the American Type Culture Collection (ATCC), a repository for cultured cell lines, and were cultured using the conditions recommended by the ATCC. The normal tissues found on Panel 1.4 are comprised of pools of samples derived from all major organ systems from 2 to 5 different adult individuals or fetuses. These samples are derived from the following organs: adult skeletal muscle, fetal skeletal muscle, adult heart, fetal heart, adult kidney, fetal kidney, adult liver, fetal liver, adult lung, fetal lung, various regions of the brain, the spleen, bone marrow, lymph node, pancreas, salivary gland, pituitary gland, adrenal gland, spinal cord, thymus, stomach, small intestine, colon, bladder, trachea, breast, ovary, uterus, placenta, prostate, testis and adipose.

### **Panels 2D and 2.2**

The plates for Panels 2D and 2.2 generally include 2 control wells and 94 test samples composed of RNA or cDNA isolated from human tissue procured by surgeons working in close cooperation with the National Cancer Institute's Cooperative Human Tissue Network (CHTN) or the National Disease Research Initiative (NDRI). The tissues are derived from human malignancies and in cases where indicated many malignant tissues have "matched margins" obtained from noncancerous tissue just adjacent to the tumor. These are termed normal adjacent tissues and are denoted "NAT" in the results below. The tumor tissue and the "matched margins" are evaluated by two independent pathologists (the surgical pathologists and again by a pathologists at NDRI or CHTN). This analysis provides a gross histopathological assessment of tumor differentiation grade. Moreover, most samples include the original surgical pathology report that provides information regarding the clinical stage of the patient. These matched margins are taken from the tissue surrounding (i.e. immediately proximal) to the zone of surgery (designated "NAT", for normal adjacent tissue, in Table RR). In addition, RNA and cDNA samples were obtained from various human tissues derived from autopsies performed on elderly people or sudden death victims (accidents, etc.). These tissues were ascertained to be free of disease and were purchased from various commercial sources such as Clontech (Palo Alto, CA), Research Genetics, and Invitrogen.

### **PANEL 3D**

The plates of Panel 3D are comprised of 94 cDNA samples and two control samples. Specifically, 92 of these samples are derived from cultured human cancer cell lines, 2 samples

of human primary cerebellar tissue and 2 controls. The human cell lines are generally obtained from ATCC (American Type Culture Collection), NCI or the German tumor cell bank and fall into the following tissue groups: Squamous cell carcinoma of the tongue, breast cancer, prostate cancer, melanoma, epidermoid carcinoma, sarcomas, bladder carcinomas, pancreatic cancers, kidney cancers, leukemias/lymphomas, ovarian/uterine/cervical, gastric, colon, lung and CNS cancer cell lines. In addition, there are two independent samples of cerebellum. These cells are all cultured under standard recommended conditions and RNA extracted using the standard procedures. The cell lines in panel 3D and 1.3D are of the most common cell lines used in the scientific literature.

#### **Panels 4D, 4R, and 4.1D**

Panel 4 includes samples on a 96 well plate (2 control wells, 94 test samples) composed of RNA (Panel 4R) or cDNA (Panels 4D/4.1D) isolated from various human cell lines or tissues related to inflammatory conditions. Total RNA from control normal tissues such as colon and lung (Stratagene, La Jolla, CA) and thymus and kidney (Clontech) were employed. Total RNA from liver tissue from cirrhosis patients and kidney from lupus patients was obtained from BioChain (Biochain Institute, Inc., Hayward, CA). Intestinal tissue for RNA preparation from patients diagnosed as having Crohn's disease and ulcerative colitis was obtained from the National Disease Research Interchange (NDRI) (Philadelphia, PA).

Astrocytes, lung fibroblasts, dermal fibroblasts, coronary artery smooth muscle cells, small airway epithelium, bronchial epithelium, microvascular dermal endothelial cells, microvascular lung endothelial cells, human pulmonary aortic endothelial cells, human umbilical vein endothelial cells were all purchased from Clonetics (Walkersville, MD) and grown in the media supplied for these cell types by Clonetics. These primary cell types were activated with various cytokines or combinations of cytokines for 6 and/or 12-14 hours, as indicated. The following cytokines were used; IL-1 beta at approximately 1-5 ng/ml, TNF alpha at approximately 5-10 ng/ml, IFN gamma at approximately 20-50 ng/ml, IL-4 at approximately 5-10 ng/ml, IL-9 at approximately 5-10 ng/ml, IL-13 at approximately 5-10 ng/ml. Endothelial cells were sometimes starved for various times by culture in the basal media from Clonetics with 0.1% serum.

Mononuclear cells were prepared from blood of employees at CuraGen Corporation, using Ficoll. LAK cells were prepared from these cells by culture in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco/Life Technologies, Rockville, MD), 1

mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), and 10 mM Hepes (Gibco) and Interleukin 2 for 4-6 days. Cells were then either activated with 10-20 ng/ml PMA and 1-2  $\mu$ g/ml ionomycin, IL-12 at 5-10 ng/ml, IFN gamma at 20-50 ng/ml and IL-18 at 5-10 ng/ml for 6 hours. In some cases, mononuclear cells were cultured for 4-5 days in  
5 DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), and 10 mM Hepes (Gibco) with PHA (phytohemagglutinin) or PWM (pokeweed mitogen) at approximately 5  $\mu$ g/ml. Samples were taken at 24, 48 and 72 hours for RNA preparation. MLR (mixed lymphocyte reaction) samples were obtained by taking blood from two donors, isolating the mononuclear cells using  
10 Ficoll and mixing the isolated mononuclear cells 1:1 at a final concentration of approximately  $2 \times 10^6$  cells/ml in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol ( $5.5 \times 10^{-5}$  M) (Gibco), and 10 mM Hepes (Gibco). The MLR was cultured and samples taken at various time points ranging from 1- 7 days for RNA preparation.

15 Monocytes were isolated from mononuclear cells using CD14 Miltenyi Beads, +ve VS selection columns and a Vario Magnet according to the manufacturer's instructions. Monocytes were differentiated into dendritic cells by culture in DMEM 5% fetal calf serum (FCS) (Hyclone, Logan, UT), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), and 10 mM Hepes (Gibco), 50 ng/ml  
20 GMCSF and 5 ng/ml IL-4 for 5-7 days. Macrophages were prepared by culture of monocytes for 5-7 days in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), 10 mM Hepes (Gibco) and 10% AB Human Serum or MCSF at approximately 50 ng/ml. Monocytes, macrophages and dendritic cells were stimulated for 6 and 12-14 hours with lipopolysaccharide (LPS) at 100  
25 ng/ml. Dendritic cells were also stimulated with anti-CD40 monoclonal antibody (Pharmingen) at 10  $\mu$ g/ml for 6 and 12-14 hours.

CD4 lymphocytes, CD8 lymphocytes and NK cells were also isolated from mononuclear cells using CD4, CD8 and CD56 Miltenyi beads, positive VS selection columns and a Vario Magnet according to the manufacturer's instructions. CD45RA and CD45RO CD4  
30 lymphocytes were isolated by depleting mononuclear cells of CD8, CD56, CD14 and CD19 cells using CD8, CD56, CD14 and CD19 Miltenyi beads and positive selection. Then CD45RO beads were used to isolate the CD45RO CD4 lymphocytes with the remaining cells being CD45RA CD4 lymphocytes. CD45RA CD4, CD45RO CD4 and CD8 lymphocytes were placed in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM



sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), and 10 mM Hepes (Gibco) and plated at  $10^6$  cells/ml onto Falcon 6 well tissue culture plates that had been coated overnight with 0.5  $\mu$ g/ml anti-CD28 (Pharmingen) and 3  $\mu$ g/ml anti-CD3 (OKT3, ATCC) in PBS. After 6 and 24 hours, the cells were harvested for RNA preparation. To prepare  
5 chronically activated CD8 lymphocytes, we activated the isolated CD8 lymphocytes for 4 days on anti-CD28 and anti-CD3 coated plates and then harvested the cells and expanded them in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), and 10 mM Hepes (Gibco) and IL-2. The expanded CD8 cells were then activated again with plate bound anti-CD3 and anti-CD28  
10 for 4 days and expanded as before. RNA was isolated 6 and 24 hours after the second activation and after 4 days of the second expansion culture. The isolated NK cells were cultured in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), and 10 mM Hepes (Gibco) and IL-2 for 4-6 days before RNA was prepared.

15 To obtain B cells, tonsils were procured from NDRI. The tonsil was cut up with sterile dissecting scissors and then passed through a sieve. Tonsil cells were then spun down and resuspended at  $10^6$  cells/ml in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), and 10 mM Hepes (Gibco). To activate the cells, we used PWM at 5  $\mu$ g/ml or anti-CD40 (Pharmingen) at  
20 approximately 10  $\mu$ g/ml and IL-4 at 5-10 ng/ml. Cells were harvested for RNA preparation at 24, 48 and 72 hours.

To prepare the primary and secondary Th1/Th2 and Tr1 cells, six-well Falcon plates were coated overnight with 10  $\mu$ g/ml anti-CD28 (Pharmingen) and 2  $\mu$ g/ml OKT3 (ATCC), and then washed twice with PBS. Umbilical cord blood CD4 lymphocytes (Poietic Systems,  
25 German Town, MD) were cultured at  $10^5$ - $10^6$  cells/ml in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), 10 mM Hepes (Gibco) and IL-2 (4 ng/ml). IL-12 (5 ng/ml) and anti-IL4 (1 g/ml) were used to direct to Th1, while IL-4 (5 ng/ml) and anti-IFN gamma (1 g/ml) were used to direct to Th2 and IL-10 at 5 ng/ml was used to direct to Tr1. After 4-5 days, the  
30 activated Th1, Th2 and Tr1 lymphocytes were washed once in DMEM and expanded for 4-7 days in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), 10 mM Hepes (Gibco) and IL-2 (1 ng/ml). Following this, the activated Th1, Th2 and Tr1 lymphocytes were re-stimulated for 5 days with anti-CD28/OKT3 and cytokines as described above, but with the addition of anti-

CD95L (1  $\mu$ g/ml) to prevent apoptosis. After 4-5 days, the Th1, Th2 and Tr1 lymphocytes were washed and then expanded again with IL-2 for 4-7 days. Activated Th1 and Th2 lymphocytes were maintained in this way for a maximum of three cycles. RNA was prepared from primary and secondary Th1, Th2 and Tr1 after 6 and 24 hours following the second and third activations with plate bound anti-CD3 and anti-CD28 mAbs and 4 days into the second and third expansion cultures in Interleukin 2.

The following leukocyte cells lines were obtained from the ATCC: Ramos, EOL-1, KU-812. EOL cells were further differentiated by culture in 0.1 mM dbcAMP at  $5 \times 10^5$  cells/ml for 8 days, changing the media every 3 days and adjusting the cell concentration to  $5 \times 10^5$  cells/ml. For the culture of these cells, we used DMEM or RPMI (as recommended by the ATCC), with the addition of 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), 10 mM Hepes (Gibco). RNA was either prepared from resting cells or cells activated with PMA at 10 ng/ml and ionomycin at 1  $\mu$ g/ml for 6 and 14 hours. Keratinocyte line CCD106 and an airway epithelial tumor line NCI-H292 were also obtained from the ATCC. Both were cultured in DMEM 5% FCS (Hyclone), 100  $\mu$ M non essential amino acids (Gibco), 1 mM sodium pyruvate (Gibco), mercaptoethanol  $5.5 \times 10^{-5}$  M (Gibco), and 10 mM Hepes (Gibco). CCD1106 cells were activated for 6 and 14 hours with approximately 5 ng/ml TNF alpha and 1 ng/ml IL-1 beta, while NCI-H292 cells were activated for 6 and 14 hours with the following cytokines: 5 ng/ml IL-4, 5 ng/ml IL-9, 5 ng/ml IL-13 and 25 ng/ml IFN gamma.

For these cell lines and blood cells, RNA was prepared by lysing approximately  $10^7$  cells/ml using Trizol (Gibco BRL). Briefly, 1/10 volume of bromochloropropane (Molecular Research Corporation) was added to the RNA sample, vortexed and after 10 minutes at room temperature, the tubes were spun at 14,000 rpm in a Sorvall SS34 rotor. The aqueous phase was removed and placed in a 15 ml Falcon Tube. An equal volume of isopropanol was added and left at -20 degrees C overnight. The precipitated RNA was spun down at 9,000 rpm for 15 min in a Sorvall SS34 rotor and washed in 70% ethanol. The pellet was redissolved in 300  $\mu$ l of RNase-free water and 35  $\mu$ l buffer (Promega) 5  $\mu$ l DTT, 7  $\mu$ l RNasin and 8  $\mu$ l DNase were added. The tube was incubated at 37 degrees C for 30 minutes to remove contaminating genomic DNA, extracted once with phenol chloroform and re-precipitated with 1/10 volume of 3 M sodium acetate and 2 volumes of 100% ethanol. The RNA was spun down and placed in RNase free water. RNA was stored at -80 degrees C.

## Panel CNSD.01

5 The plates for Panel CNSD.01 include two control wells and 94 test samples  
comprised of cDNA isolated from postmortem human brain tissue obtained from the Harvard  
Brain Tissue Resource Center. Brains are removed from calvaria of donors between 4 and 24  
hours after death, sectioned by neuroanatomists, and frozen at -80°C in liquid nitrogen vapor.  
All brains are sectioned and examined by neuropathologists to confirm diagnoses with clear  
associated neuropathology.

10 Disease diagnoses are taken from patient records. The panel contains two brains from  
each of the following diagnoses: Alzheimer's disease, Parkinson's disease, Huntington's  
disease, Progressive Supranuclear Palsy, Depression, and "Normal controls". Within each of  
these brains, the following regions are represented: cingulate gyrus, temporal pole, globus  
palladus, substantia nigra, Brodman Area 4 (primary motor strip), Brodman Area 7 (parietal  
cortex), Brodman Area 9 (prefrontal cortex), and Brodman area 17 (occipital cortex). Not all  
15 brain regions are represented in all cases; e.g., Huntington's disease is characterized in part by  
neurodegeneration in the globus palladus, thus this region is impossible to obtain from  
confirmed Huntington's cases. Likewise Parkinson's disease is characterized by degeneration  
of the substantia nigra making this region more difficult to obtain. Normal control brains were  
examined for neuropathology and found to be free of any pathology consistent with  
20 neurodegeneration.

In the labels employed to identify tissues in the CNS panel, the following abbreviations  
are used:

**PSP = Progressive supranuclear palsy**  
25 Sub Nigra = Substantia nigra  
Glob Palladus= Globus palladus  
Temp Pole = Temporal pole  
Cing Gyr = Cingulate gyrus  
BA 4 = Brodman Area 4

30

## Panel CNS\_Neurodegeneration\_V1.0

The plates for Panel CNS\_Neurodegeneration\_V1.0 include two control wells and 47  
test samples comprised of cDNA isolated from postmortem human brain tissue obtained from  
35 the Harvard Brain Tissue Resource Center (McLean Hospital) and the Human Brain and

Spinal Fluid Resource Center (VA Greater Los Angeles Healthcare System). Brains are removed from calvaria of donors between 4 and 24 hours after death, sectioned by neuroanatomists, and frozen at -80°C in liquid nitrogen vapor. All brains are sectioned and examined by neuropathologists to confirm diagnoses with clear associated neuropathology.

5           Disease diagnoses are taken from patient records. The panel contains six brains from Alzheimer's disease (AD) patients, and eight brains from "Normal controls" who showed no evidence of dementia prior to death. The eight normal control brains are divided into two categories: Controls with no dementia and no Alzheimer's like pathology (Controls) and controls with no dementia but evidence of severe Alzheimer's like pathology, (specifically  
10   senile plaque load rated as level 3 on a scale of 0-3; 0 = no evidence of plaques, 3 = severe AD senile plaque load). Within each of these brains, the following regions are represented: hippocampus, temporal cortex (Brodmann Area 21), parietal cortex (Brodmann area 7), and occipital cortex (Brodmann area 17). These regions were chosen to encompass all levels of neurodegeneration in AD. The hippocampus is a region of early and severe neuronal loss in  
15   AD; the temporal cortex is known to show neurodegeneration in AD after the hippocampus; the parietal cortex shows moderate neuronal death in the late stages of the disease; the occipital cortex is spared in AD and therefore acts as a "control" region within AD patients. Not all brain regions are represented in all cases.

20           In the labels employed to identify tissues in the CNS\_Neurodegeneration\_V1.0 panel, the following abbreviations are used:

**AD = Alzheimer's disease brain; patient was demented and showed AD-like pathology upon autopsy**

Control = Control brains; patient not demented, showing no neuropathology

25           Control (Path) = Control brains; patient not demented but showing severe AD-like pathology

SupTemporal Ctx = Superior Temporal Cortex

Inf Temporal Ctx = Inferior Temporal Cortex

30           **NOV1a and NOV1d**

            Expression of gene NOV1a and variant NOV1a was assessed using the primer-probe sets Ag4164, Ag1313b, Ag2197, and Ag708 described in Tables 12, 13, 14, and 15. Please  
35   note that Ag4164 contains a single mismatch in the probe relative to the NOV1a and NOV1d

sequences. This mismatch is not predicted to alter the RTQ-PCR results. Results from RTQ-PCR runs are shown in Tables 16, 17, 18, and 19.

**Table 12. Probe Name Ag4164**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-GCACTACAAGTGAAGCCTTAC-3'	58.1	22	822	108
Probe	FAM-5'-CTCAAGTAGAAGCCGACTTATGCAAA-3'-TAMRA	64	26	845	109
Reverse	5'-TCAAATCCTTCTGCGATACAGT-3'	58.9	22	875	110

5

**Table 13. Probe Name Ag1313b**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-CAGCTGCACGATTAATGAAGAT-3'	59.4	22	264	111
Probe	TET-5'-AGGTCTTGGACTGGCCTTCACCATT-3'-TAMRA	69	25	288	112
Reverse	5'-CCAAAGTTGTGTCCAGACTCAT-3'	59.1	22	317	113

**Table 14. Probe Name Ag2197**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-CCAAGGAAGACCTCTTCATCTT-3'	58.8	22	1022	114
Probe	FAM-5'-TCTTGCTTACGGCATAAGCGCTCTCT-3'-TAMRA	69	26	1060	115
Reverse	5'-TTCATTTCTATGGGACCTCAGA-3'	58.7	22	1086	116

**Table 15. Probe Name Ag708**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-AAAGATGGGACTCGTCATGAC-3'	59	21	232	117
Probe	TET-5'-CACGCCATCTTACTGACTGGTCTGGA-3'-TAMRA	69.5	26	253	118
Reverse	5'-GTGCAAATCCCAAAGTGCA-3'	59.5	20	306	119

10

**Table 16. Panel 1.2**

Tissue Name	Relative Expression(%)	
	1.2tm888t_ ag708	1.2tm1047t_ ag708
Endothelial cells	0.0	0.0
Heart (fetal)	1.2	0.0
Pancreas	22.5	29.5
Pancreatic ca. CAPAN 2	0.0	0.0
Adrenal Gland (new lot*)	0.9	0.0
Thyroid	0.5	0.0
Salivary gland	0.4	0.0

Pituitary gland	1.3	0.0
Brain (fetal)	4.0	0.0
Brain (whole)	7.9	6.5
Brain (amygdala)	2.1	0.0
Brain (cerebellum)	16.7	23.3
Brain (hippocampus)	4.6	3.1
Brain (thalamus)	1.5	0.0
Cerebral Cortex	10.7	5.8
Spinal cord	1.1	0.0
CNS ca. (glio/astro) U87-MG	4.4	11.7
CNS ca. (glio/astro) U-118-MG	0.6	0.1
CNS ca. (astro) SW1783	1.2	0.7
CNS ca.* (neuro; met ) SK-N-AS	0.2	0.0
CNS ca. (astro) SF-539	1.6	1.1
CNS ca. (astro) SNB-75	7.7	1.7
CNS ca. (glio) SNB-19	9.5	4.3
CNS ca. (glio) U251	<b>100.0</b>	<b>100.0</b>
CNS ca. (glio) SF-295	2.4	0.0
Heart	4.9	0.0
Skeletal Muscle (new lot*)	3.1	0.0
Bone marrow	0.0	0.0
Thymus	0.2	0.0
Spleen	0.0	0.0
Lymph node	0.2	0.0
Colorectal	0.0	0.0
Stomach	1.2	0.0
Small intestine	1.4	0.0
Colon ca. SW480	1.5	0.0
Colon ca.* (SW480 met)SW620	0.0	0.0
Colon ca. HT29	0.0	0.0
Colon ca. HCT-116	0.7	0.0
Colon ca. CaCo-2	0.0	0.0
83219 CC Well to Mod Diff (ODO3866)	0.1	0.0
Colon ca. HCC-2998	0.0	0.0
Gastric ca.* (liver met) NCI-N87	4.1	0.1
Bladder	3.5	0.0
Trachea	0.3	0.0
Kidney	5.4	10.7
Kidney (fetal)	37.9	94.0
Renal ca. 786-0	4.6	5.3
Renal ca. A498	12.2	20.2
Renal ca. RXF 393	3.3	4.2
Renal ca. ACHN	2.8	1.1
Renal ca. UO-31	2.3	0.2
Renal ca. TK-10	7.6	12.7

Liver	0.9	0.0
Liver (fetal)	0.2	0.0
Liver ca. (hepatoblast) HepG2	0.0	0.0
Lung	1.1	0.0
Lung (fetal)	1.2	0.0
Lung ca. (small cell) LX-1	0.0	0.0
Lung ca. (small cell) NCI-H69	0.1	0.0
Lung ca. (s.cell var.) SHP-77	0.0	0.0
Lung ca. (large cell) NCI-H460	0.0	0.0
Lung ca. (non-sm. cell) A549	0.5	0.0
Lung ca. (non-s.cell) NCI-H23	0.0	0.0
Lung ca (non-s.cell) HOP-62	4.9	1.1
Lung ca. (non-s.cl) NCI-H522	0.0	0.0
Lung ca. (squam.) SW 900	3.7	2.0
Lung ca. (squam.) NCI-H596	0.0	0.0
Mammary gland	3.1	4.5
Breast ca.* (pl. effusion) MCF-7	0.0	0.0
Breast ca.* (pl.ef) MDA-MB-231	3.1	6.2
Breast ca.* (pl. effusion) T47D	0.8	0.0
Breast ca. BT-549	0.7	0.0
Breast ca. MDA-N	10.7	14.6
Ovary	26.4	39.5
Ovarian ca. OVCAR-3	0.4	0.0
Ovarian ca. OVCAR-4	0.9	0.0
Ovarian ca. OVCAR-5	6.0	1.7
Ovarian ca. OVCAR-8	0.7	0.0
Ovarian ca. IGROV-1	8.1	15.2
Ovarian ca.* (ascites) SK-OV-3	2.9	0.0
Uterus	0.7	0.0
Placenta	7.1	0.0
Prostate	0.3	0.0
Prostate ca.* (bone met) PC-3	1.5	0.0
Testis	3.3	0.0
Melanoma Hs688(A).T	0.0	0.0
Melanoma* (met) Hs688(B).T	0.0	0.0
Melanoma UACC-62	0.0	0.0
Melanoma M14	0.0	0.0
Melanoma LOX IMVI	1.5	0.0
Melanoma* (met) SK-MEL-5	0.0	0.0

Table 17. Panel 1.3D

Tissue Name	Relative Expression(%)	Relative Expression(%)	Relative Expression(%)
	1.3dtm4180f am_ag2197	1.3dx4tm5365 t_ag1313b_b1	1.3dtm3301t_ ag708
Liver adenocarcinoma	2.6	0.5	2.3
Pancreas	2.8	1.4	4.4
Pancreatic ca. CAPAN 2	0.6	0.2	0.8
Adrenal gland	0.2	0.0	0.0
Thyroid	0.2	0.1	0.0
Salivary gland	0.0	0.2	0.2
Pituitary gland	0.0	0.0	0.0
Brain (fetal)	0.2	0.1	0.7
Brain (whole)	3.4	2.0	2.5
Brain (amygdala)	1.8	1.1	1.7
Brain (cerebellum)	7.4	8.0	11.2
Brain (hippocampus)	7.6	1.0	7.9
Brain (substantia nigra)	0.0	0.0	0.0
Brain (thalamus)	0.2	0.1	0.0
Cerebral Cortex	5.6	0.5	4.6
Spinal cord	0.2	0.2	0.8
CNS ca. (glio/astro) U87-MG	7.0	1.8	4.6
CNS ca. (glio/astro) U-118-MG	12.2	2.3	7.3
CNS ca. (astro) SW1783	4.9	2.0	5.7
CNS ca.* (neuro; met ) SK-N-AS	0.0	0.0	0.0
CNS ca. (astro) SF-539	4.4	1.8	3.8
CNS ca. (astro) SNB-75	33.0	13.8	36.1
CNS ca. (glio) SNB-19	6.0	3.4	6.7
CNS ca. (glio) U251	100.0	100.0	100.0
CNS ca. (glio) SF-295	2.0	0.5	4.2
Heart (fetal)	1.2	0.1	1.3
Heart	0.0	0.0	0.6
Fetal Skeletal	14.8	0.3	16.3
Skeletal muscle	0.1	0.2	0.0
Bone marrow	0.0	0.0	0.0
Thymus	0.3	0.0	0.5
Spleen	0.0	0.0	0.0
Lymph node	0.2	0.3	0.1
Colorectal	0.7	0.0	0.4
Stomach	0.7	1.2	0.7
Small intestine	1.2	0.6	2.6
Colon ca. SW480	2.2	0.1	2.4
Colon ca.* (SW480 met)SW620	0.0	0.0	0.0
Colon ca. HT29	0.0	0.0	0.0
Colon ca. HCT-116	0.6	0.0	0.6



Colon ca. CaCo-2	0.0	0.0	0.0
83219 CC Well to Mod Diff (ODO3866)	4.2	0.3	2.6
Colon ca. HCC-2998	0.0	0.0	0.0
Gastric ca.* (liver met) NCI-N87	2.6	1.3	1.0
Bladder	0.7	0.2	0.7
Trachea	0.4	0.2	0.1
Kidney	1.9	1.3	2.8
Kidney (fetal)	12.6	3.0	18.3
Renal ca. 786-0	10.7	2.1	15.1
Renal ca. A498	47.3	9.7	34.4
Renal ca. RXF 393	4.3	6.0	3.4
Renal ca. ACHN	2.6	0.8	2.5
Renal ca. UO-31	10.7	3.0	13.5
Renal ca. TK-10	7.4	1.6	9.3
Liver	0.0	0.0	0.3
Liver (fetal)	0.2	0.0	1.1
Liver ca. (hepatoblast) HepG2	0.0	0.0	0.0
Lung	0.1	0.0	0.3
Lung (fetal)	0.2	0.0	0.4
Lung ca. (small cell) LX-1	0.0	0.0	0.0
Lung ca. (small cell) NCI-H69	0.0	0.0	0.0
Lung ca. (s.cell var.) SHP-77	0.0	0.0	0.0
Lung ca. (large cell) NCI-H460	0.0	0.4	0.0
Lung ca. (non-sm. cell) A549	0.8	0.3	0.4
Lung ca. (non-s.cell) NCI-H23	0.0	0.0	0.0
Lung ca. (non-s.cell) HOP-62	4.7	1.7	4.1
Lung ca. (non-s.cl) NCI-H522	0.0	0.0	0.0
Lung ca. (squam.) SW 900	3.7	1.6	4.0
Lung ca. (squam.) NCI-H596	0.0	0.0	0.0
Mammary gland	4.0	1.1	5.8
Breast ca.* (pl. effusion) MCF-7	0.0	0.1	0.0
Breast ca.* (pl.ef) MDA-MB-231	46.3	7.8	12.4
Breast ca.* (pl. effusion) T47D	0.0	0.0	0.0
Breast ca. BT-549	6.7	2.7	2.6
Breast ca. MDA-N	8.3	1.2	11.1
Ovary	59.9	5.8	88.9
Ovarian ca. OVCAR-3	0.0	2.6	0.0
Ovarian ca. OVCAR-4	0.0	0.0	0.0
Ovarian ca. OVCAR-5	6.8	1.3	2.5
Ovarian ca. OVCAR-8	1.8	0.1	1.1
Ovarian ca. IGROV-1	4.3	0.9	4.2
Ovarian ca.* (ascites) SK-OV-3	1.8	1.7	0.6
Uterus	0.3	0.4	1.2
Placenta	0.0	0.0	0.2
Prostate	0.1	0.1	0.1

Prostate ca.* (bone met)PC-3	0.7	0.3	0.7
Testis	0.4	0.2	0.8
Melanoma Hs688(A).T	0.0	0.0	0.0
Melanoma* (met) Hs688(B).T	0.4	0.2	4.0
Melanoma UACC-62	0.0	0.0	0.0
Melanoma M14	0.0	0.1	0.0
Melanoma LOX IMVI	3.2	0.8	0.6
Melanoma* (met) SK-MEL-5	0.0	0.0	0.0
Adipose	1.7	0.1	1.5

**Table 18. Panel 2D**

Tissue Name	Relative Expression(%) 2dtm4181fam_ ag2197	Relative Expression(%) 2Dtm2694t_ ag708	Relative Expression(%) 2dx4tm4810t_ ag1313b_b2
Normal Colon GENPAK 061003	100.0	28.3	18.4
83219 CC Well to Mod Diff (ODO3866)	28.1	2.8	5.6
83220 CC NAT (ODO3866)	0.3	0.5	3.7
83221 CC Gr.2 rectosigmoid (ODO3868)	0.1	0.2	2.4
83222 CC NAT (ODO3868)	0.0	0.5	0.2
83235 CC Mod Diff (ODO3920)	0.0	0.0	0.2
83236 CC NAT (ODO3920)	0.8	0.8	1.1
83237 CC Gr.2 ascend colon (ODO3921)	0.5	1.6	3.8
83238 CC NAT (ODO3921)	0.2	2.0	1.7
83241 CC from Partial Hepatectomy (ODO4309)	0.2	0.7	0.9
83242 Liver NAT (ODO4309)	2.0	0.0	0.2
87472 Colon mets to lung (OD04451-01)	0.0	1.1	2.0
87473 Lung NAT (OD04451-02)	4.8	0.0	0.0
Normal Prostate Clontech A+ 6546-1	9.2	0.0	1.6
84140 Prostate Cancer (OD04410)	1.0	0.2	0.3
84141 Prostate NAT (OD04410)	2.2	0.0	0.0
87073 Prostate Cancer (OD04720-01)	2.2	0.2	0.0
87074 Prostate NAT (OD04720-02)	0.3	0.6	1.1
Normal Lung GENPAK 061010	0.7	3.3	4.7
83239 Lung Met to Muscle (ODO4286)	0.0	13.2	14.1
83240 Muscle NAT (ODO4286)	0.0	1.2	0.4
84136 Lung Malignant Cancer (OD03126)	0.5	8.2	3.7
84137 Lung NAT (OD03126)	0.0	0.7	0.8
84871 Lung Cancer (OD04404)	0.0	11.7	5.4
84872 Lung NAT (OD04404)	0.0	4.1	3.9
84875 Lung Cancer (OD04565)	0.2	4.8	8.9
84876 Lung NAT (OD04565)	0.0	0.0	0.0
85950 Lung Cancer (OD04237-01)	1.0	3.3	7.3
85970 Lung NAT (OD04237-02)	0.8	1.4	0.5

83255 Ocular Mel Met to Liver (ODO4310)	0.4	0.0	0.0
83256 Liver NAT (ODO4310)	0.4	0.0	0.0
84139 Melanoma Mets to Lung (OD04321)	0.9	48.3	43.1
84138 Lung NAT (OD04321)	0.0	0.2	0.1
Normal Kidney GENPAK 061008	0.4	<b>100.0</b>	<b>100.0</b>
83786 Kidney Ca, Nuclear grade 2 (OD04338)	0.9	9.6	18.7
83787 Kidney NAT (OD04338)	0.6	29.9	28.3
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	0.5	12.0	10.4
83789 Kidney NAT (OD04339)	0.5	29.7	33.9
83790 Kidney Ca, Clear cell type (OD04340)	0.2	3.0	3.4
83791 Kidney NAT (OD04340)	0.1	38.2	34.8
83792 Kidney Ca, Nuclear grade 3 (OD04348)	0.0	16.4	16.7
83793 Kidney NAT (OD04348)	0.0	34.9	35.4
87474 Kidney Cancer (OD04622-01)	1.1	0.0	0.4
87475 Kidney NAT (OD04622-03)	0.7	5.0	4.6
85973 Kidney Cancer (OD04450-01)	1.3	15.9	21.6
85974 Kidney NAT (OD04450-03)	0.0	37.9	38.6
Kidney Cancer Clontech 8120607	6.5	0.0	0.3
Kidney NAT Clontech 8120608	1.9	7.9	8.0
Kidney Cancer Clontech 8120613	4.2	2.2	2.3
Kidney NAT Clontech 8120614	1.3	23.0	15.3
Kidney Cancer Clontech 9010320	4.3	13.3	11.1
Kidney NAT Clontech 9010321	0.3	30.4	30.6
Normal Uterus GENPAK 061018	1.0	11.7	7.3
Uterus Cancer GENPAK 064011	0.0	4.0	4.2
Normal Thyroid Clontech A+ 6570-1	4.4	0.4	0.6
Thyroid Cancer GENPAK 064010	2.9	0.0	0.0
Thyroid Cancer INVITROGEN A302152	5.3	0.3	1.2
Thyroid NAT INVITROGEN A302153	0.3	2.2	1.9
Normal Breast GENPAK 061019	4.9	2.0	3.3
84877 Breast Cancer (OD04566)	2.1	3.1	3.3
85975 Breast Cancer (OD04590-01)	4.4	7.6	4.7
85976 Breast Cancer Mets (OD04590-03)	13.2	4.3	6.3
87070 Breast Cancer Metastasis (OD04655-05)	0.1	0.4	0.8
GENPAK Breast Cancer 064006	5.7	5.4	4.1
Breast Cancer Res. Gen. 1024	5.7	5.5	5.1
Breast Cancer Clontech 9100266	0.0	4.0	3.3
Breast NAT Clontech 9100265	0.0	3.7	3.5
Breast Cancer INVITROGEN A209073	0.0	4.7	6.1
Breast NAT INVITROGEN A2090734	0.9	4.1	4.9
Normal Liver GENPAK 061009	0.0	0.0	0.0
Liver Cancer GENPAK 064003	2.2	0.0	0.0
Liver Cancer Research Genetics RNA 1025	0.4	0.0	0.0
Liver Cancer Research Genetics RNA 1026	2.0	2.5	2.0
Paired Liver Cancer Tissue Research Genetics	1.4	0.0	0.0

RNA 6004-T			
Paired Liver Tissue Research Genetics RNA 6004-N	0.0	0.0	0.4
Paired Liver Cancer Tissue Research Genetics RNA 6005-T	1.0	3.2	2.0
Paired Liver Tissue Research Genetics RNA 6005-N	0.0	0.0	0.0
Normal Bladder GENPAK 061001	0.4	6.4	6.2
Bladder Cancer Research Genetics RNA 1023	0.2	1.2	2.2
Bladder Cancer INVITROGEN A302173	0.0	6.1	6.9
87071 Bladder Cancer (OD04718-01)	0.0	13.6	14.8
87072 Bladder Normal Adjacent (OD04718-03)	0.0	8.7	9.6
Normal Ovary Res. Gen.	0.0	77.4	60.2
Ovarian Cancer GENPAK 064008	0.0	32.8	32.1
87492 Ovary Cancer (OD04768-07)	0.3	0.8	0.8
87493 Ovary NAT (OD04768-08)	0.0	12.0	10.2
Normal Stomach GENPAK 061017	0.0	2.9	2.3
Gastric Cancer Clontech 9060358	0.5	1.1	1.1
NAT Stomach Clontech 9060359	0.4	5.9	3.5
Gastric Cancer Clontech 9060395	0.0	0.4	0.2
NAT Stomach Clontech 9060394	0.3	1.8	1.1
Gastric Cancer Clontech 9060397	0.1	9.3	5.7
NAT Stomach Clontech 9060396	0.0	0.2	0.8
Gastric Cancer GENPAK 064005	0.4	0.4	1.5

Table 19. Panel 4D

Tissue Name	Relative Expression(%) 4dtm4182fam_ ag2197	Tissue Name	Relative Expression(%) 4dtm4182fam_ ag2197
93768_Secondary Th1_anti-CD28/anti-CD3	0.0	93100_HUVEC (Endothelial) IL-1b	0.1
93769_Secondary Th2_anti-CD28/anti-CD3	0.0	93779_HUVEC (Endothelial) IFN gamma	0.0
93770_Secondary Tr1_anti-CD28/anti-CD3	0.0	93102_HUVEC (Endothelial)_TNF alpha + IFN gamma	0.0
93573_Secondary Th1_resting day 4-6 in IL-2	0.0	93101_HUVEC (Endothelial)_TNF alpha + IL4	0.0
93572_Secondary Th2_resting day 4-6 in IL-2	0.0	93781_HUVEC (Endothelial)_IL-11	0.0
93571_Secondary Tr1_resting day 4-6 in IL-2	0.0	93583_Lung Microvascular Endothelial Cells_none	0.0
93568_primary Th1_anti-CD28/anti-CD3	0.0	93584_Lung Microvascular Endothelial Cells_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93569_primary Th2_anti-CD28/anti-CD3	0.0	92662_Microvascular Dermal endothelium_none	0.0

93570_primary Tr1_anti-CD28/anti-CD3	0.0	92663_Microvascular Dermal endothelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.2
93565_primary Th1_resting dy 4-6 in IL-2	0.0	93773_Bronchial epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml) **	0.0
93566_primary Th2_resting dy 4-6 in IL-2	0.0	93347_Small Airway Epithelium_none	0.4
93567_primary Tr1_resting dy 4-6 in IL-2	0.0	93348_Small Airway Epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93351_CD45RA CD4 lymphocyte_anti-CD28/anti-CD3	0.0	92668_Coronary Artery SMC_resting	0.3
93352_CD45RO CD4 lymphocyte_anti-CD28/anti-CD3	0.0	92669_Coronary Artery SMC_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93251_CD8 Lymphocytes_anti-CD28/anti-CD3	0.0	93107_astrocytes_resting	<b>100.0</b>
93353_chronic CD8 Lymphocytes 2ry_resting dy 4-6 in IL-2	0.0	93108_astrocytes_TNFa (4 ng/ml) and IL1b (1 ng/ml)	16.5
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	0.0	92666_KU-812 (Basophil)_resting	0.0
93354_CD4_none	0.0	92667_KU-812 (Basophil)_PMA/ionoycin	0.0
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	0.0	93579_CCD1106 (Keratinocytes)_none	0.0
93103_LAK cells_resting	0.0	93580_CCD1106 (Keratinocytes)_TNFa and IFNg **	0.0
93788_LAK cells_IL-2	0.0	93791_Liver Cirrhosis	1.3
93787_LAK cells_IL-2+IL-12	0.0	93792_Lupus Kidney	11.3
93789_LAK cells_IL-2+IFN gamma	0.0	93577_NCI-H292	8.4
93790_LAK cells_IL-2+ IL-18	0.0	93358_NCI-H292_IL-4	12.5
93104_LAK cells_PMA/ionomycin and IL-18	0.8	93360_NCI-H292_IL-9	11.8
93578_NK Cells_IL-2_resting	0.0	93359_NCI-H292_IL-13	4.2
93109_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93357_NCI-H292_IFN gamma	3.7
93110_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93777_HPAEC_-	0.0
93111_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93778_HPAEC_IL-1 beta/TNA alpha	0.0
93112_Mononuclear Cells (PBMCs)_resting	0.0	93254_Normal Human Lung Fibroblast_none	0.3
93113_Mononuclear Cells (PBMCs)_PWM	0.0	93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	0.0
93114_Mononuclear Cells	0.1	93257_Normal Human Lung	0.0

(PBMCs)_PHA-L		Fibroblast_IL-4	
93249_Ramos (B cell)_none	0.0	93256_Normal Human Lung Fibroblast_IL-9	0.0
93250_Ramos (B cell)_ionomycin	0.0	93255_Normal Human Lung Fibroblast_IL-13	0.0
93349_B lymphocytes_PWM	0.0	93258_Normal Human Lung Fibroblast_IFN gamma	0.0
93350_B lymphocytes_CD40L and IL-4	0.0	93106_Dermal Fibroblasts CCD1070_resting	0.0
92665_EOL-1 (Eosinophil)_dbcAMP differentiated	0.0	93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	0.0
93248_EOL-1 (Eosinophil)_dbcAMP/PMAionomycin	0.0	93105_Dermal Fibroblasts CCD1070_IL-1 beta 1 ng/ml	0.0
93356_Dendritic Cells_none	0.0	93772_dermal fibroblast_IFN gamma	0.0
93355_Dendritic Cells_LPS 100 ng/ml	0.0	93771_dermal fibroblast_IL-4	0.0
93775_Dendritic Cells_anti-CD40	0.0	93260_IBD Colitis 2	0.1
93774_Monocytes_resting	0.0	93261_IBD Crohns	4.1
93776_Monocytes_LPS 50 ng/ml	0.0	735010_Colon_normal	1.2
93581_Macrophages_resting	0.3	735019_Lung_none	12.2
93582_Macrophages_LPS 100 ng/ml	0.0	64028-1_Thymus_none	62.8
93098_HUVEC (Endothelial)_none	0.0	64030-1_Kidney_none	2.2
93099_HUVEC (Endothelial)_starved	0.0		

- Panel 1.2 Summary Ag708** Expression of the NOV1a gene was assessed in two independent experiments using the same probe/primer set. There appears to be poor concordance between runs for some tissues but there is good concordance for others; only those results that are in agreement will be discussed here. In both experiments, highest expression of the NOV1a gene in a sample derived from a glioblastoma cell line (CTs = 24-26). Among normal tissues derived from the central nervous system, the NOV1a gene is also expressed at moderate levels in the cerebral cortex, cerebellum and hippocampus.
- Among tissues with metabolic function, the NOV1a gene is expressed in the pancreas. Thus, this gene may be involved in the pathogenesis and/or treatment of diseases involving the pancreas, such as pancreatitis and diabetes. In addition, NOV1a gene expression is decreased in a pancreatic cancer cell line.

The NOV1a gene appears to be overexpressed in fetal kidney when compared to the adult kidney. This result suggests that the NOV1a gene could be used to distinguish between adult and fetal kidney tissue and that this gene may play an important role in kidney development, growth and survival. Furthermore, NOV1a gene expression is higher in normal ovary, mammary gland and lung when compared to the cancer cell lines obtained from these tissues suggesting that this can be used as a marker to differentiate malignant and normal tissue.

**Panel 1.3D Summary** Ag708/Ag1313b/Ag2197 Three experiments with three different probe and primer sets produced results that were in very good agreement. One run, designated 1.3dx4tm5365t, appears to have lower absolute expression, but produces the same expression profile as the other two experiments. Highest expression of the NOV1A gene in all three runs is seen in a sample derived from a CNS cancer cell line (CTs=27). Certain glioblastoma and astrocytoma cell lines also express this gene as well so it may have a role in different types of brain cancer. Among normal tissues derived from the central nervous system, the NOV1a gene is expressed in the amygdala, cerebellum, hippocampus and cerebral cortex.

Among tissues with metabolic function, the NOV1a gene is expressed in the pancreas and in adipose. Interestingly, this gene is also expressed at higher levels in fetal heart and skeletal tissue when compared to the adult tissues.

The NOV1a gene appears to be expressed at high levels in a sample from a renal cancer cell line and from a breast cancer cell line. In addition, the NOV1a gene is expressed in ovarian tissue, but not significantly in cell lines derived from ovarian cancer.

**Panel 2D Summary** Ag708/Ag1313b In two runs using two different probe and primer sets, highest expression of the NOV1A gene is seen in the normal ovary, colon and kidney. Furthermore, in all nine matched kidney pairs and in the matched tissue pair derived from the ovary, the NOV1A gene is expressed more highly in normal tissue than in the adjacent cancer samples. This result suggests that expression of the NOV1A gene could be used as a diagnostic marker for the presence of kidney and ovarian cancer. In addition, therapeutic upregulation of the gene activity of NOV1A could be effective in the treatment of kidney and ovarian cancer. The NOV1A gene is also expressed at higher levels in lung cancer samples, when compared to normal adjacent tissue in six out of seven matched tissue pairs. Thus, therapeutic inhibition of the NOV1a gene, through the application of antibodies or small molecule drugs, could be effective in the treatment of lung cancer.

**Panel 4D Summary Ag2197** The NOV1a gene is expressed at a high level in astrocytes (CT 27.8) and its expression is down regulated upon treatment with TNF- $\alpha$  and IL-1b, suggesting that modulation of this protein could be beneficial in the treatment of CNS diseases-associated inflammation or neurodegeneration. The NOV1a gene is also expressed highly in the thymus (CT 28.4). More moderate expression of this gene is observed in the lung and in a muco-epidermoid cell line (H292). Thus, the protein encoded by the NOV1a gene could play an important role in the normal homeostasis of these tissues. Therapeutics designed with this protein could be important for maintaining or restoring normal function to these organs during inflammation.

**Panel 4.1D Summary Ag4164** Expression of this gene is low/undetectable (CT values >35) among the samples on this panel (data not shown).

**Panel CNS\_neurodegeneration\_v1.0 Summary Ag4164** Expression of this gene is low/undetectable (CT values >35) among the samples on this panel (data not shown).

## NOV1b

The NOV1b sequence is a variant of gene NOV1a annotated above. However, the NOV1b sequence only matches a subset of the probe and primer sets discussed above and is discussed independently below in section B. Expression of gene NOV1b was assessed using the primer-probe sets Ag4164 and Ag2197 described in Tables 20 and 21. Please note that Ag4164 contains a single mismatch in the probe relative to the NOV1b sequence. This mismatch is not predicted to alter the RTQ-PCR results. Results from RTQ-PCR runs are shown in Tables 22, 23, and 24.

**Table 20. Probe Name Ag4164**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-GCACTACAAGTGAAGCCTTAC-3'	58.1	22	822	120
Probe	FAM-5'-CTCAAGTAGAAGCCGACTTATGCAAA-3'-TAMRA	64	26	845	121
Reverse	5'-TCAAATCCTTCTGCGATACAGT-3'	58.9	22	875	122

**Table 21. Probe Name Ag2197**

Primers	Sequences	TM	Length	Start	SEQ ID
---------	-----------	----	--------	-------	--------



				Position	NO:
Forward	5'-CCAAGGAAGACCTCTTCATCTT-3'	58.8	22	1022	123
Probe	FAM-5'-TCTTGCTTACGGCATAAGCGCTCTCT-3'-TAMRA	69	26	1060	124
Reverse	5'-TTCATTCTATGGGACCTCAGA-3'	58.7	22	1086	125

Table 22. Panel 1.3D

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	1.3dtm4180fam_ag2197		1.3dtm4180fam_ag2197
Liver adenocarcinoma	2.6	Kidney (fetal)	12.6
Pancreas	2.8	Renal ca. 786-0	10.7
Pancreatic ca. CAPAN 2	0.6	Renal ca. A498	47.3
Adrenal gland	0.2	Renal ca. RXF 393	4.3
Thyroid	0.2	Renal ca. ACHN	2.6
Salivary gland	0.0	Renal ca. UO-31	10.7
Pituitary gland	0.0	Renal ca. TK-10	7.4
Brain (fetal)	0.2	Liver	0.0
Brain (whole)	3.4	Liver (fetal)	0.2
Brain (amygdala)	1.8	Liver ca. (hepatoblast) HepG2	0.0
Brain (cerebellum)	7.4	Lung	0.1
Brain (hippocampus)	7.6	Lung (fetal)	0.2
Brain (substantia nigra)	0.0	Lung ca. (small cell) LX-1	0.0
Brain (thalamus)	0.2	Lung ca. (small cell) NCI-H69	0.0
Cerebral Cortex	5.6	Lung ca. (s.cell var.) SHP-77	0.0
Spinal cord	0.2	Lung ca. (large cell) NCI-H460	0.0
CNS ca. (glio/astro) U87-MG	7.0	Lung ca. (non-sm. cell) A549	0.8
CNS ca. (glio/astro) U-118-MG	12.2	Lung ca. (non-s.cell) NCI-H23	0.0
CNS ca. (astro) SW1783	4.9	Lung ca. (non-s.cell) HOP-62	4.7
CNS ca.* (neuro; met ) SK-N-AS	0.0	Lung ca. (non-s.cl) NCI-H522	0.0
CNS ca. (astro) SF-539	4.4	Lung ca. (squamous) SW 900	3.7
CNS ca. (astro) SNB-75	33.0	Lung ca. (squamous) NCI-H596	0.0
CNS ca. (glio) SNB-19	6.0	Mammary gland	4.0
CNS ca. (glio) U251	100.0	Breast ca.* (pl. effusion) MCF-7	0.0
CNS ca. (glio) SF-295	2.0	Breast ca.* (pl.ef) MDA-MB-231	46.3
Heart (fetal)	1.2	Breast ca.* (pl. effusion) T47D	0.0
Heart	0.0	Breast ca. BT-549	6.7
Fetal Skeletal	14.8	Breast ca. MDA-N	8.3
Skeletal muscle	0.1	Ovary	59.9
Bone marrow	0.0	Ovarian ca. OVCAR-3	0.0
Thymus	0.3	Ovarian ca. OVCAR-4	0.0
Spleen	0.0	Ovarian ca. OVCAR-5	6.8

Lymph node	0.2	Ovarian ca. OVCAR-8	1.8
Colorectal	0.7	Ovarian ca. IGROV-1	4.3
Stomach	0.7	Ovarian ca.* (ascites) SK-OV-3	1.8
Small intestine	1.2	Uterus	0.3
Colon ca. SW480	2.2	Placenta	0.0
Colon ca.* (SW480 met)SW620	0.0	Prostate	0.1
Colon ca. HT29	0.0	Prostate ca.* (bone met)PC-3	0.7
Colon ca. HCT-116	0.6	Testis	0.4
Colon ca. CaCo-2	0.0	Melanoma Hs688(A).T	0.0
83219 CC Well to Mod Diff (ODO3866)	4.2	Melanoma* (met) Hs688(B).T	0.4
Colon ca. HCC-2998	0.0	Melanoma UACC-62	0.0
Gastric ca.* (liver met) NCI-N87	2.6	Melanoma M14	0.0
Bladder	0.7	Melanoma LOX IMVI	3.2
Trachea	0.4	Melanoma* (met) SK-MEL-5	0.0
Kidney	1.9	Adipose	1.7

**Table 23. Panel 2D**

<b>Tissue Name</b>	<b>Relative Expression(%) 2dtm4181fam _ag2197</b>	<b>Tissue Name</b>	<b>Relative Expression(%) 2dtm4181fam _ag2197</b>
Normal Colon GENPAK 061003	100.0	Kidney NAT Clontech 8120608	1.9
83219 CC Well to Mod Diff (ODO3866)	28.1	Kidney Cancer Clontech 8120613	4.2
83220 CC NAT (ODO3866)	0.3	Kidney NAT Clontech 8120614	1.3
83221 CC Gr.2 rectosigmoid (ODO3868)	0.1	Kidney Cancer Clontech 9010320	4.3
83222 CC NAT (ODO3868)	0.0	Kidney NAT Clontech 9010321	0.3
83235 CC Mod Diff (ODO3920)	0.0	Normal Uterus GENPAK 061018	1.0
83236 CC NAT (ODO3920)	0.8	Uterus Cancer GENPAK 064011	0.0
83237 CC Gr.2 ascend colon (ODO3921)	0.5	Normal Thyroid Clontech A+ 6570-1	4.4
83238 CC NAT (ODO3921)	0.2	Thyroid Cancer GENPAK 064010	2.9
83241 CC from Partial Hepatectomy (ODO4309)	0.2	Thyroid Cancer INVITROGEN A302152	5.3
83242 Liver NAT (ODO4309)	2.0	Thyroid NAT INVITROGEN A302153	0.3
87472 Colon mets to lung (OD04451-01)	0.0	Normal Breast GENPAK 061019	4.9
87473 Lung NAT (OD04451-02)	4.8	84877 Breast Cancer (OD04566)	2.1
Normal Prostate Clontech A+ 6546-1	9.2	85975 Breast Cancer (OD04590-01)	4.4

84140 Prostate Cancer (OD04410)	1.0	85976 Breast Cancer Mets (OD04590-03)	13.2
84141 Prostate NAT (OD04410)	2.2	87070 Breast Cancer Metastasis (OD04655-05)	0.1
87073 Prostate Cancer (OD04720-01)	2.2	GENPAK Breast Cancer 064006	5.7
87074 Prostate NAT (OD04720-02)	0.3	Breast Cancer Res. Gen. 1024	5.7
Normal Lung GENPAK 061010	0.7	Breast Cancer Clontech 9100266	0.0
83239 Lung Met to Muscle (ODO4286)	0.0	Breast NAT Clontech 9100265	0.0
83240 Muscle NAT (ODO4286)	0.0	Breast Cancer INVITROGEN A209073	0.0
84136 Lung Malignant Cancer (OD03126)	0.5	Breast NAT INVITROGEN A2090734	0.9
84137 Lung NAT (OD03126)	0.0	Normal Liver GENPAK 061009	0.0
84871 Lung Cancer (OD04404)	0.0	Liver Cancer GENPAK 064003	2.2
84872 Lung NAT (OD04404)	0.0	Liver Cancer Research Genetics RNA 1025	0.4
84875 Lung Cancer (OD04565)	0.2	Liver Cancer Research Genetics RNA 1026	2.0
84876 Lung NAT (OD04565)	0.0	Paired Liver Cancer Tissue Research Genetics RNA 6004- T	1.4
85950 Lung Cancer (OD04237- 01)	1.0	Paired Liver Tissue Research Genetics RNA 6004-N	0.0
85970 Lung NAT (OD04237- 02)	0.8	Paired Liver Cancer Tissue Research Genetics RNA 6005- T	1.0
83255 Ocular Mel Met to Liver (ODO4310)	0.4	Paired Liver Tissue Research Genetics RNA 6005-N	0.0
83256 Liver NAT (ODO4310)	0.4	Normal Bladder GENPAK 061001	0.4
84139 Melanoma Mets to Lung (OD04321)	0.9	Bladder Cancer Research Genetics RNA 1023	0.2
84138 Lung NAT (OD04321)	0.0	Bladder Cancer INVITROGEN A302173	0.0
Normal Kidney GENPAK 061008	0.4	87071 Bladder Cancer (OD04718-01)	0.0
83786 Kidney Ca, Nuclear grade 2 (OD04338)	0.9	87072 Bladder Normal Adjacent (OD04718-03)	0.0
83787 Kidney NAT (OD04338)	0.6	Normal Ovary Res. Gen.	0.0
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	0.5	Ovarian Cancer GENPAK 064008	0.0
83789 Kidney NAT (OD04339)	0.5	87492 Ovary Cancer (OD04768-07)	0.3
83790 Kidney Ca, Clear cell type (OD04340)	0.2	87493 Ovary NAT (OD04768- 08)	0.0
83791 Kidney NAT (OD04340)	0.1	Normal Stomach GENPAK 061017	0.0
83792 Kidney Ca, Nuclear	0.0	Gastric Cancer Clontech	0.5

grade 3 (OD04348)		9060358	
83793 Kidney NAT (OD04348)	0.0	NAT Stomach Clontech 9060359	0.4
87474 Kidney Cancer (OD04622-01)	1.1	Gastric Cancer Clontech 9060395	0.0
87475 Kidney NAT (OD04622- 03)	0.7	NAT Stomach Clontech 9060394	0.3
85973 Kidney Cancer (OD04450-01)	1.3	Gastric Cancer Clontech 9060397	0.1
85974 Kidney NAT (OD04450- 03)	0.0	NAT Stomach Clontech 9060396	0.0
Kidney Cancer Clontech 8120607	6.5	Gastric Cancer GENPAK 064005	0.4

**Table 24. Panel 4D**

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	4dtm4182fam_ ag2197		4dtm4182fam_ ag2197
93768_Secondary Th1_anti- CD28/anti-CD3	0.0	93100_HUVEC (Endothelial)_IL-1b	0.1
93769_Secondary Th2_anti- CD28/anti-CD3	0.0	93779_HUVEC (Endothelial)_IFN gamma	0.0
93770_Secondary Tr1_anti- CD28/anti-CD3	0.0	93102_HUVEC (Endothelial)_TNF alpha + IFN gamma	0.0
93573_Secondary Th1_resting day 4-6 in IL-2	0.0	93101_HUVEC (Endothelial)_TNF alpha + IL4	0.0
93572_Secondary Th2_resting day 4-6 in IL-2	0.0	93781_HUVEC (Endothelial)_IL-11	0.0
93571_Secondary Tr1_resting day 4-6 in IL-2	0.0	93583_Lung Microvascular Endothelial Cells_none	0.0
93568_primary Th1_anti- CD28/anti-CD3	0.0	93584_Lung Microvascular Endothelial Cells_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93569_primary Th2_anti- CD28/anti-CD3	0.0	92662_Microvascular Dermal endothelium_none	0.0
93570_primary Tr1_anti- CD28/anti-CD3	0.0	92663_Microvascular Dermal endothelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.2
93565_primary Th1_resting dy 4-6 in IL-2	0.0	93773_Bronchial epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml) **	0.0
93566_primary Th2_resting dy 4-6 in IL-2	0.0	93347_Small Airway Epithelium_none	0.4
93567_primary Tr1_resting dy 4-6 in IL-2	0.0	93348_Small Airway Epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93351_CD45RA CD4 lymphocyte_anti-CD28/anti- CD3	0.0	92668_Coronary Artery SMC_resting	0.3
93352_CD45RO CD4	0.0	92669_Coronary Artery	0.0

lymphocyte_anti-CD28/anti-CD3		SMC_TNFa (4 ng/ml) and IL1b (1 ng/ml)	
93251_CD8 Lymphocytes_anti-CD28/anti-CD3	0.0	93107_astrocytes_resting	<b>100.0</b>
93353_chronic CD8 Lymphocytes 2ry_resting dy 4-6 in IL-2	0.0	93108_astrocytes_TNFa (4 ng/ml) and IL1b (1 ng/ml)	16.5
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	0.0	92666_KU-812 (Basophil)_resting	0.0
93354_CD4_none	0.0	92667_KU-812 (Basophil)_PMA/ionomycin	0.0
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	0.0	93579_CCD1106 (Keratinocytes)_none	0.0
93103_LAK cells_resting	0.0	93580_CCD1106 (Keratinocytes)_TNFa and IFNg **	0.0
93788_LAK cells_IL-2	0.0	93791_Liver Cirrhosis	1.3
93787_LAK cells_IL-2+IL-12	0.0	93792_Lupus Kidney	11.3
93789_LAK cells_IL-2+IFN gamma	0.0	93577_NCI-H292	8.4
93790_LAK cells_IL-2+ IL-18	0.0	93358_NCI-H292_IL-4	12.5
93104_LAK cells_PMA/ionomycin and IL-18	0.8	93360_NCI-H292_IL-9	11.8
93578_NK Cells_IL-2_resting	0.0	93359_NCI-H292_IL-13	4.2
93109_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93357_NCI-H292_IFN gamma	3.7
93110_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93777_HPAEC_-	0.0
93111_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93778_HPAEC_IL-1 beta/TNA alpha	0.0
93112_Mononuclear Cells (PBMcs)_resting	0.0	93254_Normal Human Lung Fibroblast_none	0.3
93113_Mononuclear Cells (PBMcs)_PWM	0.0	93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	0.0
93114_Mononuclear Cells (PBMcs)_PHA-L	0.1	93257_Normal Human Lung Fibroblast_IL-4	0.0
93249_Ramos (B cell)_none	0.0	93256_Normal Human Lung Fibroblast_IL-9	0.0
93250_Ramos (B cell)_ionomycin	0.0	93255_Normal Human Lung Fibroblast_IL-13	0.0
93349_B lymphocytes_PWM	0.0	93258_Normal Human Lung Fibroblast_IFN gamma	0.0
93350_B lymphocytes_CD40L and IL-4	0.0	93106_Dermal Fibroblasts CCD1070_resting	0.0
92665_EOL-1 (Eosinophil)_dbcAMP differentiated	0.0	93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	0.0
93248_EOL-1 (Eosinophil)_dbcAMP/PMAionomycin	0.0	93105_Dermal Fibroblasts CCD1070_IL-1 beta 1 ng/ml	0.0

93356_Dendritic Cells_none	0.0	93772_dermal fibroblast_IFN gamma	0.0
93355_Dendritic Cells_LPS 100 ng/ml	0.0	93771_dermal fibroblast_IL-4	0.0
93775_Dendritic Cells_anti-CD40	0.0	93260_IBD Colitis 2	0.1
93774_Monocytes_resting	0.0	93261_IBD Crohns	4.1
93776_Monocytes_LPS 50 ng/ml	0.0	735010_Colon_normal	1.2
93581_Macrophages_resting	0.3	735019_Lung_none	12.2
93582_Macrophages_LPS 100 ng/ml	0.0	64028-1_Thymus_none	62.8
93098_HUVEC (Endothelial)_none	0.0	64030-1_Kidney_none	2.2
93099_HUVEC (Endothelial)_starved	0.0		

**Panel 1.3D Summary Ag2197** Highest expression of the NOV1a gene is seen in a sample derived from a CNS cancer cell line (CT=27). Certain glioblastoma and astrocytoma cell lines also express this gene as well so it may have a role in different types of brain cancer. Among normal tissues derived from the central nervous system, the NOV1a gene is expressed in the amygdala, cerebellum, hippocampus and cerebral cortex.

Among tissues with metabolic function, the NOV1a gene is expressed in the pancreas and in adipose. Interestingly, this gene is also expressed at higher levels in fetal heart and skeletal tissue when compared to the adult tissues.

The NOV1a gene appears to be expressed at high levels in a sample from a renal cancer cell line and from a breast cancer cell line. In addition, the NOV1a gene is expressed in ovarian tissue, but not significantly in cell lines derived from ovarian cancer.

**Panel 2D Summary Ag2197** Expression of the NOV1b gene is highest in normal colon tissue (CT=25). The gene is also expressed at high levels in the uterus, but not in uterine cancer. Thus, expression of NOV1b could be used as a diagnostic marker for the presence of uterine cancer. Furthermore, therapeutic upregulation of the activity of the protein product could potentially be useful in the treatment of uterine cancer. The NOV1b gene also appears to be expressed at higher levels in liver cancers (two out of two matched tissue pairs) and kidney cancers (seven out nine matched tissue pairs) when compared to normal adjacent tissue. Thus, expression of the NOV1b gene could be used to differentiate between kidney and liver cancers and other cancers or normal tissue. Furthermore, thereapeutic inhibition of the activity of the

protein encoded by the NOV1b gene, through the use of antibodies or small molecule drugs, could be effective in treating kidney and liver cancers.

**Panel 4D Summary Ag2197** The NOV1a gene is expressed at a high level in astrocytes (CT 27.8) and its expression is down regulated upon treatment with TNF-a and IL-1b, suggesting that modulation of this protein could be beneficial in the treatment of CNS diseases-associated inflammation or neurodegeneration. The NOV1a gene is also expressed highly in the thymus (CT 28.4). More moderate expression of this gene is observed in the lung and in a muco-epidermoid cell line (H292). Thus, the protein encoded by the NOV1a gene could play an important role in the normal homeostasis of these tissues. Therapeutics designed with this protein could be important for maintaining or restoring normal function to these organs during inflammation.

**Panel 4.1D Summary Ag4164** Expression of this gene is low/undetectable (CT values >35) among the samples on this panel (data not shown).

**Panel CNS\_neurodegeneration\_v1.0 Summary Ag4164** Expression of this gene is low/undetectable (CT values >35) among the samples on this panel (data not shown).

20

#### NOV1c

Please note that the NOV1c sequence is a variant of gene NOV1a annotated in section A. However, the NOV1c sequence only matches a subset of the probe and primer sets discussed above and is discussed independently below in section C. Expression of gene NOV1c was assessed using the primer-probe sets Ag2197, Ag708 and Ag1313b described in Tables 25, 26, and 27. Results from RTQ-PCR runs are shown in Tables 28, 29, 30, and 31.

**Table 25. Probe Name Ag2197**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5' - CCAAGGAAGACCTCTTCATCTT - 3'	58.8	22	1022	126
Probe	FAM-5' - TCTTGCTTACGGCATAAGCGCTCTCT-3' - TAMRA	69	26	1060	127
Reverse	5' - TTCATTCTATGGGACCTCAGA - 3'	58.7	22	1086	128

Table 26. Probe Name Ag708

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-AAAGATGGGACTCGTCATGAC-3'	59	21	232	129
Probe	TET-5'-CACGCCATCTTACTGACTGGTCTGGA-3'-TAMRA	69.5	26	253	130
Reverse	5'-GTGCAAATCCCAAAGTGCA-3'	59.5	20	306	131

Table 27. Probe Name Ag1313b

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-CAGCTGCACGATTAATGAAGAT-3'	59.4	22	264	132
Probe	TET-5'-AGGTCTTGGACTGGCCTTCACCATT-3'-TAMRA	69	25	288	133
Reverse	5'-CCAAAGTTGTGTCCAGACTCAT-3'	59.1	22	317	134

5

Table 28. Panel 1.2

Tissue Name	Relative Expression(%)	
	1.2tm888t_ ag708	1.2tm1047t_ ag708
Endothelial cells	0.0	0.0
Heart (fetal)	1.2	0.0
Pancreas	22.5	29.5
Pancreatic ca. CAPAN 2	0.0	0.0
Adrenal Gland (new lot*)	0.9	0.0
Thyroid	0.5	0.0
Salivary gland	0.4	0.0
Pituitary gland	1.3	0.0
Brain (fetal)	4.0	0.0
Brain (whole)	7.9	6.5
Brain (amygdala)	2.1	0.0
Brain (cerebellum)	16.7	23.3
Brain (hippocampus)	4.6	3.1
Brain (thalamus)	1.5	0.0
Cerebral Cortex	10.7	5.8
Spinal cord	1.1	0.0
CNS ca. (glio/astro) U87-MG	4.4	11.7
CNS ca. (glio/astro) U-118-MG	0.6	0.1
CNS ca. (astro) SW1783	1.2	0.7
CNS ca.* (neuro; met ) SK-N-AS	0.2	0.0
CNS ca. (astro) SF-539	1.6	1.1
CNS ca. (astro) SNB-75	7.7	1.7
CNS ca. (glio) SNB-19	9.5	4.3



CNS ca. (glio) U251	100.0	100.0
CNS ca. (glio) SF-295	2.4	0.0
Heart	4.9	0.0
Skeletal Muscle (new lot*)	3.1	0.0
Bone marrow	0.0	0.0
Thymus	0.2	0.0
Spleen	0.0	0.0
Lymph node	0.2	0.0
Colorectal	0.0	0.0
Stomach	1.2	0.0
Small intestine	1.4	0.0
Colon ca. SW480	1.5	0.0
Colon ca. * (SW480 met)SW620	0.0	0.0
Colon ca. HT29	0.0	0.0
Colon ca. HCT-116	0.7	0.0
Colon ca. CaCo-2	0.0	0.0
83219 CC Well to Mod Diff (ODO3866)	0.1	0.0
Colon ca. HCC-2998	0.0	0.0
Gastric ca. * (liver met) NCI-N87	4.1	0.1
Bladder	3.5	0.0
Trachea	0.3	0.0
Kidney	5.4	10.7
Kidney (fetal)	37.9	94.0
Renal ca. 786-0	4.6	5.3
Renal ca. A498	12.2	20.2
Renal ca. RXF 393	3.3	4.2
Renal ca. ACHN	2.8	1.1
Renal ca. UO-31	2.3	0.2
Renal ca. TK-10	7.6	12.7
Liver	0.9	0.0
Liver (fetal)	0.2	0.0
Liver ca. (hepatoblast) HepG2	0.0	0.0
Lung	1.1	0.0
Lung (fetal)	1.2	0.0
Lung ca. (small cell) LX-1	0.0	0.0
Lung ca. (small cell) NCI-H69	0.1	0.0
Lung ca. (s.cell var.) SHP-77	0.0	0.0
Lung ca. (large cell)NCI-H460	0.0	0.0
Lung ca. (non-sm. cell) A549	0.5	0.0
Lung ca. (non-s.cell) NCI-H23	0.0	0.0
Lung ca (non-s.cell) HOP-62	4.9	1.1
Lung ca. (non-s.cl) NCI-H522	0.0	0.0
Lung ca. (squam.) SW 900	3.7	2.0
Lung ca. (squam.) NCI-H596	0.0	0.0
Mammary gland	3.1	4.5

Breast ca.* (pl. effusion) MCF-7	0.0	0.0
Breast ca.* (pl.ef) MDA-MB-231	3.1	6.2
Breast ca.* (pl. effusion) T47D	0.8	0.0
Breast ca. BT-549	0.7	0.0
Breast ca. MDA-N	10.7	14.6
Ovary	26.4	39.5
Ovarian ca. OVCAR-3	0.4	0.0
Ovarian ca. OVCAR-4	0.9	0.0
Ovarian ca. OVCAR-5	6.0	1.7
Ovarian ca. OVCAR-8	0.7	0.0
Ovarian ca. IGROV-1	8.1	15.2
Ovarian ca.* (ascites) SK-OV-3	2.9	0.0
Uterus	0.7	0.0
Placenta	7.1	0.0
Prostate	0.3	0.0
Prostate ca.* (bone met)PC-3	1.5	0.0
Testis	3.3	0.0
Melanoma Hs688(A).T	0.0	0.0
Melanoma* (met) Hs688(B).T	0.0	0.0
Melanoma UACC-62	0.0	0.0
Melanoma M14	0.0	0.0
Melanoma LOX IMVI	1.5	0.0
Melanoma* (met) SK-MEL-5	0.0	0.0

Table 29. Panel 1.3D

Tissue Name	Relative Expression(%)	Relative Expression(%)	Relative Expression(%)
	1.3dtm4180f am_ag2197	1.3dx4tm5365 t_ag1313b_b1	1.3dtm3301t_ ag708
Liver adenocarcinoma	2.6	0.5	2.3
Pancreas	2.8	1.4	4.4
Pancreatic ca. CAPAN 2	0.6	0.2	0.8
Adrenal gland	0.2	0.0	0.0
Thyroid	0.2	0.1	0.0
Salivary gland	0.0	0.2	0.2
Pituitary gland	0.0	0.0	0.0
Brain (fetal)	0.2	0.1	0.7
Brain (whole)	3.4	2.0	2.5
Brain (amygdala)	1.8	1.1	1.7
Brain (cerebellum)	7.4	8.0	11.2
Brain (hippocampus)	7.6	1.0	7.9
Brain (substantia nigra)	0.0	0.0	0.0
Brain (thalamus)	0.2	0.1	0.0
Cerebral Cortex	5.6	0.5	4.6
Spinal cord	0.2	0.2	0.8

CNS ca. (glio/astro) U87-MG	7.0	1.8	4.6
CNS ca. (glio/astro) U-118-MG	12.2	2.3	7.3
CNS ca. (astro) SW1783	4.9	2.0	5.7
CNS ca.* (neuro; met ) SK-N-AS	0.0	0.0	0.0
CNS ca. (astro) SF-539	4.4	1.8	3.8
CNS ca. (astro) SNB-75	33.0	13.8	36.1
CNS ca. (glio) SNB-19	6.0	3.4	6.7
CNS ca. (glio) U251	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>
CNS ca. (glio) SF-295	2.0	0.5	4.2
Heart (fetal)	1.2	0.1	1.3
Heart	0.0	0.0	0.6
Fetal Skeletal	14.8	0.3	16.3
Skeletal muscle	0.1	0.2	0.0
Bone marrow	0.0	0.0	0.0
Thymus	0.3	0.0	0.5
Spleen	0.0	0.0	0.0
Lymph node	0.2	0.3	0.1
Colorectal	0.7	0.0	0.4
Stomach	0.7	1.2	0.7
Small intestine	1.2	0.6	2.6
Colon ca. SW480	2.2	0.1	2.4
Colon ca.* (SW480 met)SW620	0.0	0.0	0.0
Colon ca. HT29	0.0	0.0	0.0
Colon ca. HCT-116	0.6	0.0	0.6
Colon ca. CaCo-2	0.0	0.0	0.0
83219 CC Well to Mod Diff (ODO3866)	4.2	0.3	2.6
Colon ca. HCC-2998	0.0	0.0	0.0
Gastric ca.* (liver met) NCI-N87	2.6	1.3	1.0
Bladder	0.7	0.2	0.7
Trachea	0.4	0.2	0.1
Kidney	1.9	1.3	2.8
Kidney (fetal)	12.6	3.0	18.3
Renal ca. 786-0	10.7	2.1	15.1
Renal ca. A498	47.3	9.7	34.4
Renal ca. RXF 393	4.3	6.0	3.4
Renal ca. ACHN	2.6	0.8	2.5
Renal ca. UO-31	10.7	3.0	13.5
Renal ca. TK-10	7.4	1.6	9.3
Liver	0.0	0.0	0.3
Liver (fetal)	0.2	0.0	1.1
Liver ca. (hepatoblast) HepG2	0.0	0.0	0.0
Lung	0.1	0.0	0.3
Lung (fetal)	0.2	0.0	0.4
Lung ca. (small cell) LX-1	0.0	0.0	0.0
Lung ca. (small cell) NCI-H69	0.0	0.0	0.0

Lung ca. (s.cell var.) SHP-77	0.0	0.0	0.0
Lung ca. (large cell) NCI-H460	0.0	0.4	0.0
Lung ca. (non-sm. cell) A549	0.8	0.3	0.4
Lung ca. (non-s.cell) NCI-H23	0.0	0.0	0.0
Lung ca (non-s.cell) HOP-62	4.7	1.7	4.1
Lung ca. (non-s.cl) NCI-H522	0.0	0.0	0.0
Lung ca. (squam.) SW 900	3.7	1.6	4.0
Lung ca. (squam.) NCI-H596	0.0	0.0	0.0
Mammary gland	4.0	1.1	5.8
Breast ca.* (pl. effusion) MCF-7	0.0	0.1	0.0
Breast ca.* (pl.ef) MDA-MB-231	46.3	7.8	12.4
Breast ca.* (pl. effusion) T47D	0.0	0.0	0.0
Breast ca. BT-549	6.7	2.7	2.6
Breast ca. MDA-N	8.3	1.2	11.1
Ovary	59.9	5.8	88.9
Ovarian ca. OVCAR-3	0.0	2.6	0.0
Ovarian ca. OVCAR-4	0.0	0.0	0.0
Ovarian ca. OVCAR-5	6.8	1.3	2.5
Ovarian ca. OVCAR-8	1.8	0.1	1.1
Ovarian ca. IGROV-1	4.3	0.9	4.2
Ovarian ca.* (ascites) SK-OV-3	1.8	1.7	0.6
Uterus	0.3	0.4	1.2
Placenta	0.0	0.0	0.2
Prostate	0.1	0.1	0.1
Prostate ca.* (bone met) PC-3	0.7	0.3	0.7
Testis	0.4	0.2	0.8
Melanoma Hs688(A).T	0.0	0.0	0.0
Melanoma* (met) Hs688(B).T	0.4	0.2	4.0
Melanoma UACC-62	0.0	0.0	0.0
Melanoma M14	0.0	0.1	0.0
Melanoma LOX IMVI	3.2	0.8	0.6
Melanoma* (met) SK-MEL-5	0.0	0.0	0.0
Adipose	1.7	0.1	1.5

Table 30. Panel 2D

Tissue Name	Relative Expression(%) 2dtm4181fam_ ag2197	Relative Expression(%) 2Dtm2694t_ ag708	Relative Expression(%) 2dx4tm4810t_ ag1313b_b2
Normal Colon GENPAK 061003	100.0	28.3	18.4
83219 CC Well to Mod Diff (ODO3866)	28.1	2.8	5.6
83220 CC NAT (ODO3866)	0.3	0.5	3.7
83221 CC Gr.2 rectosigmoid (ODO3868)	0.1	0.2	2.4
83222 CC NAT (ODO3868)	0.0	0.5	0.2
83235 CC Mod Diff (ODO3920)	0.0	0.0	0.2

83236 CC NAT (ODO3920)	0.8	0.8	1.1
83237 CC Gr.2 ascend colon (ODO3921)	0.5	1.6	3.8
83238 CC NAT (ODO3921)	0.2	2.0	1.7
83241 CC from Partial Hepatectomy (ODO4309)	0.2	0.7	0.9
83242 Liver NAT (ODO4309)	2.0	0.0	0.2
87472 Colon mets to lung (OD04451-01)	0.0	1.1	2.0
87473 Lung NAT (OD04451-02)	4.8	0.0	0.0
Normal Prostate Clontech A+ 6546-1	9.2	0.0	1.6
84140 Prostate Cancer (OD04410)	1.0	0.2	0.3
84141 Prostate NAT (OD04410)	2.2	0.0	0.0
87073 Prostate Cancer (OD04720-01)	2.2	0.2	0.0
87074 Prostate NAT (OD04720-02)	0.3	0.6	1.1
Normal Lung GENPAK 061010	0.7	3.3	4.7
83239 Lung Met to Muscle (ODO4286)	0.0	13.2	14.1
83240 Muscle NAT (ODO4286)	0.0	1.2	0.4
84136 Lung Malignant Cancer (OD03126)	0.5	8.2	3.7
84137 Lung NAT (OD03126)	0.0	0.7	0.8
84871 Lung Cancer (OD04404)	0.0	11.7	5.4
84872 Lung NAT (OD04404)	0.0	4.1	3.9
84875 Lung Cancer (OD04565)	0.2	4.8	8.9
84876 Lung NAT (OD04565)	0.0	0.0	0.0
85950 Lung Cancer (OD04237-01)	1.0	3.3	7.3
85970 Lung NAT (OD04237-02)	0.8	1.4	0.5
83255 Ocular Mel Met to Liver (ODO4310)	0.4	0.0	0.0
83256 Liver NAT (ODO4310)	0.4	0.0	0.0
84139 Melanoma Mets to Lung (OD04321)	0.9	48.3	43.1
84138 Lung NAT (OD04321)	0.0	0.2	0.1
Normal Kidney GENPAK 061008	0.4	<b>100.0</b>	<b>100.0</b>
83786 Kidney Ca, Nuclear grade 2 (OD04338)	0.9	9.6	18.7
83787 Kidney NAT (OD04338)	0.6	29.9	28.3
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	0.5	12.0	10.4
83789 Kidney NAT (OD04339)	0.5	29.7	33.9
83790 Kidney Ca, Clear cell type (OD04340)	0.2	3.0	3.4
83791 Kidney NAT (OD04340)	0.1	38.2	34.8
83792 Kidney Ca, Nuclear grade 3 (OD04348)	0.0	16.4	16.7
83793 Kidney NAT (OD04348)	0.0	34.9	35.4
87474 Kidney Cancer (OD04622-01)	1.1	0.0	0.4
87475 Kidney NAT (OD04622-03)	0.7	5.0	4.6
85973 Kidney Cancer (OD04450-01)	1.3	15.9	21.6
85974 Kidney NAT (OD04450-03)	0.0	37.9	38.6
Kidney Cancer Clontech 8120607	6.5	0.0	0.3
Kidney NAT Clontech 8120608	1.9	7.9	8.0
Kidney Cancer Clontech 8120613	4.2	2.2	2.3
Kidney NAT Clontech 8120614	1.3	23.0	15.3

Kidney Cancer Clontech 9010320	4.3	13.3	11.1
Kidney NAT Clontech 9010321	0.3	30.4	30.6
Normal Uterus GENPAK 061018	1.0	11.7	7.3
Uterus Cancer GENPAK 064011	0.0	4.0	4.2
Normal Thyroid Clontech A+ 6570-1	4.4	0.4	0.6
Thyroid Cancer GENPAK 064010	2.9	0.0	0.0
Thyroid Cancer INVITROGEN A302152	5.3	0.3	1.2
Thyroid NAT INVITROGEN A302153	0.3	2.2	1.9
Normal Breast GENPAK 061019	4.9	2.0	3.3
84877 Breast Cancer (OD04566)	2.1	3.1	3.3
85975 Breast Cancer (OD04590-01)	4.4	7.6	4.7
85976 Breast Cancer Mets (OD04590-03)	13.2	4.3	6.3
87070 Breast Cancer Metastasis (OD04655-05)	0.1	0.4	0.8
GENPAK Breast Cancer 064006	5.7	5.4	4.1
Breast Cancer Res. Gen. 1024	5.7	5.5	5.1
Breast Cancer Clontech 9100266	0.0	4.0	3.3
Breast NAT Clontech 9100265	0.0	3.7	3.5
Breast Cancer INVITROGEN A209073	0.0	4.7	6.1
Breast NAT INVITROGEN A2090734	0.9	4.1	4.9
Normal Liver GENPAK 061009	0.0	0.0	0.0
Liver Cancer GENPAK 064003	2.2	0.0	0.0
Liver Cancer Research Genetics RNA 1025	0.4	0.0	0.0
Liver Cancer Research Genetics RNA 1026	2.0	2.5	2.0
Paired Liver Cancer Tissue Research Genetics RNA 6004-T	1.4	0.0	0.0
Paired Liver Tissue Research Genetics RNA 6004-N	0.0	0.0	0.4
Paired Liver Cancer Tissue Research Genetics RNA 6005-T	1.0	3.2	2.0
Paired Liver Tissue Research Genetics RNA 6005-N	0.0	0.0	0.0
Normal Bladder GENPAK 061001	0.4	6.4	6.2
Bladder Cancer Research Genetics RNA 1023	0.2	1.2	2.2
Bladder Cancer INVITROGEN A302173	0.0	6.1	6.9
87071 Bladder Cancer (OD04718-01)	0.0	13.6	14.8
87072 Bladder Normal Adjacent (OD04718-03)	0.0	8.7	9.6
Normal Ovary Res. Gen.	0.0	77.4	60.2
Ovarian Cancer GENPAK 064008	0.0	32.8	32.1
87492 Ovary Cancer (OD04768-07)	0.3	0.8	0.8
87493 Ovary NAT (OD04768-08)	0.0	12.0	10.2
Normal Stomach GENPAK 061017	0.0	2.9	2.3
Gastric Cancer Clontech 9060358	0.5	1.1	1.1
NAT Stomach Clontech 9060359	0.4	5.9	3.5
Gastric Cancer Clontech 9060395	0.0	0.4	0.2
NAT Stomach Clontech 9060394	0.3	1.8	1.1
Gastric Cancer Clontech 9060397	0.1	9.3	5.7

NAT Stomach Clontech 9060396	0.0	0.2	0.8
Gastric Cancer GENPAK 064005	0.4	0.4	1.5

**Table 31. Panel 4D**

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	4dtm4182fam-ag2197		4dtm4182fam-ag2197
93768_Secondary Th1_anti-CD28/anti-CD3	0.0	93100_HUVEC (Endothelial) IL-1b	0.1
93769_Secondary Th2_anti-CD28/anti-CD3	0.0	93779_HUVEC (Endothelial) IFN gamma	0.0
93770_Secondary Tr1_anti-CD28/anti-CD3	0.0	93102_HUVEC (Endothelial) TNF alpha + IFN gamma	0.0
93573_Secondary Th1_resting day 4-6 in IL-2	0.0	93101_HUVEC (Endothelial) TNF alpha + IL4	0.0
93572_Secondary Th2_resting day 4-6 in IL-2	0.0	93781_HUVEC (Endothelial) IL-11	0.0
93571_Secondary Tr1_resting day 4-6 in IL-2	0.0	93583_Lung Microvascular Endothelial Cells none	0.0
93568_primary Th1_anti-CD28/anti-CD3	0.0	93584_Lung Microvascular Endothelial Cells TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93569_primary Th2_anti-CD28/anti-CD3	0.0	92662_Microvascular Dermal endothelium none	0.0
93570_primary Tr1_anti-CD28/anti-CD3	0.0	92663_Microvascular Dermal endothelium TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.2
93565_primary Th1_resting dy 4-6 in IL-2	0.0	93773_Bronchial epithelium TNFa (4 ng/ml) and IL1b (1 ng/ml) **	0.0
93566_primary Th2_resting dy 4-6 in IL-2	0.0	93347_Small Airway Epithelium none	0.4
93567_primary Tr1_resting dy 4-6 in IL-2	0.0	93348_Small Airway Epithelium TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93351_CD45RA CD4 lymphocyte_anti-CD28/anti-CD3	0.0	92668_Coronary Artery SMC_resting	0.3
93352_CD45RO CD4 lymphocyte_anti-CD28/anti-CD3	0.0	92669_Coronary Artery SMC TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.0
93251_CD8 Lymphocytes_anti-CD28/anti-CD3	0.0	93107_astrocytes_resting	100.0
93353_chronic CD8 Lymphocytes 2ry_resting dy 4-6 in IL-2	0.0	93108_astrocytes TNFa (4 ng/ml) and IL1b (1 ng/ml)	16.5
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	0.0	92666_KU-812 (Basophil) resting	0.0
93354_CD4 none	0.0	92667_KU-812	0.0

		(Basophil)_PMA/ionomycin	
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	0.0	93579_CCD1106 (Keratinocytes)_none	0.0
93103_LAK cells_resting	0.0	93580_CCD1106 (Keratinocytes)_TNFa and IFNg **	0.0
93788_LAK cells_IL-2	0.0	93791_Liver Cirrhosis	1.3
93787_LAK cells_IL-2+IL-12	0.0	93792_Lupus Kidney	11.3
93789_LAK cells_IL-2+IFN gamma	0.0	93577_NCI-H292	8.4
93790_LAK cells_IL-2+ IL-18	0.0	93358_NCI-H292_IL-4	12.5
93104_LAK cells_PMA/ionomycin and IL- 18	0.8	93360_NCI-H292_IL-9	11.8
93578_NK Cells IL-2_resting	0.0	93359_NCI-H292_IL-13	4.2
93109_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93357_NCI-H292_IFN gamma	3.7
93110_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93777_HPAEC_-	0.0
93111_Mixed Lymphocyte Reaction_Two Way MLR	0.0	93778_HPAEC_IL-1 beta/TNA alpha	0.0
93112_Mononuclear Cells (PBMcs)_resting	0.0	93254_Normal Human Lung Fibroblast_none	0.3
93113_Mononuclear Cells (PBMcs)_PWM	0.0	93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	0.0
93114_Mononuclear Cells (PBMcs)_PHA-L	0.1	93257_Normal Human Lung Fibroblast_IL-4	0.0
93249_Ramos (B cell)_none	0.0	93256_Normal Human Lung Fibroblast_IL-9	0.0
93250_Ramos (B cell)_ionomycin	0.0	93255_Normal Human Lung Fibroblast_IL-13	0.0
93349_B lymphocytes_PWM	0.0	93258_Normal Human Lung Fibroblast_IFN gamma	0.0
93350_B lymphocytes_CD40L and IL-4	0.0	93106_Dermal Fibroblasts CCD1070_resting	0.0
92665_EOL-1 (Eosinophil)_dbcAMP differentiated	0.0	93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	0.0
93248_EOL-1 (Eosinophil)_dbcAMP/PMAion omycin	0.0	93105_Dermal Fibroblasts CCD1070_IL-1 beta 1 ng/ml	0.0
93356_Dendritic Cells_none	0.0	93772_dermal fibroblast_IFN gamma	0.0
93355_Dendritic Cells_LPS 100 ng/ml	0.0	93771_dermal fibroblast_IL-4	0.0
93775_Dendritic Cells_anti- CD40	0.0	93260_IBD Colitis 2	0.1
93774_Monocytes_resting	0.0	93261_IBD Crohns	4.1
93776_Monocytes_LPS 50 ng/ml	0.0	735010_Colon_normal	1.2
93581_Macrophages_resting	0.3	735019_Lung_none	12.2



93582_Macrophages_LPS 100 ng/ml	0.0	64028-1_Thymus_none	62.8
93098_HUVEC (Endothelial)_none	0.0	64030-1_Kidney_none	2.2
93099_HUVEC (Endothelial)_starved	0.0		

**Panel 1.2 Summary Ag708** Expression of the NOV1a gene was assessed in two independent experiments using the same probe/primer set. There appears to be poor concordance between runs for some tissues but there is good concordance for others; only those results that are in agreement will be discussed here. In both experiments, highest expression of the NOV1a gene in a sample derived from a glioblastoma cell line (CTs = 24-26). Among normal tissues derived from the central nervous system, the NOV1a gene is also expressed at moderate levels in the cerebral cortex, cerebellum and hippocampus.

Among tissues with metabolic function, the NOV1a gene is expressed in the pancreas. Thus, this gene may be involved in the pathogenesis and/or treatment of diseases involving the pancreas, such as pancreatitis and diabetes. In addition, NOV1a gene expression is decreased in a pancreatic cancer cell line.

The NOV1a gene appears to be overexpressed in fetal kidney when compared to the adult kidney. This result suggests that the NOV1a gene could be used to distinguish between adult and fetal kidney tissue and that this gene may play an important role in kidney development, growth and survival. Furthermore, NOV1a gene expression is higher in normal ovary, mammary gland and lung when compared to the cancer cell lines obtained from these tissues suggesting that this can be used as a marker to differentiate malignant and normal tissue.

**Panel 1.3D Summary Ag708/ Ag1313b/Ag2197** Three experiments with three different probe and primer sets produced results that were in very good agreement. One run, designated 1.3dx4tm5365t, appears to have lower absolute expression, but produces the same expression profile as the other two experiments. Highest expression of the NOV1a gene in all three runs is seen in a sample derived from a CNS cancer cell line (CTs=27). Certain glioblastoma and astrocytoma cell lines also express this gene as well so it may have a role in different types of brain cancer. Among normal tissues derived from the central nervous system, the NOV1a gene is expressed in the amygdala, cerebellum, hippocampus and cerebral cortex.

Among tissues with metabolic function, the NOV1a gene is expressed in the pancreas and in adipose. Interestingly, this gene is also expressed at higher levels in fetal heart and skeletal tissue when compared to the adult tissues.

5 The NOV1a gene appears to be expressed at high levels in a sample from a renal cancer cell line and from a breast cancer cell line. In addition, the NOV1a gene is expressed in ovarian tissue, but not significantly in cell lines derived from ovarian cancer.

**Panel 2D Summary Ag708/Ag1313b/Ag2197** In two runs using two different probe and primer sets, highest expression of the NOV1a gene is seen in the normal ovary, colon and  
10 kidney. Furthermore, in all nine matched kidney pairs and in the matched tissue pair derived from the ovary, the NOV1a gene is expressed more highly in normal tissue than in the adjacent cancer samples. This result suggests that expression of the NOV1a gene could be used as a diagnostic marker for the presence of kidney and ovarian cancer. In addition, therapeutic upregulation of the gene activity of NOV1a could be effective in the treatment of  
15 kidney and ovarian cancer. The NOV1a gene is also expressed at higher levels in lung cancer samples, when compared to normal adjacent tissue in six out of seven matched tissue pairs. Thus, therapeutic inhibition of the NOV1a gene, through the application of antibodies or small molecule drugs, could be effective in the treatment of lung cancer.

20 **Panel 4D Summary Ag2197** The NOV1a gene is expressed at a high level in astrocytes (CT 27.8) and its expression is down regulated upon treatment with TNF- $\alpha$  and IL-1 $\beta$ , suggesting that modulation of this protein could be beneficial in the treatment of CNS diseases-associated inflammation or neurodegeneration. The NOV1a gene is also expressed highly in the thymus (CT 28.4). More moderate expression of this gene is observed in the lung  
25 and in a muco-epidermoid cell line (H292). Thus, the protein encoded by the NOV1a gene could play an important role in the normal homeostasis of these tissues. Therapeutics designed with this protein could be important for maintaining or restoring normal function to these organs during inflammation.

30

### NOV3

Expression of gene NOV3 was assessed using the primer-probe set Ag1534 described in Table 32. Results from RTQ-PCR runs are shown in Tables 33, 34, 35, and 36.

Table 32. Probe Name Ag1534

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-TTTCAAGACACCCTGTGATACC-3'	59	22	765	135
Probe	FAM-5'-ACTTCGTGTCCTGAATGTTCCAGGCT-3'-TAMRA	69.1	26	799	136
Reverse	5'-CAGAGGAATGAAGGCATAGATG-3'	58.8	22	825	137

Table 33. Panel 1.2

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	1.2tm2210f_ag1534		1.2tm2210f_ag1534
Endothelial cells	2.7	Renal ca. 786-0	48.0
Heart (fetal)	2.1	Renal ca. A498	5.4
Pancreas	0.5	Renal ca. RXF 393	4.8
Pancreatic ca. CAPAN 2	5.9	Renal ca. ACHN	40.6
Adrenal Gland (new lot*)	12.6	Renal ca. UO-31	17.4
Thyroid	1.2	Renal ca. TK-10	53.2
Salivary gland	46.3	Liver	5.4
Pituitary gland	0.6	Liver (fetal)	0.9
Brain (fetal)	3.1	Liver ca. (hepatoblast) HepG2	0.3
Brain (whole)	0.2	Lung	1.2
Brain (amygdala)	4.5	Lung (fetal)	1.2
Brain (cerebellum)	0.8	Lung ca. (small cell) LX-1	2.3
Brain (hippocampus)	18.6	Lung ca. (small cell) NCI-H69	23.2
Brain (thalamus)	8.5	Lung ca. (s.cell var.) SHP-77	0.6
Cerebral Cortex	29.9	Lung ca. (large cell) NCI-H460	18.0
Spinal cord	4.8	Lung ca. (non-sm. cell) A549	13.0
CNS ca. (glio/astro) U87-MG	14.3	Lung ca. (non-s.cell) NCI-H23	42.6
CNS ca. (glio/astro) U-118-MG	6.6	Lung ca. (non-s.cell) HOP-62	24.3
CNS ca. (astro) SW1783	2.4	Lung ca. (non-s.cl) NCI-H522	18.8
CNS ca.* (neuro; met ) SK-N-AS	6.4	Lung ca. (squam.) SW 900	90.8
CNS ca. (astro) SF-539	2.6	Lung ca. (squam.) NCI-H596	40.6
CNS ca. (astro) SNB-75	6.0	Mammary gland	12.6
CNS ca. (glio) SNB-19	11.0	Breast ca.* (pl. effusion) MCF-7	17.8
CNS ca. (glio) U251	5.4	Breast ca.* (pl.ef) MDA-MB-231	1.5
CNS ca. (glio) SF-295	41.8	Breast ca.* (pl. effusion) T47D	26.6
Heart	32.1	Breast ca. BT-549	2.4
Skeletal Muscle (new lot*)	2.7	Breast ca. MDA-N	0.2
Bone marrow	1.0	Ovary	3.7
Thymus	0.0	Ovarian ca. OVCAR-3	10.3
Spleen	1.9	Ovarian ca. OVCAR-4	6.8

Lymph node	1.6	Ovarian ca. OVCAR-5	44.8
Colorectal	2.9	Ovarian ca. OVCAR-8	21.3
Stomach	1.0	Ovarian ca. IGROV-1	18.0
Small intestine	6.8	Ovarian ca.* (ascites) SK-OV-3	22.5
Colon ca. SW480	0.6	Uterus	6.2
Colon ca.* (SW480 met)SW620	2.2	Placenta	1.4
Colon ca. HT29	4.2	Prostate	50.3
Colon ca. HCT-116	6.3	Prostate ca.* (bone met)PC-3	47.0
Colon ca. CaCo-2	2.1	Testis	1.1
83219 CC Well to Mod Diff (ODO3866)	0.1	Melanoma Hs688(A).T	11.3
Colon ca. HCC-2998	6.3	Melanoma* (met) Hs688(B).T	7.8
Gastric ca.* (liver met) NCI-N87	0.0	Melanoma UACC-62	20.2
Bladder	28.5	Melanoma M14	16.7
Trachea	3.6	Melanoma LOX IMVI	1.3
Kidney	100.0	Melanoma* (met) SK-MEL-5	33.2
Kidney (fetal)	65.5		

Table 34. Panel 1.3D

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	1.3Dtm2922f_ ag1534		1.3Dtm2922f_ ag1534
Liver adenocarcinoma	6.2	Kidney (fetal)	36.9
Pancreas	0.0	Renal ca. 786-0	98.6
Pancreatic ca. CAPAN 2	14.8	Renal ca. A498	1.5
Adrenal gland	3.5	Renal ca. RXF 393	4.6
Thyroid	10.4	Renal ca. ACHN	28.3
Salivary gland	3.9	Renal ca. UO-31	13.1
Pituitary gland	18.8	Renal ca. TK-10	21.9
Brain (fetal)	35.8	Liver	1.8
Brain (whole)	13.9	Liver (fetal)	3.4
Brain (amygdala)	21.2	Liver ca. (hepatoblast) HepG2	0.0
Brain (cerebellum)	5.2	Lung	10.4
Brain (hippocampus)	100.0	Lung (fetal)	47.3
Brain (substantia nigra)	1.7	Lung ca. (small cell) LX-1	3.9
Brain (thalamus)	7.2	Lung ca. (small cell) NCI-H69	29.9
Cerebral Cortex	20.9	Lung ca. (s.cell var.) SHP-77	1.5
Spinal cord	5.4	Lung ca. (large cell)NCI-H460	9.0
CNS ca. (glio/astro) U87-MG	5.1	Lung ca. (non-sm. cell) A549	16.6
CNS ca. (glio/astro) U-118-MG	27.5	Lung ca. (non-s.cell) NCI-H23	41.8
CNS ca. (astro) SW1783	9.0	Lung ca. (non-s.cell) HOP-62	15.3
CNS ca.* (neuro; met ) SK-N-AS	39.5	Lung ca. (non-s.cl) NCI-H522	2.4
CNS ca. (astro) SF-539	14.2	Lung ca. (squam.) SW 900	65.1

CNS ca. (astro) SNB-75	35.4	Lung ca. (squam.) NCI-H596	9.0
CNS ca. (glio) SNB-19	28.7	Mammary gland	5.7
CNS ca. (glio) U251	12.5	Breast ca.* (pl. effusion) MCF-7	14.8
CNS ca. (glio) SF-295	26.8	Breast ca.* (pl.ef) MDA-MB-231	4.9
Heart (fetal)	0.0	Breast ca.* (pl. effusion) T47D	17.4
Heart	0.0	Breast ca. BT-549	4.4
Fetal Skeletal	21.2	Breast ca. MDA-N	0.0
Skeletal muscle	0.0	Ovary	8.5
Bone marrow	0.0	Ovarian ca. OVCAR-3	10.4
Thymus	0.0	Ovarian ca. OVCAR-4	0.0
Spleen	7.2	Ovarian ca. OVCAR-5	12.4
Lymph node	3.9	Ovarian ca. OVCAR-8	17.4
Colorectal	8.4	Ovarian ca. IGROV-1	10.9
Stomach	10.4	Ovarian ca.* (ascites) SK-OV-3	27.7
Small intestine	3.7	Uterus	13.1
Colon ca. SW480	2.8	Placenta	0.0
Colon ca.* (SW480 met)SW620	3.1	Prostate	21.6
Colon ca. HT29	5.5	Prostate ca.* (bone met)PC-3	17.9
Colon ca. HCT-116	1.3	Testis	36.9
Colon ca. CaCo-2	1.7	Melanoma Hs688(A).T	27.0
83219 CC Well to Mod Diff (ODO3866)	5.6	Melanoma* (met) Hs688(B).T	37.6
Colon ca. HCC-2998	0.0	Melanoma UACC-62	0.0
Gastric ca.* (liver met) NCI-N87	0.0	Melanoma M14	8.0
Bladder	8.8	Melanoma LOX IMVI	0.0
Trachea	40.3	Melanoma* (met) SK-MEL-5	48.3
Kidney	14.6	Adipose	32.3

Table 35. Panel 2D

Tissue Name	Relative Expression(%) 2Dtm2414f_ ag1534
Normal Colon GENPAK 061003	20.3
83219 CC Well to Mod Diff (ODO3866)	1.6
83220 CC NAT (ODO3866)	1.7
83221 CC Gr.2 rectosigmoid (ODO3868)	2.7
83222 CC NAT (ODO3868)	2.9
83235 CC Mod Diff (ODO3920)	16.0
83236 CC NAT (ODO3920)	0.5
83237 CC Gr.2 ascend colon (ODO3921)	11.0
83238 CC NAT (ODO3921)	1.7
83241 CC from Partial Hepatectomy (ODO4309)	4.5

83242 Liver NAT (ODO4309)	0.0
87472 Colon mets to lung (OD04451-01)	2.3
87473 Lung NAT (OD04451-02)	16.4
Normal Prostate Clontech A+ 6546-1	75.3
84140 Prostate Cancer (OD04410)	29.9
84141 Prostate NAT (OD04410)	24.1
87073 Prostate Cancer (OD04720-01)	29.9
87074 Prostate NAT (OD04720-02)	16.2
Normal Lung GENPAK 061010	25.5
83239 Lung Met to Muscle (ODO4286)	1.4
83240 Muscle NAT (ODO4286)	1.8
84136 Lung Malignant Cancer (OD03126)	14.1
84137 Lung NAT (OD03126)	43.8
84871 Lung Cancer (OD04404)	18.9
84872 Lung NAT (OD04404)	33.7
84875 Lung Cancer (OD04565)	7.2
84876 Lung NAT (OD04565)	8.4
85950 Lung Cancer (OD04237-01)	6.0
85970 Lung NAT (OD04237-02)	6.6
83255 Ocular Mel Met to Liver (ODO4310)	0.0
83256 Liver NAT (ODO4310)	3.2
84139 Melanoma Mets to Lung (OD04321)	8.0
84138 Lung NAT (OD04321)	17.4
Normal Kidney GENPAK 061008	62.0
83786 Kidney Ca, Nuclear grade 2 (OD04338)	20.3
83787 Kidney NAT (OD04338)	19.6
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	100.0
83789 Kidney NAT (OD04339)	25.7
83790 Kidney Ca, Clear cell type (OD04340)	7.9
83791 Kidney NAT (OD04340)	24.7
83792 Kidney Ca, Nuclear grade 3 (OD04348)	0.6
83793 Kidney NAT (OD04348)	24.3
87474 Kidney Cancer (OD04622-01)	4.1
87475 Kidney NAT (OD04622-03)	0.9
85973 Kidney Cancer (OD04450-01)	4.2
85974 Kidney NAT (OD04450-03)	24.0
Kidney Cancer Clontech 8120607	2.1
Kidney NAT Clontech 8120608	0.0
Kidney Cancer Clontech 8120613	0.8
Kidney NAT Clontech 8120614	90.8
Kidney Cancer Clontech 9010320	4.1
Kidney NAT Clontech 9010321	4.2
Normal Uterus GENPAK 061018	1.9
Uterus Cancer GENPAK 064011	12.7
Normal Thyroid Clontech A+ 6570-1	25.3

Thyroid Cancer GENPAK 064010	2.4
Thyroid Cancer INVITROGEN A302152	3.0
Thyroid NAT INVITROGEN A302153	20.3
Normal Breast GENPAK 061019	14.3
84877 Breast Cancer (OD04566)	15.5
85975 Breast Cancer (OD04590-01)	4.5
85976 Breast Cancer Mets (OD04590-03)	5.0
87070 Breast Cancer Metastasis (OD04655-05)	68.8
GENPAK Breast Cancer 064006	11.8
Breast Cancer Res. Gen. 1024	19.8
Breast Cancer Clontech 9100266	0.3
Breast NAT Clontech 9100265	0.0
Breast Cancer INVITROGEN A209073	11.0
Breast NAT INVITROGEN A2090734	9.9
Normal Liver GENPAK 061009	0.8
Liver Cancer GENPAK 064003	0.6
Liver Cancer Research Genetics RNA 1025	1.5
Liver Cancer Research Genetics RNA 1026	0.0
Paired Liver Cancer Tissue Research Genetics RNA 6004-T	1.5
Paired Liver Tissue Research Genetics RNA 6004-N	0.4
Paired Liver Cancer Tissue Research Genetics RNA 6005-T	0.0
Paired Liver Tissue Research Genetics RNA 6005-N	0.0
Normal Bladder GENPAK 061001	6.0
Bladder Cancer Research Genetics RNA 1023	3.4
Bladder Cancer INVITROGEN A302173	13.0
87071 Bladder Cancer (OD04718-01)	14.9
87072 Bladder Normal Adjacent (OD04718-03)	9.9
Normal Ovary Res. Gen.	4.4
Ovarian Cancer GENPAK 064008	37.1
87492 Ovary Cancer (OD04768-07)	0.9
87493 Ovary NAT (OD04768-08)	0.3
Normal Stomach GENPAK 061017	5.5
Gastric Cancer Clontech 9060358	0.0
NAT Stomach Clontech 9060359	0.0
Gastric Cancer Clontech 9060395	0.4
NAT Stomach Clontech 9060394	0.0
Gastric Cancer Clontech 9060397	0.0
NAT Stomach Clontech 9060396	0.0
Gastric Cancer GENPAK 064005	7.7

Table 36. Panel 3D

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	3dtm4931f_ ag1534		3dtm4931f_ ag1534

94905_Daoy_Medulloblastoma/ Cerebellum_sscDNA	0.6	94954_Ca Ski_Cervical epidermoid carcinoma (metastasis)_sscDNA	3.4
94906_TE671_Medulloblastom /Cerebellum_sscDNA	1.2	94955_ES-2_Ovarian clear cell carcinoma_sscDNA	2.3
94907_D283 Med_Medulloblastoma/Cerebell um_sscDNA	100.0	94957_Ramos/6h stim_ Stimulated with PMA/ionomycin 6h_sscDNA	0.0
94908_PFSK-1_Primitive Neuroectodermal/Cerebellum_s scDNA	1.6	94958_Ramos/14h stim_ Stimulated with PMA/ionomycin 14h_sscDNA	0.0
94909_XF-498_CNS_sscDNA	5.1	94962_MEG-01_Chronic myelogenous leukemia (megokaryoblast)_sscDNA	6.4
94910_SNB- 78_CNS/glioma_sscDNA	2.6	94963_Raji_Burkitt's lymphoma_sscDNA	0.0
94911_SF- 268_CNS/glioblastoma_sscDN A	1.1	94964_Daudi_Burkitt's lymphoma_sscDNA	1.2
94912_T98G_Glioblastoma_ssc DNA	4.1	94965_U266_B-cell plasmacytoma/myeloma_sscDN A	0.0
96776_SK-N- SH_Neuroblastoma (metastasis)_sscDNA	2.5	94968_CA46_Burkitt's lymphoma_sscDNA	0.0
94913_SF- 295_CNS/glioblastoma_sscDN A	2.1	94970_RL_non-Hodgkin's B- cell lymphoma_sscDNA	0.0
94914_Cerebellum_sscDNA	6.3	94972_JM1_pre-B-cell lymphoma/leukemia_sscDNA	0.7
96777_Cerebellum_sscDNA	0.0	94973_Jurkat_T cell leukemia_sscDNA	0.0
94916_NCI- H292_Mucoepidermoid lung carcinoma_sscDNA	6.7	94974_TF- 1_Erythroleukemia_sscDNA	3.5
94917_DMS-114_Small cell lung cancer_sscDNA	0.0	94975_HUT 78_T-cell lymphoma_sscDNA	0.6
94918_DMS-79_Small cell lung cancer/neuroendocrine_sscDNA	0.8	94977_U937_Histiocytic lymphoma_sscDNA	0.8
94919_NCI-H146_Small cell lung cancer/neuroendocrine_sscDNA	1.5	94980_KU-812_Myelogenous leukemia_sscDNA	2.8
94920_NCI-H526_Small cell lung cancer/neuroendocrine_sscDNA	2.5	94981_769-P_Clear cell renal carcinoma_sscDNA	9.0
94921_NCI-N417_Small cell lung cancer/neuroendocrine_sscDNA	1.7	94983_Caki-2_Clear cell renal carcinoma_sscDNA	1.1
94923_NCI-H82_Small cell lung cancer/neuroendocrine_sscDNA	18.6	94984_SW 839_Clear cell renal carcinoma_sscDNA	12.5
94924_NCI-H157_Squamous cell lung cancer	1.6	94986_G401_Wilms' tumor_sscDNA	0.0



(metastasis)_sscDNA			
94925_NCI-H1155_Large cell lung cancer/neuroendocrine_sscDNA	5.2	94987_Hs766T_Pancreatic carcinoma (LN metastasis)_sscDNA	2.0
94926_NCI-H1299_Large cell lung cancer/neuroendocrine_sscDNA	3.3	94988_CAPAN-1_Pancreatic adenocarcinoma (liver metastasis)_sscDNA	0.0
94927_NCI-H727_Lung carcinoid_sscDNA	0.0	94989_SU86.86_Pancreatic carcinoma (liver metastasis)_sscDNA	0.7
94928_NCI-UMC-11_Lung carcinoid_sscDNA	2.9	94990_BxPC-3_Pancreatic adenocarcinoma_sscDNA	3.1
94929_LX-1_Small cell lung cancer_sscDNA	0.0	94991_HPAC_Pancreatic adenocarcinoma_sscDNA	0.4
94930_Colo-205_Colon cancer_sscDNA	0.0	94992_MIA PaCa-2_Pancreatic carcinoma_sscDNA	0.7
94931_KM12_Colon cancer_sscDNA	0.7	94993_CFPAC-1_Pancreatic ductal adenocarcinoma_sscDNA	11.8
94932_KM20L2_Colon cancer_sscDNA	0.0	94994_PANC-1_Pancreatic epithelioid ductal carcinoma_sscDNA	2.3
94933_NCI-H716_Colon cancer_sscDNA	0.0	94996_T24_Bladder carcinoma (transitional cell)_sscDNA	4.7
94935_SW-48_Colon adenocarcinoma_sscDNA	0.0	94997_5637_Bladder carcinoma_sscDNA	3.1
94936_SW1116_Colon adenocarcinoma_sscDNA	0.4	94998_HT-1197_Bladder carcinoma_sscDNA	2.9
94937_LS 174T_Colon adenocarcinoma_sscDNA	8.7	94999_UM-UC-3_Bladder carcinoma (transitional cell)_sscDNA	0.0
94938_SW-948_Colon adenocarcinoma_sscDNA	0.7	95000_A204_Rhabdomyosarcoma_sscDNA	1.3
94939_SW-480_Colon adenocarcinoma_sscDNA	4.8	95001_HT-1080_Fibrosarcoma_sscDNA	4.2
94940_NCI-SNU-5_Gastric carcinoma_sscDNA	1.3	95002_MG-63_Osteosarcoma (bone)_sscDNA	1.4
94941_KATO III_Gastric carcinoma_sscDNA	0.0	95003_SK-LMS-1_Leiomyosarcoma (vulva)_sscDNA	3.2
94943_NCI-SNU-16_Gastric carcinoma_sscDNA	0.4	95004_SJRH30_Rhabdomyosarcoma (met to bone marrow)_sscDNA	1.7
94944_NCI-SNU-1_Gastric carcinoma_sscDNA	0.3	95005_A431_Epidermoid carcinoma_sscDNA	4.8
94946_RF-1_Gastric adenocarcinoma_sscDNA	9.2	95007_WM266-4_Melanoma_sscDNA	0.0
94947_RF-48_Gastric adenocarcinoma_sscDNA	5.1	95010_DU 145_Prostate carcinoma (brain metastasis)_sscDNA	0.0
96778_MKN-45_Gastric carcinoma_sscDNA	1.1	95012_MDA-MB-468_Breast adenocarcinoma_sscDNA	2.8
94949_NCI-N87_Gastric carcinoma_sscDNA	0.8	95013_SCC-4_Squamous cell carcinoma of tongue_sscDNA	0.0

94951_OVCAR-5_Ovarian carcinoma_sscDNA	0.0	95014_SCC-9_Squamous cell carcinoma of tongue_sscDNA	0.0
94952_RL95-2_Uterine carcinoma_sscDNA	6.4	95015_SCC-15_Squamous cell carcinoma of tongue_sscDNA	0.8
94953_HelaS3_Cervical adenocarcinoma_sscDNA	0.2	95017_CAL 27_Squamous cell carcinoma of tongue_sscDNA	3.7

**Panel 1.2 Summary Ag1534** The NOV3 gene encodes a protein with homology to the ileal sodium/bile cotransporter. Highest expression of this gene is detected in the kidney (CT=28.1). The NOV3 gene appears to be expressed in clusters of cell lines derived from breast cancer, ovarian cancer, lung cancer, renal cancer and melanoma. Thus expression of this gene could be used to detect the presence of any of these cancers. Furthermore, therapeutic modulation of the expression of the NOV3 gene or the activity of its protein product may be beneficial in the treatment of breast cancer, ovarian cancer, lung cancer, renal cancer and melanoma.

Among tissues involved in metabolic function, the NOV3 gene is expressed in the thyroid, adrenal gland, heart, liver, and skeletal muscle. Thus, the protein encoded by the NOV3 gene could be involved in the pathogenesis and/or treatment of diseases that involve any of these tissues. Furthermore, the NOV3 gene is expressed at higher levels in adult heart tissue (CT=29.7) than in fetal heart tissue (CT=33.6). Therefore, expression of the NOV3 gene could also be used to differentiate between adult and fetal heart tissue.

The NOV3 gene is also widely expressed in tissues originating in the central nervous system. These tissues include the fetal brain, amygdala, hippocampus, thalamus, cerebral cortex and spinal cord. This transporter gene most likely plays a role in the uptake of nutrients. Blockade of this transporter may decrease the loss of neurons due to excitotoxicity during ischemic stroke.

**Panel 1.3D Summary Ag1534** Highest expression of the NOV3 gene in Panel 1.3D is detected in the hippocampus (CT=31.2). This gene is also expressed in the amygdala and cerebral cortex. Please see Panel 1.2 summary for discussion of potential utility of this gene with respect to CNS function.

Among tissues with metabolic function, the NOV3 gene is expressed in the thyroid, pituitary gland, adipose, and fetal skeletal muscle. Interestingly, this gene is much more highly expressed in fetal skeletal muscle (CT=33.4) than in adult skeletal muscle (CT=40), suggesting that this gene could be used to distinguish the two. In addition, the increased NOV3

gene expression in fetal skeletal muscle when compared to adult suggests that the protein product may enhance muscular growth or development in the fetus and thus may also act in a regenerative capacity in the adult. Therefore, therapeutic modulation of the NOV3 gene could be useful in treatment of muscular related disease. More specifically, treatment of weak or dystrophic muscle with the protein encoded by this gene could restore muscle mass or function. This gene is also more highly expressed in fetal kidney, lung and brain when compared to the corresponding adult tissues.

**Panel 2D Summary Ag1534** The NOV3 gene is most highly expressed in a kidney cancer (CT=31.1). In general, however, this gene is more commonly expressed at higher levels in normal tissues than the adjacent tumor tissues. Specifically, this gene is expressed at higher levels in normal adjacent tissues next to thyroid cancer as well as adjacent to some kidney and lung cancers. These data indicate that NOV3 gene expression might be used to distinguish normal tissue from malignant tissue and also that therapeutic modulation of this gene product might be of use in the treatment of these types of cancer.

**Panel 3D Summary Ag1534** Highest expression of the NOV3 gene is detected in a cell line derived from a medulloblastoma (CT=30.1). Additionally there is expression in a chronic myelogenous leukemia (megakaryoblast) cell line, a gastric adenocarcinoma cell line, a clear cell renal carcinoma cell line, a pancreatic ductal adenocarcinoma cell line and a small cell lung cancer cell line. Thus, the expression of this gene could be used to distinguish some cancer cell lines from others. In addition, these data indicate that the expression of the NOV3 gene might be associated with these forms of cancer and thus, therapeutic modulation of this gene might be of use in the treatment of cancer.

#### References:

1. Oelkers P, Kirby LC, Heubi JE, Dawson PA. Primary bile acid malabsorption caused by mutations in the ileal sodium-dependent bile acid transporter gene (SLC10A2). J Clin Invest 1997 Apr 15;99(8):1880-7.

Primary bile acid malabsorption (PBAM) is an idiopathic intestinal disorder associated with congenital diarrhea, steatorrhea, interruption of the enterohepatic circulation of bile acids, and reduced plasma cholesterol levels. The molecular basis of PBAM is unknown, and several conflicting mechanisms have been postulated. In this study, we cloned the human ileal Na<sup>+</sup>/bile acid cotransporter gene (SLC10A2) and employed single-stranded conformation

polymorphism analysis to screen for PBAM-associated mutations. Four polymorphisms were identified and sequenced in a family with congenital PBAM. One allele encoded an A171S missense mutation and a mutated donor splice site for exon 3. The other allele encoded two missense mutations at conserved amino acid positions, L243P and T262M. In transfected COS cells, the L243P, T262M, and double mutant (L243P/T262M) did not affect transporter protein expression or trafficking to the plasma membrane; however, transport of taurocholate and other bile acids was abolished. In contrast, the A171S mutation had no effect on taurocholate uptake. The dysfunctional mutations were not detected in 104 unaffected control subjects, whereas the A171S was present in 28% of that population. These findings establish that SLC10A2 mutations can cause PBAM and underscore the ileal Na<sup>+</sup>/bile acid cotransporter's role in intestinal reclamation of bile acids.

#### NOV4

Expression of gene NOV4 was assessed using the primer-probe sets Ag2432 and Ag1250 described in Tables 37 and 38. Results from RTQ-PCR runs are shown in Tables 39, 40, 41, and 42.

**Table 37. Probe Name Ag2432**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-GTAGGCAAAGGGACTCACTGT-3'	58 3	21	153	138
Probe	FAM-5'-CAGAAATCAATAATCTTTGACTGCCG-3'-TAMRA	64 1	26	189	139
Reverse	5'-GCACATTACGTGGCTGAGA-3'	58 4	19	216	140

**Table 38. Probe Name Ag1250**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-CGTGGTGAACCTCTGCCTTATAT-3'	58 3	22	112	141
Probe	TET-5'-CACAGACCTCTCGTCTTTGACCGATT-3'-TAMRA	68 7	26	149	142
Reverse	5'-AGTCCCTTTGCCTACCACAAT-3'	59 9	21	191	143

**Table 39. Panel 1.2**

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
-------------	------------------------	-------------	------------------------

	1.2tm1412t_ ag1250		1.2tm1412t_ ag1250
Endothelial cells	24.8	Renal ca. 786-0	8.4
Heart (fetal)	12.7	Renal ca. A498	27.4
Pancreas	13.3	Renal ca. RXF 393	4.1
Pancreatic ca. CAPAN 2	10.4	Renal ca. ACHN	15.2
Adrenal Gland (new lot*)	23.8	Renal ca. UO-31	11.0
Thyroid	10.0	Renal ca. TK-10	27.5
Salivary gland	31.6	Liver	18.8
Pituitary gland	10.9	Liver (fetal)	13.4
Brain (fetal)	7.1	Liver ca. (hepatoblast) HepG2	25.0
Brain (whole)	12.7	Lung	5.2
Brain (amygdala)	5.7	Lung (fetal)	6.7
Brain (cerebellum)	10.1	Lung ca. (small cell) LX-1	38.4
Brain (hippocampus)	14.2	Lung ca. (small cell) NCI-H69	12.8
Brain (thalamus)	9.1	Lung ca. (s.cell var.) SHP-77	12.3
Cerebral Cortex	15.0	Lung ca. (large cell) NCI-H460	32.1
Spinal cord	7.3	Lung ca. (non-sm. cell) A549	25.9
CNS ca. (glio/astro) U87-MG	15.7	Lung ca. (non-s.cell) NCI-H23	11.3
CNS ca. (glio/astro) U-118-MG	9.7	Lung ca. (non-s.cell) HOP-62	40.3
CNS ca. (astro) SW1783	7.1	Lung ca. (non-s.cl) NCI-H522	100.0
CNS ca.* (neuro; met ) SK-N-AS	45.7	Lung ca. (squam.) SW 900	18.4
CNS ca. (astro) SF-539	5.0	Lung ca. (squam.) NCI-H596	27.9
CNS ca. (astro) SNB-75	3.7	Mammary gland	13.8
CNS ca. (glio) SNB-19	12.3	Breast ca.* (pl. effusion) MCF-7	23.3
CNS ca. (glio) U251	6.4	Breast ca.* (pl.ef) MDA-MB-231	13.6
CNS ca. (glio) SF-295	14.5	Breast ca.* (pl. effusion) T47D	21.3
Heart	39.8	Breast ca. BT-549	14.1
Skeletal Muscle (new lot*)	74.2	Breast ca. MDA-N	0.0
Bone marrow	5.0	Ovary	11.7
Thymus	3.6	Ovarian ca. OVCAR-3	29.1
Spleen	3.5	Ovarian ca. OVCAR-4	19.2
Lymph node	6.0	Ovarian ca. OVCAR-5	21.3
Colorectal	2.6	Ovarian ca. OVCAR-8	13.5
Stomach	19.3	Ovarian ca. IGROV-1	18.9
Small intestine	15.1	Ovarian ca.* (ascites) SK-OV-3	43.8
Colon ca. SW480	19.1	Uterus	6.0
Colon ca.* (SW480 met)SW620	15.0	Placenta	12.0
Colon ca. HT29	23.0	Prostate	18.2
Colon ca. HCT-116	41.2	Prostate ca.* (bone met)PC-3	75.8
Colon ca. CaCo-2	50.0	Testis	8.7
83219 CC Well to Mod Diff (ODO3866)	4.6	Melanoma Hs688(A).T	6.5
Colon ca. HCC-2998	32.8	Melanoma* (met) Hs688(B).T	4.7

Gastric ca.* (liver met) NCI-N87	23.3	Melanoma UACC-62	54.7
Bladder	14.7	Melanoma M14	19.6
Trachea	5.9	Melanoma LOX IMVI	31.0
Kidney	25.7	Melanoma* (met) SK-MEL-5	43.8
Kidney (fetal)	23.2		

**Table 40. Panel 1.3D**

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	1.3dtm4246f_ag2432		1.3dtm4246f_ag2432
Liver adenocarcinoma	0.0	Kidney (fetal)	0.0
Pancreas	0.0	Renal ca. 786-0	0.0
Pancreatic ca. CAPAN 2	0.0	Renal ca. A498	0.0
Adrenal gland	0.0	Renal ca. RXF 393	0.0
Thyroid	0.0	Renal ca. ACHN	0.0
Salivary gland	0.0	Renal ca. UO-31	0.0
Pituitary gland	0.0	Renal ca. TK-10	0.0
Brain (fetal)	0.5	Liver	0.0
Brain (whole)	0.0	Liver (fetal)	0.0
Brain (amygdala)	0.2	Liver ca. (hepatoblast) HepG2	0.0
Brain (cerebellum)	0.4	Lung	0.0
Brain (hippocampus)	100.0	Lung (fetal)	0.0
Brain (substantia nigra)	0.0	Lung ca. (small cell) LX-1	0.0
Brain (thalamus)	0.3	Lung ca. (small cell) NCI-H69	0.0
Cerebral Cortex	0.2	Lung ca. (s.cell var.) SHP-77	0.0
Spinal cord	0.0	Lung ca. (large cell) NCI-H460	0.0
CNS ca. (glio/astro) U87-MG	0.0	Lung ca. (non-sm. cell) A549	0.0
CNS ca. (glio/astro) U-118-MG	0.0	Lung ca. (non-s.cell) NCI-H23	0.0
CNS ca. (astro) SW1783	0.0	Lung ca. (non-s.cell) HOP-62	0.0
CNS ca.* (neuro; met ) SK-N-AS	0.0	Lung ca. (non-s.cl) NCI-H522	0.0
CNS ca. (astro) SF-539	0.0	Lung ca. (squamous) SW 900	0.0
CNS ca. (astro) SNB-75	0.0	Lung ca. (squamous) NCI-H596	0.0
CNS ca. (glio) SNB-19	0.0	Mammary gland	0.0
CNS ca. (glio) U251	0.0	Breast ca.* (pl. effusion) MCF-7	0.0
CNS ca. (glio) SF-295	0.0	Breast ca.* (pl.ef) MDA-MB-231	0.0
Heart (fetal)	0.0	Breast ca.* (pl. effusion) T47D	0.0
Heart	0.0	Breast ca. BT-549	0.0
Fetal Skeletal	0.0	Breast ca. MDA-N	0.0
Skeletal muscle	0.0	Ovary	0.0
Bone marrow	0.0	Ovarian ca. OVCAR-3	0.0
Thymus	0.0	Ovarian ca. OVCAR-4	0.0

Spleen	0.0	Ovarian ca. OVCAR-5	0.0
Lymph node	0.0	Ovarian ca. OVCAR-8	0.0
Colorectal	0.0	Ovarian ca. IGROV-1	0.0
Stomach	0.0	Ovarian ca.* (ascites) SK-OV-3	0.2
Small intestine	0.0	Uterus	0.0
Colon ca. SW480	0.0	Placenta	0.0
Colon ca.* (SW480 met)SW620	0.0	Prostate	0.0
Colon ca. HT29	0.0	Prostate ca.* (bone met)PC-3	0.0
Colon ca. HCT-116	0.0	Testis	0.0
Colon ca. CaCo-2	4.7	Melanoma Hs688(A).T	0.0
83219 CC Well to Mod Diff (ODO3866)	0.0	Melanoma* (met) Hs688(B).T	0.0
Colon ca. HCC-2998	0.0	Melanoma UACC-62	0.0
Gastric ca.* (liver met) NCI- N87	0.0	Melanoma M14	0.0
Bladder	0.3	Melanoma LOX IMVI	0.0
Trachea	0.0	Melanoma* (met) SK-MEL-5	0.0
Kidney	0.0	Adipose	0.0

**Table 41. Panel 2D**

<b>Tissue Name</b>	<b>Relative Expression(%) 2dtm4247f_ ag2432</b>	<b>Tissue Name</b>	<b>Relative Expression(%) 2dtm4247f_ ag2432</b>
Normal Colon GENPAK 061003	100.0	Kidney NAT Clontech 8120608	0.0
83219 CC Well to Mod Diff (ODO3866)	7.5	Kidney Cancer Clontech 8120613	0.0
83220 CC NAT (ODO3866)	7.6	Kidney NAT Clontech 8120614	0.0
83221 CC Gr.2 rectosigmoid (ODO3868)	0.0	Kidney Cancer Clontech 9010320	0.0
83222 CC NAT (ODO3868)	15.5	Kidney NAT Clontech 9010321	2.8
83235 CC Mod Diff (ODO3920)	36.1	Normal Uterus GENPAK 061018	0.0
83236 CC NAT (ODO3920)	22.8	Uterus Cancer GENPAK 064011	37.1
83237 CC Gr.2 ascend colon (ODO3921)	38.4	Normal Thyroid Clontech A+ 6570-1	4.9
83238 CC NAT (ODO3921)	0.0	Thyroid Cancer GENPAK 064010	0.0
83241 CC from Partial Hepatectomy (ODO4309)	0.0	Thyroid Cancer INVITROGEN A302152	9.4
83242 Liver NAT (ODO4309)	0.0	Thyroid NAT INVITROGEN A302153	18.9
87472 Colon mets to lung (OD04451-01)	3.0	Normal Breast GENPAK 061019	5.8
87473 Lung NAT (OD04451- 02)	6.6	84877 Breast Cancer (OD04566)	6.9
Normal Prostate Clontech A+	6.3	85975 Breast Cancer	0.0

6546-1		(OD04590-01)	
84140 Prostate Cancer (OD04410)	55.9	85976 Breast Cancer Mets (OD04590-03)	6.7
84141 Prostate NAT (OD04410)	4.9	87070 Breast Cancer Metastasis (OD04655-05)	39.0
87073 Prostate Cancer (OD04720-01)	65.1	GENPAK Breast Cancer 064006	24.0
87074 Prostate NAT (OD04720-02)	59.9	Breast Cancer Res. Gen. 1024	12.7
Normal Lung GENPAK 061010	37.4	Breast Cancer Clontech 9100266	5.3
83239 Lung Met to Muscle (ODO4286)	3.7	Breast NAT Clontech 9100265	6.1
83240 Muscle NAT (ODO4286)	28.9	Breast Cancer INVITROGEN A209073	6.4
84136 Lung Malignant Cancer (OD03126)	0.0	Breast NAT INVITROGEN A2090734	17.2
84137 Lung NAT (OD03126)	0.0	Normal Liver GENPAK 061009	9.5
84871 Lung Cancer (OD04404)	0.0	Liver Cancer GENPAK 064003	15.2
84872 Lung NAT (OD04404)	4.8	Liver Cancer Research Genetics RNA 1025	0.0
84875 Lung Cancer (OD04565)	0.0	Liver Cancer Research Genetics RNA 1026	0.0
84876 Lung NAT (OD04565)	9.7	Paired Liver Cancer Tissue Research Genetics RNA 6004- T	0.0
85950 Lung Cancer (OD04237- 01)	12.4	Paired Liver Tissue Research Genetics RNA 6004-N	0.0
85970 Lung NAT (OD04237- 02)	3.9	Paired Liver Cancer Tissue Research Genetics RNA 6005- T	2.6
83255 Ocular Mel Met to Liver (ODO4310)	4.8	Paired Liver Tissue Research Genetics RNA 6005-N	0.0
83256 Liver NAT (ODO4310)	2.5	Normal Bladder GENPAK 061001	87.7
84139 Melanoma Mets to Lung (OD04321)	0.0	Bladder Cancer Research Genetics RNA 1023	0.0
84138 Lung NAT (OD04321)	5.7	Bladder Cancer INVITROGEN A302173	14.9
Normal Kidney GENPAK 061008	23.0	87071 Bladder Cancer (OD04718-01)	6.0
83786 Kidney Ca, Nuclear grade 2 (OD04338)	4.2	87072 Bladder Normal Adjacent (OD04718-03)	34.6
83787 Kidney NAT (OD04338)	4.2	Normal Ovary Res. Gen.	0.0
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	9.1	Ovarian Cancer GENPAK 064008	7.4
83789 Kidney NAT (OD04339)	10.3	87492 Ovary Cancer (OD04768-07)	0.0
83790 Kidney Ca, Clear cell type (OD04340)	3.0	87493 Ovary NAT (OD04768- 08)	3.8
83791 Kidney NAT (OD04340)	4.6	Normal Stomach GENPAK 061017	27.5



83792 Kidney Ca, Nuclear grade 3 (OD04348)	0.0	Gastric Cancer Clontech 9060358	0.0
83793 Kidney NAT (OD04348)	6.9	NAT Stomach Clontech 9060359	3.8
87474 Kidney Cancer (OD04622-01)	0.0	Gastric Cancer Clontech 9060395	14.9
87475 Kidney NAT (OD04622-03)	0.0	NAT Stomach Clontech 9060394	27.5
85973 Kidney Cancer (OD04450-01)	5.3	Gastric Cancer Clontech 9060397	0.0
85974 Kidney NAT (OD04450-03)	7.6	NAT Stomach Clontech 9060396	0.0
Kidney Cancer Clontech 8120607	0.0	Gastric Cancer GENPAK 064005	38.2

**Table 42. Panel 4D**

Tissue Name	Relative Expression (%)	Tissue Name	Relative Expression (%)
	4Dtm2109t_ag1250		4Dtm2109t_ag1250
93768_Secondary Th1_anti-CD28/anti-CD3	77.9	93100_HUVEC (Endothelial)_IL-1b	11.6
93769_Secondary Th2_anti-CD28/anti-CD3	44.4	93779_HUVEC (Endothelial)_IFN gamma	14.6
93770_Secondary Tr1_anti-CD28/anti-CD3	68.3	93102_HUVEC (Endothelial)_TNF alpha + IFN gamma	6.7
93573_Secondary Th1_resting day 4-6 in IL-2	5.9	93101_HUVEC (Endothelial)_TNF alpha + IL4	13.4
93572_Secondary Th2_resting day 4-6 in IL-2	12.4	93781_HUVEC (Endothelial)_IL-11	3.9
93571_Secondary Tr1_resting day 4-6 in IL-2	4.4	93583_Lung Microvascular Endothelial Cells_none	17.1
93568_primary Th1_anti-CD28/anti-CD3	92.0	93584_Lung Microvascular Endothelial Cells_TNFa (4 ng/ml) and IL1b (1 ng/ml)	16.8
93569_primary Th2_anti-CD28/anti-CD3	85.3	92662_Microvascular Dermal endothelium_none	23.3
93570_primary Tr1_anti-CD28/anti-CD3	61.1	92663_Microvascular Dermal endothelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	18.9
93565_primary Th1_resting dy 4-6 in IL-2	12.3	93773_Bronchial epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml) **	24.3
93566_primary Th2_resting dy 4-6 in IL-2	7.7	93347_Small Airway Epithelium_none	10.2
93567_primary Tr1_resting dy 4-6 in IL-2	10.2	93348_Small Airway Epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	33.0
93351_CD45RA CD4 lymphocyte_anti-CD28/anti-	37.4	92668_Coronary Artery SMC_resting	6.9

CD3			
93352_CD45RO CD4 lymphocyte_anti-CD28/anti-CD3	28.5	92669_Coronary Artery SMC_TNFa (4 ng/ml) and IL1b (1 ng/ml)	7.2
93251_CD8 Lymphocytes_anti-CD28/anti-CD3	42.9	93107_astrocytes_resting	4.7
93353_chronic CD8 Lymphocytes 2ry_resting dy 4-6 in IL-2	47.0	93108_astrocytes_TNFa (4 ng/ml) and IL1b (1 ng/ml)	8.8
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	25.7	92666_KU-812 (Basophil)_resting	20.4
93354_CD4_none	3.2	92667_KU-812 (Basophil)_PMA/ionoycin	31.2
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	10.1	93579_CCD1106 (Keratinocytes)_none	19.2
93103_LAK cells_resting	14.9	93580_CCD1106 (Keratinocytes)_TNFa and IFNg **	58.6
93788_LAK cells_IL-2	20.7	93791_Liver Cirrhosis	6.3
93787_LAK cells_IL-2+IL-12	29.3	93792_Lupus Kidney	5.7
93789_LAK cells_IL-2+IFN gamma	19.5	93577_NCI-H292	19.5
93790_LAK cells_IL-2+ IL-18	14.0	93358_NCI-H292_IL-4	24.7
93104_LAK cells_PMA/ionomycin and IL-18	10.6	93360_NCI-H292_IL-9	30.1
93578_NK Cells_IL-2_resting	13.3	93359_NCI-H292_IL-13	24.8
93109_Mixed Lymphocyte Reaction_Two Way MLR	13.4	93357_NCI-H292_IFN gamma	17.3
93110_Mixed Lymphocyte Reaction_Two Way MLR	19.3	93777_HPAEC_-	9.3
93111_Mixed Lymphocyte Reaction_Two Way MLR	13.5	93778_HPAEC_IL-1 beta/TNA alpha	7.7
93112_Mononuclear Cells (PBMCs)_resting	4.9	93254_Normal Human Lung Fibroblast_none	6.6
93113_Mononuclear Cells (PBMCs)_PWM	56.6	93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	4.2
93114_Mononuclear Cells (PBMCs)_PHA-L	34.6	93257_Normal Human Lung Fibroblast_IL-4	21.0
93249_Ramos (B cell)_none	63.3	93256_Normal Human Lung Fibroblast_IL-9	13.2
93250_Ramos (B cell)_ionomycin	73.7	93255_Normal Human Lung Fibroblast_IL-13	29.3
93349_B lymphocytes_PWM	57.4	93258_Normal Human Lung Fibroblast_IFN gamma	25.2
93350_B lymphocytes_CD40L and IL-4	17.2	93106_Dermal Fibroblasts CCD1070_resting	28.3
92665_EOI-1 (Eosinophil)_dbcAMP differentiated	10.2	93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	25.5
93248_EOI-1	5.2	93105_Dermal Fibroblasts	24.8

(Eosinophil)_dbcAMP/PMAionomycin		CCD1070_IL-1 beta 1 ng/ml	
93356_Dendritic Cells_none	7.5	93772_dermal fibroblast_IFN gamma	10.7
93355_Dendritic Cells_LPS 100 ng/ml	6.1	93771_dermal fibroblast_IL-4	12.7
93775_Dendritic Cells_anti-CD40	9.7	93260_IBD Colitis 2	0.3
93774_Monocytes_resting	4.8	93261_IBD Crohns	0.9
93776_Monocytes_LPS 50 ng/ml	3.5	735010_Colon_normal	8.7
93581_Macrophages_resting	23.2	735019_Lung_none	15.8
93582_Macrophages_LPS 100 ng/ml	13.1	64028-1_Thymus_none	<b>100.0</b>
93098_HUVEC (Endothelial)_none	19.5	64030-1_Kidney_none	11.5
93099_HUVEC (Endothelial)_starved	17.1		

**Panel 1.2 Summary Ag1250** The NOV4 gene is expressed at high levels in lung cancer cell lines. Overall, there is a predominant expression pattern that shows higher expression of this gene in cancer cell lines when compared to normal tissues. Specifically, the NOV4 gene is expressed at higher levels in samples derived from colon cancer, ovarian cancer, breast cancer, prostate cancer and melanoma cell lines. Thus, expression of the NOV4 gene could be used to distinguish cultured cell lines from normal tissues. In addition, these data indicate that the expression of this gene might be associated with these forms of cancer and thus, therapeutic modulation of the NOV4 gene product might be of use in the treatment of these cancers.

**Panel 1.3D Summary Ag2432** Expression of the NOV4 gene is limited to the hippocampus (CT=27.6), where it is expressed at high levels. Therefore, expression of this gene could be used to distinguish hippocampus from other tissues.

**Panel 2D Summary Ag2432** Significant but low expression of the NOV4 gene is detected in normal colon and bladder tissues. Therefore, expression of this gene could be used to distinguish colon and bladder from other tissues.

**Panel 4D Summary Ag1250** The NOV4 gene encodes a protein with homology to prohibitins, which are proteins that have been shown to be involved various functions, including cell cycle regulation, apoptosis, assembly of mitochondrial respiratory chain

enzymes, and aging (ref. 1). The NOV4 gene is expressed at moderate levels throughout the samples on this panel (CTs=30.8–33.6). Interestingly, however, this gene is expressed at highest levels in the thymus as well as in activated Th1 and Th2 T cells. Given this expression pattern, the NOV4 gene product may play an important role in the normal homeostasis of the thymus and might be associated with the activation process of T cells. Therefore, modulation of this protein by small molecule drugs might be important for controlling T cell activation and could have some benefit for treatment of diseases associated with hyperactive T cells, such as autoimmune disease, delayed type hypersensitivity, and other T cell mediated diseases (such as asthma and psoriasis). In addition, the NOV4 gene is also expressed in activated B cells and in a Ramos B cell line. It has been suggested that increased prohibitin expression is associated with and may facilitate B-cell maturation (ref. 2). Thus, modulation of this protein by small molecule drugs might be important for controlling B differentiation and the generation of immunoglobulins by B cells and could therefore have some therapeutic benefit in the treatment of hypoglobulinemia.

**Panel CNS\_neurodegeneration\_v1.0 Summary Ag2432** Expression of this gene is low/undetectable (CT values >35) among the samples on this panel (data not shown).

#### References:

1. Coates PJ, Nenutil R, McGregor A, Picksley SM, Crouch DH, Hall PA, Wright EG. Mammalian prohibitin proteins respond to mitochondrial stress and decrease during cellular senescence. *Exp Cell Res* 2001 May 1;265(2):262-73

The two prohibitin proteins, Phb1p and Phb2p(BAP37), have been ascribed various functions, including cell cycle regulation, apoptosis, assembly of mitochondrial respiratory chain enzymes, and aging. We show that the mammalian prohibitins are present in the inner mitochondrial membrane and are always bound to each other, with no free protein detectable. They are coexpressed during development and in adult mammalian tissues, and expression levels are indicative of a role in mitochondrial metabolism, but are not compatible with roles in the regulation of cellular proliferation or apoptosis. High level expression of the proteins is consistently seen in primary human tumors, while cellular senescence of human and chick fibroblasts is accompanied by heterogeneous decreases in both proteins. The two proteins are induced by metabolic stress caused by an imbalance in the synthesis of mitochondrial- and nuclear-encoded mitochondrial proteins, but do not respond to oxidative stress, heat shock, or

other cellular stresses. The gene promoter sequences contain binding sites for the Myc oncoprotein and overexpression of Myc induces expression of the prohibitins. The data support conserved roles for the prohibitins in regulating mitochondrial respiratory activity and in aging.

5

PMID: 11302691

2. Woodlock TJ, Bethlenny G, Segel GB. Prohibitin expression is increased in phorbol ester-treated chronic leukemic B-lymphocytes. *Blood Cells Mol Dis* 2001 Jan-Feb;27(1):27-34

10

Chronic lymphocytic leukemia (CLL) is characterized by the gradual accumulation of immature B-lymphocytes. CLL B-lymphocytes mature to a plasmacytoid phenotype when treated in vitro with phorbol esters. CLL B-cell apparent maturation is associated with altered expression of specific plasma membrane and mitochondrial proteins including heightened expression of a 30-kDa heat shock protein 60 (hsp60) analog. During our efforts to further characterize this hsp60 analog by mass spectrometry, we detected the mitochondrial protein prohibitin in phorbol-ester-matured CLL B-lymphocytes. Prohibitin modulates cell proliferation and inhibits cell cycle traverse in several systems, although few data are available for lymphocytes. A twofold increase in prohibitin concentration was observed in phorbol-ester-matured compared to resting CLL B-cells as determined by quantitative Western immunoblot analysis. A similar increase in prohibitin was observed in phorbol-ester-treated normal human B-lymphocyte populations. An antisense oligonucleotide complementary to the 5' coding region of the prohibitin gene blunted the increase in prohibitin protein in phorbol-ester-treated CLL B-cells by 42%. These data suggest that increased prohibitin expression is associated with and may facilitate B-cell maturation.

15

20

25

PMID: 11162143

30 **NOV5**

Expression of gene NOV5 was assessed using the primer-probe sets Ag3086 and Ag3797 described in Tables 43 and 44. Results from RTQ-PCR runs are shown in Tables 45, 46, 47, 48, 49, and 50.

**Table 45. Probe Name Ag3086**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-GGACCCCATTCGACTACTGT-3'	20	20	1309	144
Probe	FAM-5'-CTGATGACCAGCGCCATCAATC-3'-TAMRA	23	23	1345	145
Reverse	5'-TTCTCAAACGACCTGGTC-3'	20	20	1399	146

**Table 46. Probe Name Ag3797**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-TCTGGACGACAACTATTGCC-3'	58.7	20	627	147
Probe	FAM-5'-ATGGTGCTACACTACGGATCCGCAG-3'-TAMRA	69.2	25	672	148
Reverse	5'-GTCACAGAATTCTCGCTCGA-3'	59.1	20	698	149

5

**Table 47. Panel 1.3D**

Tissue Name	Relative Expression(%) 1.3dx4tm5430 f_ag3086_a1	Tissue Name	Relative Expression(%) 1.3dx4tm5430 f_ag3086_a1
Liver adenocarcinoma	0.7	Kidney (fetal)	31.1
Pancreas	17.9	Renal ca. 786-0	0.2
Pancreatic ca. CAPAN 2	0.6	Renal ca. A498	0.5
Adrenal gland	2.7	Renal ca. RXF 393	0.7
Thyroid	3.3	Renal ca. ACHN	0.8
Salivary gland	1.2	Renal ca. UO-31	0.4
Pituitary gland	3.6	Renal ca. TK-10	0.2
Brain (fetal)	3.2	Liver	94.2
Brain (whole)	3.4	Liver (fetal)	100.0
Brain (amygdala)	2.1	Liver ca. (hepatoblast) HepG2	58.4
Brain (cerebellum)	1.5	Lung	2.8
Brain (hippocampus)	3.0	Lung (fetal)	12.9
Brain (substantia nigra)	1.7	Lung ca. (small cell) LX-1	1.3
Brain (thalamus)	3.0	Lung ca. (small cell) NCI-H69	0.2
Cerebral Cortex	0.9	Lung ca. (s.cell var.) SHP-77	1.2
Spinal cord	2.9	Lung ca. (large cell) NCI-H460	1.4
CNS ca. (glio/astro) U87-MG	0.7	Lung ca. (non-sm. cell) A549	0.2
CNS ca. (glio/astro) U-118-MG	0.9	Lung ca. (non-s.cell) NCI-H23	0.9
CNS ca. (astro) SW1783	0.4	Lung ca. (non-s.cell) HOP-62	0.5
CNS ca.* (neuro; met) SK-N-AS	0.7	Lung ca. (non-s.cl) NCI-H522	0.6
CNS ca. (astro) SF-539	0.5	Lung ca. (squam.) SW 900	0.4
CNS ca. (astro) SNB-75	1.2	Lung ca. (squam.) NCI-H596	0.5

CNS ca. (glio) SNB-19	1.6	Mammary gland	3.1
CNS ca. (glio) U251	2.4	Breast ca.* (pl. effusion) MCF-7	0.7
CNS ca. (glio) SF-295	0.7	Breast ca.* (pl.ef) MDA-MB-231	0.7
Heart (fetal)	0.6	Breast ca.* (pl. effusion) T47D	2.7
Heart	0.5	Breast ca. BT-549	0.7
Fetal Skeletal	0.2	Breast ca. MDA-N	0.0
Skeletal muscle	1.4	Ovary	0.4
Bone marrow	2.0	Ovarian ca. OVCAR-3	0.6
Thymus	1.2	Ovarian ca. OVCAR-4	0.4
Spleen	4.0	Ovarian ca. OVCAR-5	0.4
Lymph node	3.1	Ovarian ca. OVCAR-8	0.7
Colorectal	1.6	Ovarian ca. IGROV-1	0.7
Stomach	10.3	Ovarian ca.* (ascites) SK-OV-3	0.0
Small intestine	29.6	Uterus	3.2
Colon ca. SW480	0.7	Placenta	4.6
Colon ca.* (SW480 met)SW620	0.2	Prostate	2.1
Colon ca. HT29	0.2	Prostate ca.* (bone met)PC-3	0.9
Colon ca. HCT-116	1.2	Testis	12.4
Colon ca. CaCo-2	2.4	Melanoma Hs688(A).T	0.1
83219 CC Well to Mod Diff (ODO3866)	2.1	Melanoma* (met) Hs688(B).T	0.2
Colon ca. HCC-2998	1.4	Melanoma UACC-62	0.4
Gastric ca.* (liver met) NCI-N87	1.8	Melanoma M14	0.8
Bladder	4.5	Melanoma LOX IMVI	0.0
Trachea	2.6	Melanoma* (met) SK-MEL-5	0.4
Kidney	26.1	Adipose	2.2

Table 48. General Screening Panel \_v1.4

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	1.4x4tm7355f_ag3797_a1		1.4x4tm7355f_ag3797_a1
D6005-01_Human adipose	1.4	Renal ca. TK-10	28.9
112193_Metastatic melanoma	0.4	Bladder	8.4
112192_Metastatic melanoma	0.5	Gastric ca.(liver met)_NCI-N87	2.7
95280_Epidermis (metastatic melanoma)	0.3	112197_Stomach	1.4
95279_Epidermis (metastatic melanoma)	0.3	94938_Colon Adenocarcinoma	1.0
Melanoma (met)_SK-MEL-5	0.5	Colon ca._SW480	3.9
112196_Tongue (oncology)	0.8	Colon ca.(SW480 met)_SW620	1.2
113461_Testis Pool	2.0	Colon ca._HT29	0.2
Prostate ca.(bone met)_PC-3	1.5	Colon ca._HCT-116	4.3
113455_Prostate Pool	1.8	Colon ca._CaCo-2	11.5

103396 Placenta	1.7	83219_CC Well to Mod Diff (ODO3866)	2.8
113463 Uterus Pool	0.5	94936 Colon Adenocarcinoma	2.9
Ovarian carcinoma_OVCAR-3	1.0	94930 Colon	0.5
Ovarian carcinoma(ascites)_SK-OV-3	0.8	94935 Colon Adenocarcinoma	0.2
95297_Adenocarcinoma (ovary)	0.3	113468 Colon Pool	1.6
Ovarian carcinoma_OVCAR-5	6.4	113457 Small Intestine Pool	2.0
Ovarian carcinoma_IGROV-1	4.6	113460 Stomach Pool	1.9
Ovarian carcinoma_OVCAR-8	2.9	113467 Bone Marrow Pool	0.4
103368 Ovary	1.9	103371 Fetal Heart	0.8
MCF7_breast carcinoma(pleural effusion)	2.3	113451 Heart Pool	0.7
Breast ca. (pleural effusion)_MDA-MB-231	2.2	113466 Lymph Node Pool	1.7
112189_ductal cell carcinoma(breast)	3.0	103372 Fetal Skeletal Muscle	0.7
Breast ca. (pleural effusion)_T47D	18.5	113456 Skeletal Muscle Pool	1.1
Breast carcinoma_MDA-N	0.7	113459 Spleen Pool	2.5
113452_Breast Pool	1.5	113462 Thymus Pool	2.4
103398 Trachea	1.2	CNS ca. (glio/astro)_U87-MG	2.7
112354_lung	0.4	CNS ca. (glio/astro)_U-118-MG	3.0
103374 Fetal Lung	2.3	CNS ca. (neuro;met)_SK-N-AS	2.1
94921_Small cell carcinoma of the lung	0.2	95264 Brain astrocytoma	0.6
Lung ca.(small cell)_LX-1	3.3	CNS ca. (astro)_SNB-75	1.8
94919_Small cell carcinoma of the lung	0.5	CNS ca. (glio)_SNB-19	4.1
Lung ca.(s.cell var.)_SHP-77	2.4	CNS ca. (glio)_SF-295	2.1
95268_Lung (Large cell carcinoma)	0.6	113447 Brain (Amygdala) Pool	0.9
94920_Small cell carcinoma of the lung	0.6	103382 Brain (cerebellum)	1.9
Lung ca.(non-s.cell)_NCI-H23	3.7	64019-1_brain(fetal)	2.8
Lung ca.(large cell)_NCI-H460	0.9	113448 Brain (Hippocampus) Pool	1.0
Lung ca.(non-s.cell)_HOP-62	1.2	113464 Cerebral Cortex Pool	0.7
Lung ca.(non-s.cl)_NCI-H522	1.7	113449 Brain (Substantia nigra) Pool	0.9
103392 Liver	26.6	113450 Brain (Thalamus) Pool	1.0
103393 Fetal Liver	45.5	103384 Brain (whole)	1.6
Liver ca.(hepatoblast)_HepG2	100.0	113458 Spinal Cord Pool	1.7
113465 Kidney Pool	1.7	103375 Adrenal Gland	3.1
103373 Fetal Kidney	11.1	113454 Pituitary gland Pool	1.7
Renal ca. 786-0	1.0	103397 Salivary Gland	1.0
112188_renal cell carcinoma	0.3	103369 Thyroid (female)	2.8



Renal ca. ACHN	1.4	Pancreatic ca. CAPAN2	0.8
112190 Renal cell carcinoma	1.8	113453 Pancreas Pool	7.5

**Table 49. Panel 2.2**

<b>Tissue Name</b>	<b>Relative Expression(%) 2.2x4tm6408f_ ag3086_a1</b>	<b>Tissue Name</b>	<b>Relative Expression(%) 2.2x4tm6408f_ ag3086_a1</b>
Normal Colon GENPAK 061003	1.4	83793 Kidney NAT (OD04348)	40.9
97759 Colon cancer (OD06064)	0.1	98938 Kidney malignant cancer (OD06204B)	0.4
97760 Colon cancer NAT (OD06064)	0.0	98939 Kidney normal adjacent tissue (OD06204E)	5.1
97778 Colon cancer (OD06159)	0.0	85973 Kidney Cancer (OD04450-01)	5.7
97779 Colon cancer NAT (OD06159)	1.4	85974 Kidney NAT (OD04450-03)	15.0
98861 Colon cancer (OD06297-04)	0.0	Kidney Cancer Clontech 8120613	0.2
98862 Colon cancer NAT (OD06297-015)	1.1	Kidney NAT Clontech 8120614	5.9
83237 CC Gr.2 ascend colon (ODO3921)	0.4	Kidney Cancer Clontech 9010320	0.4
83238 CC NAT (ODO3921)	0.2	Kidney NAT Clontech 9010321	1.5
97766 Colon cancer metastasis (OD06104)	0.0	Kidney Cancer Clontech 8120607	0.1
97767 Lung NAT (OD06104)	0.6	Kidney NAT Clontech 8120608	3.5
87472 Colon mets to lung (OD04451-01)	1.3	Normal Uterus GENPAK 061018	0.2
87473 Lung NAT (OD04451-02)	0.4	Uterus Cancer GENPAK 064011	0.1
Normal Prostate Clontech A+ 6546-1 (8090438)	0.6	Normal Thyroid Clontech A+ 6570-1 (7080817)	0.5
84140 Prostate Cancer (OD04410)	0.2	Thyroid Cancer GENPAK 064010	0.3
84141 Prostate NAT (OD04410)	0.5	Thyroid Cancer INVITROGEN A302152	2.1
Normal Ovary Res. Gen.	0.4	Thyroid NAT INVITROGEN A302153	0.4
98863 Ovarian cancer (OD06283-03)	0.2	Normal Breast GENPAK 061019	0.7
98865 Ovarian cancer NAT/fallopian tube (OD06283-07)	1.2	84877 Breast Cancer (OD04566)	2.3
Ovarian Cancer GENPAK 064008	1.5	Breast Cancer Res. Gen. 1024	1.9
97773 Ovarian cancer (OD06145)	0.9	85975 Breast Cancer (OD04590-01)	5.1
97775 Ovarian cancer NAT (OD06145)	1.8	85976 Breast Cancer Mets (OD04590-03)	1.3

98853 Ovarian cancer (OD06455-03)	0.6	87070 Breast Cancer Metastasis (OD04655-05)	0.7
98854 Ovarian NAT (OD06455-07) Fallopian tube	0.2	GENPAK Breast Cancer 064006	0.5
Normal Lung GENPAK 061010	0.8	Breast Cancer Clontech 9100266	0.2
92337 Invasive poor diff. lung adeno (ODO4945-01)	0.3	Breast NAT Clontech 9100265	0.5
92338 Lung NAT (ODO4945-03)	1.1	Breast Cancer INVITROGEN A209073	0.2
84136 Lung Malignant Cancer (OD03126)	1.6	Breast NAT INVITROGEN A2090734	1.4
84137 Lung NAT (OD03126)	0.3	97763 Breast cancer (OD06083)	0.7
90372 Lung Cancer (OD05014A)	0.5	97764 Breast cancer node metastasis (OD06083)	0.2
90373 Lung NAT (OD05014B)	0.6	Normal Liver GENPAK 061009	28.7
97761 Lung cancer (OD06081)	0.5	Liver Cancer Research Genetics RNA 1026	7.5
97762 Lung cancer NAT (OD06081)	1.2	Liver Cancer Research Genetics RNA 1025	45.0
85950 Lung Cancer (OD04237-01)	0.3	Paired Liver Cancer Tissue Research Genetics RNA 6004-T	35.7
85970 Lung NAT (OD04237-02)	1.1	Paired Liver Tissue Research Genetics RNA 6004-N	5.1
83255 Ocular Mel Met to Liver (ODO4310)	0.2	Paired Liver Cancer Tissue Research Genetics RNA 6005-T	14.7
83256 Liver NAT (ODO4310)	21.6	Paired Liver Tissue Research Genetics RNA 6005-N	65.0
84139 Melanoma Mets to Lung (OD04321)	0.2	Liver Cancer GENPAK 064003	<b>100.0</b>
84138 Lung NAT (OD04321)	0.2	Normal Bladder GENPAK 061001	2.8
Normal Kidney GENPAK 061008	5.5	Bladder Cancer Research Genetics RNA 1023	0.2
83786 Kidney Ca, Nuclear grade 2 (OD04338)	20.6	Bladder Cancer INVITROGEN A302173	0.7
83787 Kidney NAT (OD04338)	4.5	Normal Stomach GENPAK 061017	2.5
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	6.5	Gastric Cancer Clontech 9060397	0.0
83789 Kidney NAT (OD04339)	6.0	NAT Stomach Clontech 9060396	0.1
83790 Kidney Ca, Clear cell type (OD04340)	0.4	Gastric Cancer Clontech 9060395	0.0
83791 Kidney NAT (OD04340)	8.7	NAT Stomach Clontech 9060394	0.3
83792 Kidney Ca, Nuclear grade 3 (OD04348)	0.3	Gastric Cancer GENPAK 064005	2.8

Table 50. Panel 4D

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	4dx4tm5510f_ag3086_a2		4dx4tm5510f_ag3086_a2
93768_Secondary Th1_anti-CD28/anti-CD3	0.7	93100_HUVEC (Endothelial) IL-1b	0.4
93769_Secondary Th2_anti-CD28/anti-CD3	0.9	93779_HUVEC (Endothelial) IFN gamma	1.2
93770_Secondary Tr1_anti-CD28/anti-CD3	1.1	93102_HUVEC (Endothelial) TNF alpha + IFN gamma	0.3
93573_Secondary Th1_resting day 4-6 in IL-2	2.8	93101_HUVEC (Endothelial) TNF alpha + IL4	0.2
93572_Secondary Th2_resting day 4-6 in IL-2	1.4	93781_HUVEC (Endothelial) IL-11	0.8
93571_Secondary Tr1_resting day 4-6 in IL-2	1.3	93583_Lung Microvascular Endothelial Cells_none	1.0
93568_primary Th1_anti-CD28/anti-CD3	0.7	93584_Lung Microvascular Endothelial Cells_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.9
93569_primary Th2_anti-CD28/anti-CD3	0.9	92662_Microvascular Dermal endothelium_none	1.6
93570_primary Tr1_anti-CD28/anti-CD3	1.1	92663_Microvascular Dermal endothelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.8
93565_primary Th1_resting dy 4-6 in IL-2	6.4	93773_Bronchial epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml) **	3.2
93566_primary Th2_resting dy 4-6 in IL-2	3.1	93347_Small Airway Epithelium_none	1.8
93567_primary Tr1_resting dy 4-6 in IL-2	2.0	93348_Small Airway Epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	3.3
93351_CD45RA CD4 lymphocyte_anti-CD28/anti-CD3	0.4	92668_Coronary Artery SMC_resting	1.0
93352_CD45RO CD4 lymphocyte_anti-CD28/anti-CD3	1.4	92669_Coronary Artery SMC_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.6
93251_CD8 Lymphocytes_anti-CD28/anti-CD3	1.4	93107_astrocytes_resting	3.2
93353_chronic CD8 Lymphocytes 2ry_resting dy 4-6 in IL-2	1.7	93108_astrocytes_TNFa (4 ng/ml) and IL1b (1 ng/ml)	2.7
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	0.6	92666_KU-812 (Basophil)_resting	1.5
93354_CD4_none	2.1	92667_KU-812 (Basophil)_PMA/ionoycin	1.2
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	4.2	93579_CCD1106 (Keratinocytes)_none	1.0
93103_LAK cells_resting	1.2	93580_CCD1106	3.6

		(Keratinocytes)_TNFa and IFNg **	
93788_LAK cells_IL-2	3.7	93791_Liver Cirrhosis	84.6
93787_LAK cells_IL-2+IL-12	2.2	93792_Lupus Kidney	33.9
93789_LAK cells_IL-2+IFN gamma	3.0	93577_NCI-H292	8.6
93790_LAK cells_IL-2+ IL-18	2.6	93358_NCI-H292_IL-4	7.1
93104_LAK cells_PMA/ionomycin and IL-18	1.0	93360_NCI-H292_IL-9	6.5
93578_NK Cells_IL-2_resting	1.5	93359_NCI-H292_IL-13	2.8
93109_Mixed Lymphocyte Reaction_Two Way MLR	2.2	93357_NCI-H292_IFN gamma	3.1
93110_Mixed Lymphocyte Reaction_Two Way MLR	1.2	93777_HPAEC_-	1.7
93111_Mixed Lymphocyte Reaction_Two Way MLR	1.3	93778_HPAEC_IL-1 beta/TNA alpha	1.4
93112_Mononuclear Cells (PBMCs)_resting	0.7	93254_Normal Human Lung Fibroblast_none	4.3
93113_Mononuclear Cells (PBMCs)_PWM	0.8	93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	4.9
93114_Mononuclear Cells (PBMCs)_PHA-L	0.6	93257_Normal Human Lung Fibroblast_IL-4	2.2
93249_Ramos (B cell)_none	1.9	93256_Normal Human Lung Fibroblast_IL-9	1.2
93250_Ramos (B cell)_ionomycin	1.4	93255_Normal Human Lung Fibroblast_IL-13	1.6
93349_B lymphocytes_PWM	1.2	93258_Normal Human Lung Fibroblast_IFN gamma	1.9
93350_B lymphocytes_CD40L and IL-4	2.2	93106_Dermal Fibroblasts CCD1070_resting	3.3
92665_EOL-1 (Eosinophil)_dbcAMP differentiated	2.4	93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	4.7
93248_EOL-1 (Eosinophil)_dbcAMP/PMA ionomycin	2.6	93105_Dermal Fibroblasts CCD1070_IL-1 beta 1 ng/ml	0.7
93356_Dendritic Cells_none	2.2	93772_dermal fibroblast_IFN gamma	0.8
93355_Dendritic Cells_LPS 100 ng/ml	2.1	93771_dermal fibroblast_IL-4	2.4
93775_Dendritic Cells_anti-CD40	1.8	93260_IBD Colitis 2	11.6
93774_Monocytes_resting	1.5	93261_IBD Crohns	14.2
93776_Monocytes_LPS 50 ng/ml	0.6	735010_Colon_normal	61.0
93581_Macrophages_resting	1.7	735019_Lung_none	3.6
93582_Macrophages_LPS 100 ng/ml	1.1	64028-1_Thymus_none	100.0
93098_HUVEC (Endothelial)_none	1.3	64030-1_Kidney_none	5.7

93099_HUVEC (Endothelial)_starved	1.2		
--------------------------------------	-----	--	--

Table 51. Panel 4.1D

Tissue Name	Relative Expression(%)	
	4.1dx4tm5986 f_ag3797_a1	4.1dtm6034f_ ag3797
93768_Secondary Th1_anti-CD28/anti-CD3	8.9	1.3
93769_Secondary Th2_anti-CD28/anti-CD3	2.6	1.3
93770_Secondary Tr1_anti-CD28/anti-CD3	3.1	0.9
93573_Secondary Th1_resting day 4-6 in IL-2	1.5	1.6
93572_Secondary Th2_resting day 4-6 in IL-2	3.4	0.7
93571_Secondary Tr1_resting day 4-6 in IL-2	3.4	0.9
93568_primary Th1_anti-CD28/anti-CD3	3.2	0.3
93569_primary Th2_anti-CD28/anti-CD3	2.0	1.1
93570_primary Tr1_anti-CD28/anti-CD3	2.7	1.0
93565_primary Th1_resting dy 4-6 in IL-2	3.2	0.4
93566_primary Th2_resting dy 4-6 in IL-2	4.5	0.0
93567_primary Tr1_resting dy 4-6 in IL-2	2.3	0.7
93351_CD45RA CD4 lymphocyte_anti-CD28/anti-CD3	2.0	0.7
93352_CD45RO CD4 lymphocyte_anti-CD28/anti-CD3	1.8	2.0
93251_CD8 Lymphocytes_anti-CD28/anti-CD3	5.6	0.9
93353_chronic CD8 Lymphocytes 2ry_resting dy 4-6 in IL-2	4.6	1.1
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	1.8	0.2
93354_CD4_none	7.2	1.3
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	6.6	1.1
93103_LAK cells_resting	7.6	0.4
93788_LAK cells_IL-2	4.8	0.3
93787_LAK cells_IL-2+IL-12	5.7	0.8
93789_LAK cells_IL-2+IFN gamma	5.5	0.2
93790_LAK cells_IL-2+ IL-18	1.6	0.4
93104_LAK cells_PMA/ionomycin and IL-18	2.7	1.3
93578_NK Cells IL-2_resting	5.6	2.0
93109_Mixed Lymphocyte Reaction_Two Way MLR	4.9	2.0
93110_Mixed Lymphocyte Reaction_Two Way MLR	0.5	0.8
93111_Mixed Lymphocyte Reaction_Two Way MLR	6.0	0.2
93112_Mononuclear Cells (PBMCs)_resting	1.3	0.4
93113_Mononuclear Cells (PBMCs)_PWM	7.9	0.5
93114_Mononuclear Cells (PBMCs)_PHA-L	5.1	0.5
93249_Ramos (B cell)_none	5.7	0.6
93250_Ramos (B cell)_ionomycin	4.4	0.3
93349_B lymphocytes_PWM	1.1	0.2
93350_B lymphocytes_CD40L and IL-4	4.3	0.6
92665_EOL-1 (Eosinophil)_dbcAMP differentiated	8.4	3.5
93248_EOL-1 (Eosinophil)_dbcAMP/PMAionomycin	7.2	5.1

93356_Dendritic Cells_none	3.4	1.0
93355_Dendritic Cells_LPS 100 ng/ml	5.5	0.5
93775_Dendritic Cells_anti-CD40	2.6	0.3
93774_Monocytes_resting	1.1	0.9
93776_Monocytes_LPS 50 ng/ml	2.6	0.3
93581_Macrophages_resting	5.2	0.2
93582_Macrophages_LPS 100 ng/ml	1.4	0.3
93098_HUVEC (Endothelial)_none	1.2	0.2
93099_HUVEC (Endothelial)_starved	3.4	0.2
93100_HUVEC (Endothelial)_IL-1b	2.4	0.1
93779_HUVEC (Endothelial)_IFN gamma	2.6	1.2
93102_HUVEC (Endothelial)_TNF alpha + IFN gamma	0.0	0.3
93101_HUVEC (Endothelial)_TNF alpha + IL4	1.6	0.4
93781_HUVEC (Endothelial)_IL-11	2.2	0.4
93583_Lung Microvascular Endothelial Cells_none	1.8	0.9
93584_Lung Microvascular Endothelial Cells_TNFa (4 ng/ml) and IL1b (1 ng/ml)	2.2	0.3
92662_Microvascular Dermal endothelium_none	1.4	0.5
92663_Microvascular Dermal endothelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	2.2	0.3
93773_Bronchial epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml) **	17.8	0.5
93347_Small Airway Epithelium_none	1.5	0.2
93348_Small Airway Epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	3.0	0.6
92668_Coronary Artery SMC_resting	1.1	0.6
92669_Coronary Artery SMC_TNFa (4 ng/ml) and IL1b (1 ng/ml)	1.9	0.8
93107_astrocytes_resting	2.5	1.5
93108_astrocytes_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.5	1.2
92666_KU-812 (Basophil)_resting	4.3	0.8
92667_KU-812 (Basophil)_PMA/ionoycin	3.0	0.6
93579_CCD1106 (Keratinocytes)_none	1.6	0.9
93580_CCD1106 (Keratinocytes)_TNFa and IFNg **	2.4	0.0
93791_Liver Cirrhosis	76.6	9.8
93577_NCI-H292	5.4	4.1
93358_NCI-H292_IL-4	10.3	0.6
93360_NCI-H292_IL-9	16.5	1.3
93359_NCI-H292_IL-13	8.5	3.6
93357_NCI-H292_IFN gamma	5.8	3.4
93777_HPAEC_-	1.2	1.0
93778_HPAEC_IL-1 beta/TNA alpha	1.5	0.3
93254_Normal Human Lung Fibroblast_none	2.5	0.8
93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	5.7	0.9
93257_Normal Human Lung Fibroblast_IL-4	2.9	0.4
93256_Normal Human Lung Fibroblast_IL-9	2.5	0.4

93255_Normal Human Lung Fibroblast_IL-13	2.7	1.9
93258_Normal Human Lung Fibroblast_IFN gamma	0.0	2.2
93106_Dermal Fibroblasts CCD1070_resting	6.1	3.4
93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	2.1	3.8
93105_Dermal Fibroblasts CCD1070_IL-1 beta 1 ng/ml	4.0	1.5
93772_dermal fibroblast_IFN gamma	1.6	0.9
93771_dermal fibroblast_IL-4	1.4	1.3
93892_Dermal fibroblasts_none	2.3	1.5
99202_Neutrophils_TNFa+LPS	0.0	1.7
99203_Neutrophils_none	0.8	0.6
735010_Colon_normal	21.7	6.7
735019_Lung_none	3.7	10.6
64028-1_Thymus_none	11.7	27.0
64030-1_Kidney_none	<b>100.0</b>	<b>100.0</b>

**Table 52. Panel CNS\_Neurodegeneration\_v1.0**

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	tm7142f_ ag3797_b2		tm7142f_ ag3797_b2
AD 1 Hippo	53.6	Control (Path) 3 Temporal Ctx	12.5
AD 2 Hippo	69.7	Control (Path) 4 Temporal Ctx	62.2
AD 3 Hippo	25.6	AD 1 Occipital Ctx	48.0
AD 4 Hippo	33.9	AD 2 Occipital Ctx (Missing)	0.0
AD 5 Hippo	91.6	AD 3 Occipital Ctx	13.0
AD 6 Hippo	39.7	AD 4 Occipital Ctx	40.7
Control 2 Hippo	38.4	AD 5 Occipital Ctx	51.9
Control 4 Hippo	59.4	AD 6 Occipital Ctx	28.4
Control (Path) 3 Hippo	7.8	Control 1 Occipital Ctx	2.6
AD 1 Temporal Ctx	41.0	Control 2 Occipital Ctx	<b>100.0</b>
AD 2 Temporal Ctx	70.4	Control 3 Occipital Ctx	33.1
AD 3 Temporal Ctx	21.3	Control 4 Occipital Ctx	18.6
AD 4 Temporal Ctx	46.7	Control (Path) 1 Occipital Ctx	82.7
AD 5 Inf Temporal Ctx	92.1	Control (Path) 2 Occipital Ctx	18.7
AD 5 Sup Temporal Ctx	74.8	Control (Path) 3 Occipital Ctx	3.4
AD 6 Inf Temporal Ctx	44.5	Control (Path) 4 Occipital Ctx	49.0
AD 6 Sup Temporal Ctx	57.9	Control 1 Parietal	19.8

Control 1 Temporal Ctx	22.8	Control 2 Parietal	60.6
Control 2 Temporal Ctx	45.7	Control 3 Parietal	31.0
Control 3 Temporal Ctx	13.8	Control (Path) 1 Parietal	57.3
Control 3 Temporal Ctx	51.4	Control (Path) 2 Parietal	31.0
Control (Path) 1 Temporal Ctx	62.8	Control (Path) 3 Parietal	5.7
Control (Path) 2 Temporal Ctx	41.4	Control (Path) 4 Parietal	52.1

**Panel 1.3D Summary Ag3086** The NOV5gene is highly expressed in both fetal and adult liver tissue (CTs = 26) and liver cancer cell lines (CT = 27). The gene is also expressed at moderate to low levels in most of the other tissues in the panel. Thus, since the NOV5gene appears to be highly expressed in liver tissue, it could therefore be used to distinguish liver derived tissue from other tissues. The NOV5gene product may also be a potential therapeutic treatment of liver disease.

Among tissues involved in the central nervous system, the NOV5gene is moderately expressed in the fetal and adult brain, including the adult thalamus, substantia nigra, hippocampus, amygdala and is also expressed at low but significant levels in the cerebellum and cerebral cortex. This expression profile suggests that the NOV5gene has functional significance in the CNS. The close homologue to the NOV5 gene product, hepatocyte growth factor, has numerous therapeutic applications in the CNS, including prevention of neuronal death in animal models of stroke and ischemia. Hepatocyte growth factor has mitogenic activity, crossing the blood brain barrier when disrupted, and thus has potential application as a protein therapeutic to treat brain pathologies when administered directly to the cortico spinal fluid or systemically when the blood brain barrier is disrupted. Hepatocyte growth factor-like protein is a neurotrophic factor useful in the prevention of motoneuron atrophy upon axotomy. Therefore, the protein encoded by the NOV5gene may be useful as a therapeutic agent in treating stroke and neurodegenerative diseases including Alzheimer's disease, Parkinson's disease, and Huntington's disease. The potential role of the NOV5gene or its protein product in brain plasticity and regeneration affords utility in treating brain damage and aging related disorders, such as memory impairment that has hippocampal dysfunction as its primary focus.

25

**General\_Screening\_Panel\_1.4 Ag3797** The expression of the NOV5gene in panel 1.4 appears to be highest in a sample derived from a liver cancer cell line (HepG2) (CT = 25.3). In addition there is substantial expression of this gene associated with other liver derived material (adult liver CT=27.2; fetal liver CT=26.5). Thus, the expression of the NOV5gene



could be used to distinguish liver derived specimens from other samples. In addition, therapeutic modulation of this gene might be of benefit in the treatment of liver related disorders, such as cirrhosis.

5           **Panel 2.2 Summary Ag3086** The expression of the NOV5gene appears to be highest in a sample derived from a liver cancer specimen (CT=26) and is also significant in a number of samples derived from liver tissue. This result is consistent with what is seen in Panels 1.4 and 2D. In addition there appears to be substantial expression of this gene associated with normal kidney tissue (CT=27.2) when compared to adjacent kidney cancer specimens. Thus, 10 this gene could be used to distinguish liver tissue from non-liver tissue as well as distinguish normal kidney tissue when compared to adjacent kidney cancer. Moreover, therapeutic modulation of the expression of the NOV5gene or function of its product might be of benefit in the treatment of kidney cancer.

15           **Panel 4D Summary Ag3086** The NOV5gene is highly expressed in the thymus (CT = 24), colon (CT = 28.4), and IBD Colitis 2 (CT = 27.2) and is expressed at lower levels in mature T cells. The NOV5gene encodes a putative hepatocyte like growth factor homologue. There are reports that hepatocyte growth factor (HGF) is expressed in the thymus and colon. In the thymus, HGF may promote T cell production and in the colon, overexpression of HGF 20 has been shown to leads to IBD like disease in mice. Therapies designed with the protein encoded for by the NOV5gene could be important in the regulation of T cell development and immune function and be useful in organ transplantation. In addition, blocking the function of the NOV5 gene product could help in the treatment of IBD colitis.

25           **Panel 4.1D Summary Ag3797** Results from two experiments using the same probe and primer set are in very good agreement. In both experiments, highest expression of the NOV5gene is detected in kidney (CT=29, 27.4). Moderate expression is also detected in liver cirrhosis (CT=29.4, 30.7). Moderate to low expression of the gene is detected in many of the tissues in this panel. Thus, expression of the NOV5gene could be used to distinguish those 30 tissues from other tissues.

**Panel CNS\_Neurodegeneration\_v1.0 Summary Ag3797** Highest expression of the NOV5 gene is detected in the occipital cortex of a control patient (CT=31.3). Moderate to low expression is detected throughout the tissue samples in this panel. Please see panel 1.3 for a 35 discussion of potential utility of this gene with regards to the CNS.

## References:

1. Korhonen L, Sjöholm U, Takei N, Kern MA, Schirmacher P, Castren E, Lindholm  
5 D. (2000) *Eur J Neurosci.* 12:3453-61.

Hepatocyte growth factor-scatter factor (HGF) is expressed in different parts of the nervous system, and has been shown to exhibit neurotrophic activity. Here we show that c-Met, the receptor for HGF, is expressed in developing rat hippocampus, with the highest levels  
10 during the first postnatal weeks. To study the function of HGF, hippocampal neurons were prepared from embryonic rats and treated with different HGF concentrations. In these cultures, HGF increased the number of neurons expressing the 28-kDa calcium-binding protein (calbindin D) in a dose-dependent manner. The effect of HGF was larger than that observed with either brain-derived neurotrophic factor (BDNF) or neurotrophin-3 (NT-3), and  
15 cotreatment of the cultures with HGF and the neurotrophins was additive with respect to calbindin D neurons. Besides affecting the number of neurons, HGF significantly increased the degree of sprouting of calbindin D-positive neurons, suggesting an influence on neuronal maturation. BDNF and NT-3 stimulated neurite outgrowth of calbindin D neurons to a much smaller degree. In contrast to calbindin D neurons, HGF did not significantly increase the  
20 number of neurons immunoreactive with the neurotransmitter gamma-aminobutyric acid (GABA) in the hippocampal cultures. Immunohistochemical studies showed that c-Met-, calbindin D- and HGF-immunoreactive cells are all present in the dentate gyrus and partly colocalize within neurons. These results show that HGF acts on calbindin D-containing hippocampal neurons and increases their neurite outgrowth, suggesting that HGF plays an  
25 important role for the maturation and function of these neurons in the hippocampus.

PMID: 11029614

2. Powell EM, Mars WM, Levitt P. (2001) *Neuron* 30:79-89.

Cortical interneurons arise from the proliferative zone of the ventral telencephalon, the  
30 ganglionic eminence, and migrate into the developing neocortex. The spatial patterns of migratory interneurons reflect the complementary expression of hepatocyte growth factor/scatter factor (HGF/SF) and its receptor, MET, in the forebrain. Scatter assays on forebrain explants demonstrate regionally specific motogenic activity due to HGF/SF. In addition, exogenous ligand disrupts normal cell migration. Mice lacking the urokinase-type  
35 plasminogen activator receptor (u-PAR), a key component of HGF/SF activation, exhibit

deficient scatter activity in the forebrain, abnormal interneuron migration from the ganglionic eminence, and reduced interneurons in the frontal and parietal cortex. The data suggest that HGF/SF motogenic activity, which is essential for normal development of other organ systems, is a conserved mechanism that regulates trans-telencephalic migration of interneurons.

PMID: 11343646

3. Stella MC, Vercelli A, Repici M, Follenzi A, Comoglio PM. (2001) *Mol Biol Cell* 12:1341-52.

Macrophage stimulating protein (MSP), also known as hepatocyte growth factor-like, is a soluble cytokine that belongs to the family of the plasminogen-related growth factors (PRGFs). PRGFs are alpha/beta heterodimers that bind to transmembrane tyrosine kinase receptors. MSP was originally isolated as a chemotactic factor for peritoneal macrophages. Through binding to its receptor, encoded by the RON gene, it stimulates dissociation of epithelia and works as an inflammatory mediator by repressing the production of nitric oxide (NO). Here, we identify a novel role for MSP in the central nervous system. As a paradigm to analyze this function we chose the hypoglossal system of adult mice. We demonstrate in vivo that either administration of exogenous MSP or transplantation of MSP-producing cells at the proximal stump of the resected nerve is sufficient to prevent motoneuron atrophy upon axotomy. We also show that the MSP gene is expressed in the tongue, the target of the hypoglossal nerve, and that MSP induces biosynthesis of Ron receptor in the motoneuron somata. Finally, we show that MSP suppresses NO production in the injured hypoglossal nuclei. Together, these data suggest that MSP is a novel neurotrophic factor for cranial motoneurons and, by regulating the production of NO, may have a role in brain plasticity and regeneration.

PMID: 11359926

4. Kern MA, Bamborschke S, Nekic M, Schubert D, Rydin C, Lindholm D, Schirmacher P. (2001) *Cytokine* 14:170-6.

Hepatocyte growth factor (HGF) and its specific receptor, MET, are expressed in the developing and adult mammalian brain. Recent studies have shown a neurotrophic activity of HGF in the nervous system. The present study focused on HGF concentrations in the cerebrospinal fluid (CSF) and serum in normal persons and in different central nervous system

(CNS) diseases considering blood-CSF barrier (BCB) function. Concentrations of HGF were analyzed using an enzyme-linked immunosorbent assay (ELISA). HGF was present in normal human CSF (346+/-126 pg/ml) representing approximately half of the HGF serum concentrations. The CSF HGF levels were not significantly changed in chronic CNS disease and in aseptic meningitis (419+/-71 pg/ml), but significantly increased in patients with bacterial meningitis (6101+/- 5200 pg/ml). The HGF levels in CSF were not influenced by increased serum concentrations in patients with normal or mildly affected BCB function. The results show that HGF is present in normal CSF and does not appear to cross the CSF barrier significantly unless it is severely disrupted. So far, strong increases of HGF concentration in CSF are only present in acute bacterial meningitis. Copyright 2001 Academic Press.

PMID: 11396995

5. Hayashi K, Morishita R, Nakagami H, Yoshimura S, Hara A, Matsumoto K, Nakamura T, Ogihara T, Kaneda Y, Sakai N. (2001) *Gene Ther* 8:1167-73.

To develop a novel strategy to prevent delayed neuronal death (DND) following transient occlusion of arteries, the gene of hepatocyte growth factor (HGF), a novel neurotrophic factor, was transfected into the subarachnoid space of gerbils after transient forebrain ischemia. Importantly, transfection of HGF gene into the subarachnoid space prevented DND, accompanied by a significant increase in HGF in the cerebrospinal fluid. Prevention of DND by HGF is due to the inhibition of apoptosis through the blockade of bax translocation from the cytoplasm to the nucleus. HGF gene transfer into the subarachnoid space may provide a new therapeutic strategy for cerebrovascular disease.

PMID: 11509947

6. Tamura S, Sugawara T, Tokoro Y, Taniguchi H, Fukao K, Nakauchi H, Takahama Y. (1998) *Scand J Immunol*. 47:296-301.

The c-Met oncoprotein is a cell-surface receptor for hepatocyte growth factor (HGF). Signals through HGF and c-Met have been appreciated for their crucial roles in the development of many cell types, including liver cells. The present study examined whether c-Met is expressed in the thymus and whether c-Met/HGF signals can regulate T-cell development in the thymus. We have found that mRNA transcripts encoding c-Met are expressed in mouse thymus. The c-Met transcripts were expressed at higher levels in fetal and

neonatal thymus than in adult thymus, and were mostly expressed by lymphoid cells rather than by stromal cells. Interestingly, the addition of HGF to fetal thymus organ cultures increased the generation of mature T cells expressing high levels of T-cell antigen receptors. These results indicate that c-Met is expressed in the thymus during early ontogeny, and that c-Met/HGF signals can promote T-cell development.

PMID: 9600310

7. Takayama H, Takagi H, Larochelle WJ, Kapur RP, Merlino G. (2001) Lab Invest. 81:297-305.

Hepatocyte growth factor/scatter factor (HGF/SF) can stimulate growth of gastrointestinal epithelial cells in vitro; however, the physiological role of HGF/SF in the digestive tract is poorly understood. To elucidate this in vivo function, mice were analyzed in which an HGF/SF transgene was overexpressed throughout the digestive tract. Nearly a third of all HGF/SF transgenic mice in this study (28 of 87) died by 6 months of age as a result of sporadic intestinal obstruction of unknown etiology. Enteric ganglia were not overtly affected, indicating that the pathogenesis of this intestinal lesion was different from that operating in Hirschsprung's disease. Transgenic mice also exhibited a rectal inflammatory bowel disease (IBD) with a high incidence of anorectal prolapse. Expression of interleukin-2 was decreased in the transgenic colon, indicating that HGF/SF may influence regulation of the local intestinal immune system within the colon. These results suggest that HGF/SF plays an important role in the development of gastrointestinal paresis and chronic intestinal inflammation. HGF/SF transgenic mice may represent a useful model for the study of molecular mechanisms associated with a subset of IBD and intestinal pseudo-obstruction. Moreover, our data identify previously unappreciated side effects that may be encountered when using HGF/SF as a therapeutic agent.

PMID: 11310823

## NOV6

Expression of gene NOV6 was assessed using the primer-probe set Ag2439 described in Table 53. Results from RTQ-PCR runs are shown in Tables 54, 55, 56, and 57.

Table 53. Probe Name Ag2439

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5' - TATCATCACTTGTGATGGCAA - 3'	59	22	189	150
Probe	FAM-5' - AAAACCGAGAGCACTTTGAAAACACA-3' - TAMRA	66.1	26	223	151
Reverse	5' - AAACCTCTCTCCAGGGTACAA - 3'	59.1	22	255	152

5

Table 54. Panel 1.3D

Tissue Name	Relative Expression(%) 1.3dtm3781t_ ag2439	Tissue Name	Relative Expression(%) 1.3dtm3781t_ ag2439
Liver adenocarcinoma	11.3	Kidney (fetal)	7.6
Pancreas	3.7	Renal ca. 786-0	4.9
Pancreatic ca. CAPAN 2	4.9	Renal ca. A498	10.4
Adrenal gland	5.3	Renal ca. RXF 393	4.1
Thyroid	7.4	Renal ca. ACHN	2.2
Salivary gland	5.3	Renal ca. UO-31	0.6
Pituitary gland	7.6	Renal ca. TK-10	5.4
Brain (fetal)	23.5	Liver	4.2
Brain (whole)	6.2	Liver (fetal)	12.9
Brain (amygdala)	12.9	Liver ca. (hepatoblast) HepG2	11.9
Brain (cerebellum)	8.5	Lung	35.1
Brain (hippocampus)	100.0	Lung (fetal)	12.3
Brain (substantia nigra)	4.9	Lung ca. (small cell) LX-1	8.8
Brain (thalamus)	9.2	Lung ca. (small cell) NCI-H69	5.6
Cerebral Cortex	12.9	Lung ca. (s.cell var.) SHP-77	21.8
Spinal cord	5.1	Lung ca. (large cell) NCI-H460	5.8
CNS ca. (glio/astro) U87-MG	11.8	Lung ca. (non-sm. cell) A549	14.9
CNS ca. (glio/astro) U-118-MG	15.5	Lung ca. (non-s.cell) NCI-H23	11.9
CNS ca. (astro) SW1783	3.7	Lung ca. (non-s.cell) HOP-62	4.6
CNS ca.* (neuro; met ) SK-N-AS	46.7	Lung ca. (non-s.cl) NCI-H522	2.1
CNS ca. (astro) SF-539	8.8	Lung ca. (squamous) SW 900	3.1
CNS ca. (astro) SNB-75	4.2	Lung ca. (squamous) NCI-H596	3.1
CNS ca. (glio) SNB-19	9.9	Mammary gland	3.9
CNS ca. (glio) U251	5.7	Breast ca.* (pl. effusion) MCF-7	8.6
CNS ca. (glio) SF-295	6.2	Breast ca.* (pl.ef) MDA-MB-231	23.3
Heart (fetal)	6.1	Breast ca.* (pl. effusion) T47D	3.0
Heart	22.2	Breast ca. BT-549	31.2
Fetal Skeletal	40.1	Breast ca. MDA-N	7.6

Skeletal muscle	2.5	Ovary	3.5
Bone marrow	11.3	Ovarian ca. OVCAR-3	5.0
Thymus	6.7	Ovarian ca. OVCAR-4	0.3
Spleen	12.2	Ovarian ca. OVCAR-5	8.9
Lymph node	14.1	Ovarian ca. OVCAR-8	9.3
Colorectal	22.8	Ovarian ca. IGROV-1	5.4
Stomach	10.4	Ovarian ca.* (ascites) SK-OV-3	12.6
Small intestine	21.2	Uterus	9.3
Colon ca. SW480	2.1	Placenta	17.7
Colon ca.* (SW480 met)SW620	9.3	Prostate	5.1
Colon ca. HT29	8.4	Prostate ca.* (bone met)PC-3	3.7
Colon ca. HCT-116	6.0	Testis	5.1
Colon ca. CaCo-2	16.7	Melanoma Hs688(A).T	1.5
83219 CC Well to Mod Diff (ODO3866)	13.5	Melanoma* (met) Hs688(B).T	1.0
Colon ca. HCC-2998	32.1	Melanoma UACC-62	0.7
Gastric ca.* (liver met) NCI-N87	34.2	Melanoma M14	4.6
Bladder	14.0	Melanoma LOX IMVI	4.4
Trachea	24.8	Melanoma* (met) SK-MEL-5	45.4
Kidney	1.7	Adipose	11.7

Table 55. Panel 2D

Tissue Name	Relative Expression(%) 2dtm3782t_ ag2439	Tissue Name	Relative Expression(%) 2dtm3782t_ ag2439
Normal Colon GENPAK 061003	100.0	Kidney NAT Clontech 8120608	1.2
83219 CC Well to Mod Diff (ODO3866)	17.2	Kidney Cancer Clontech 8120613	2.8
83220 CC NAT (ODO3866)	13.8	Kidney NAT Clontech 8120614	1.6
83221 CC Gr.2 rectosigmoid (ODO3868)	16.6	Kidney Cancer Clontech 9010320	7.7
83222 CC NAT (ODO3868)	3.7	Kidney NAT Clontech 9010321	3.3
83235 CC Mod Diff (ODO3920)	31.2	Normal Uterus GENPAK 061018	5.2
83236 CC NAT (ODO3920)	17.2	Uterus Cancer GENPAK 064011	22.1
83237 CC Gr.2 ascend colon (ODO3921)	74.2	Normal Thyroid Clontech A+ 6570-1	10.5
83238 CC NAT (ODO3921)	15.3	Thyroid Cancer GENPAK 064010	17.3
83241 CC from Partial Hepatectomy (ODO4309)	32.5	Thyroid Cancer INVITROGEN A302152	14.5
83242 Liver NAT (ODO4309)	8.2	Thyroid NAT INVITROGEN A302153	16.2
87472 Colon mets to lung	5.3	Normal Breast GENPAK	16.7

(OD04451-01)		061019	
87473 Lung NAT (OD04451-02)	11.1	84877 Breast Cancer (OD04566)	7.3
Normal Prostate Clontech A+ 6546-1	11.6	85975 Breast Cancer (OD04590-01)	22.2
84140 Prostate Cancer (OD04410)	21.0	85976 Breast Cancer Mets (OD04590-03)	24.8
84141 Prostate NAT (OD04410)	21.3	87070 Breast Cancer Metastasis (OD04655-05)	15.3
87073 Prostate Cancer (OD04720-01)	24.1	GENPAK Breast Cancer 064006	18.0
87074 Prostate NAT (OD04720-02)	22.5	Breast Cancer Res. Gen. 1024	17.7
Normal Lung GENPAK 061010	84.1	Breast Cancer Clontech 9100266	13.3
83239 Lung Met to Muscle (ODO4286)	11.3	Breast NAT Clontech 9100265	6.9
83240 Muscle NAT (ODO4286)	9.4	Breast Cancer INVITROGEN A209073	16.3
84136 Lung Malignant Cancer (OD03126)	19.9	Breast NAT INVITROGEN A2090734	11.0
84137 Lung NAT (OD03126)	36.9	Normal Liver GENPAK 061009	7.1
84871 Lung Cancer (OD04404)	28.7	Liver Cancer GENPAK 064003	8.0
84872 Lung NAT (OD04404)	11.7	Liver Cancer Research Genetics RNA 1025	5.1
84875 Lung Cancer (OD04565)	37.4	Liver Cancer Research Genetics RNA 1026	1.3
84876 Lung NAT (OD04565)	13.3	Paired Liver Cancer Tissue Research Genetics RNA 6004-T	5.4
85950 Lung Cancer (OD04237-01)	25.9	Paired Liver Tissue Research Genetics RNA 6004-N	11.9
85970 Lung NAT (OD04237-02)	21.3	Paired Liver Cancer Tissue Research Genetics RNA 6005-T	1.6
83255 Ocular Mel Met to Liver (ODO4310)	9.5	Paired Liver Tissue Research Genetics RNA 6005-N	0.5
83256 Liver NAT (ODO4310)	6.8	Normal Bladder GENPAK 061001	39.8
84139 Melanoma Mets to Lung (OD04321)	11.2	Bladder Cancer Research Genetics RNA 1023	8.9
84138 Lung NAT (OD04321)	30.4	Bladder Cancer INVITROGEN A302173	93.3
Normal Kidney GENPAK 061008	23.2	87071 Bladder Cancer (OD04718-01)	22.5
83786 Kidney Ca, Nuclear grade 2 (OD04338)	14.2	87072 Bladder Normal Adjacent (OD04718-03)	37.1
83787 Kidney NAT (OD04338)	9.7	Normal Ovary Res. Gen.	2.0
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	26.2	Ovarian Cancer GENPAK 064008	21.3
83789 Kidney NAT (OD04339)	11.3	87492 Ovary Cancer (OD04768-07)	33.4



83790 Kidney Ca, Clear cell type (OD04340)	24.5	87493 Ovary NAT (OD04768-08)	4.9
83791 Kidney NAT (OD04340)	17.2	Normal Stomach GENPAK 061017	23.7
83792 Kidney Ca, Nuclear grade 3 (OD04348)	6.7	Gastric Cancer Clontech 9060358	5.6
83793 Kidney NAT (OD04348)	9.2	NAT Stomach Clontech 9060359	18.9
87474 Kidney Cancer (OD04622-01)	7.7	Gastric Cancer Clontech 9060395	34.6
87475 Kidney NAT (OD04622-03)	0.9	NAT Stomach Clontech 9060394	27.4
85973 Kidney Cancer (OD04450-01)	10.1	Gastric Cancer Clontech 9060397	64.2
85974 Kidney NAT (OD04450-03)	7.2	NAT Stomach Clontech 9060396	5.3
Kidney Cancer Clontech 8120607	1.2	Gastric Cancer GENPAK 064005	48.0

Table 56. Panel 4D

Tissue Name	Relative Expression (%)	Tissue Name	Relative Expression (%)
	4dtm3783t_ag2439		4dtm3783t_ag2439
93768_Secondary Th1_anti-CD28/anti-CD3	51.8	93100_HUVEC (Endothelial) IL-1b	5.9
93769_Secondary Th2_anti-CD28/anti-CD3	39.0	93779_HUVEC (Endothelial) IFN gamma	6.0
93770_Secondary Tr1_anti-CD28/anti-CD3	54.0	93102_HUVEC (Endothelial) TNF alpha + IFN gamma	7.2
93573_Secondary Th1_resting day 4-6 in IL-2	4.2	93101_HUVEC (Endothelial) TNF alpha + IL4	18.9
93572_Secondary Th2_resting day 4-6 in IL-2	5.7	93781_HUVEC (Endothelial) IL-11	6.2
93571_Secondary Tr1_resting day 4-6 in IL-2	7.4	93583_Lung Microvascular Endothelial Cells_none	21.8
93568_primary Th1_anti-CD28/anti-CD3	82.4	93584_Lung Microvascular Endothelial Cells_TNFa (4 ng/ml) and IL1b (1 ng/ml)	21.2
93569_primary Th2_anti-CD28/anti-CD3	52.1	92662_Microvascular Dermal endothelium_none	33.7
93570_primary Tr1_anti-CD28/anti-CD3	76.8	92663_Microvascular Dermal endothelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	15.1
93565_primary Th1_resting dy 4-6 in IL-2	36.3	93773_Bronchial epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml) **	1.0
93566_primary Th2_resting dy 4-6 in IL-2	16.8	93347_Small Airway Epithelium_none	4.1
93567_primary Tr1_resting dy	9.1	93348_Small Airway	34.2

4-6 in IL-2		Epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	
93351_CD45RA CD4 lymphocyte_anti-CD28/anti- CD3	18.0	92668_Coronary Artery SMC_resting	4.9
93352_CD45RO CD4 lymphocyte_anti-CD28/anti- CD3	37.1	92669_Coronary Artery SMC_TNFa (4 ng/ml) and IL1b (1 ng/ml)	0.6
93251_CD8 Lymphocytes_anti- CD28/anti-CD3	40.6	93107_astrocytes_resting	5.0
93353_chronic CD8 Lymphocytes 2ry_resting dy 4- 6 in IL-2	41.2	93108_astrocytes_TNFa (4 ng/ml) and IL1b (1 ng/ml)	6.6
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	23.5	92666_KU-812 (Basophil)_resting	17.9
93354_CD4_none	4.8	92667_KU-812 (Basophil)_PMA/ionomycin	43.5
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	8.5	93579_CCD1106 (Keratinocytes)_none	10.2
93103_LAK cells_resting	32.8	93580_CCD1106 (Keratinocytes)_TNFa and IFNg **	0.7
93788_LAK cells_IL-2	29.1	93791_Liver Cirrhosis	2.6
93787_LAK cells_IL-2+IL-12	26.2	93792_Lupus Kidney	1.7
93789_LAK cells_IL-2+IFN gamma	25.3	93577_NCI-H292	10.0
93790_LAK cells_IL-2+ IL-18	28.5	93358_NCI-H292_IL-4	18.0
93104_LAK cells_PMA/ionomycin and IL- 18	12.8	93360_NCI-H292_IL-9	15.6
93578_NK Cells IL-2_resting	10.5	93359_NCI-H292_IL-13	5.8
93109_Mixed Lymphocyte Reaction_Two Way MLR	21.5	93357_NCI-H292_IFN gamma	7.1
93110_Mixed Lymphocyte Reaction_Two Way MLR	14.8	93777_HPAEC_-	12.2
93111_Mixed Lymphocyte Reaction_Two Way MLR	14.4	93778_HPAEC_IL-1 beta/TNA alpha	13.8
93112_Mononuclear Cells (PBMCs)_resting	3.9	93254_Normal Human Lung Fibroblast_none	4.8
93113_Mononuclear Cells (PBMCs)_PWM	75.8	93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	2.3
93114_Mononuclear Cells (PBMCs)_PHA-L	36.3	93257_Normal Human Lung Fibroblast_IL-4	12.5
93249_Ramos (B cell)_none	44.1	93256_Normal Human Lung Fibroblast_IL-9	10.2
93250_Ramos (B cell)_ionomycin	100.0	93255_Normal Human Lung Fibroblast_IL-13	7.6
93349_B lymphocytes_PWM	93.3	93258_Normal Human Lung Fibroblast_IFN gamma	10.9
93350_B lymphocytes_CD40L and IL-4	22.4	93106_Dermal Fibroblasts CCD1070_resting	12.7

92665_EOL-1 (Eosinophil)_dbcAMP differentiated	6.1	93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	32.3
93248_EOL-1 (Eosinophil)_dbcAMP/PMAion omycin	16.2	93105_Dermal Fibroblasts CCD1070_IL-1 beta 1 ng/ml	4.9
93356_Dendritic Cells_none	16.2	93772_dermal fibroblast_IFN gamma	6.3
93355_Dendritic Cells_LPS 100 ng/ml	30.8	93771_dermal fibroblast_IL-4	9.8
93775_Dendritic Cells_anti- CD40	39.0	93260_IBD Colitis 2	1.5
93774_Monocytes_resting	4.9	93261_IBD Crohns	1.2
93776_Monocytes_LPS 50 ng/ml	6.8	735010_Colon_normal	11.0
93581_Macrophages_resting	41.5	735019_Lung_none	17.9
93582_Macrophages_LPS 100 ng/ml	7.1	64028-1_Thymus_none	11.4
93098_HUVEC (Endothelial)_none	12.2	64030-1_Kidney_none	41.5
93099_HUVEC (Endothelial)_starved	20.0		

Table 57.Panel CNS\_neurodegeneration\_v1.0

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
	tm6902t ag2439_a2s1		tm6902t ag2439_a2s1
106655_4951 Hippo	1.7	106677_4624 BA21	13.8
106657_4986 Hippo	5.4	106681_4640 BA21	2.5
106652_4933 Hippo	10.3	106654_4951 BA17	20.2
106649_4901 Hippo	2.4	ens_water	0.0
110138_3087 hippo	100.0	106651_4933 BA17	14.8
110121_3027 Hippo	0.0	106648_4901 BA17	2.2
106670_4971 Hippo	2.4	110140_3087 occ ctx	4.3
106666_4867 Hippo	0.9	110123_3027 Occ Ctx	0.0
106680_4624 Hippo	0.0	106659_4595 BA17	23.7
106653_4951 BA21	32.6	106668_4971 BA17	5.2
106656_4986 BA21	1.7	106662_4737 BA17	31.4
106650_4933 BA21	26.2	106665_4867 BA17	11.1
106647_4901 BA21	24.3	106675_3975 BA17	56.4
110136_3087 inf temp ctx	1.6	106672_3954 BA17	6.8
110137_3087 sup temp ctx	8.1	106678_4624 BA17	9.0
110118_3027 Inf Temp Ctx	1.5	106682_4640 BA17	29.9
110119_3027 Sup Temp Ctx	0.0	106660_4595 BA7	14.7
106658_4595 BA21	14.7	113670_106669 pool	53.3
106667_4971 BA21	10.4	106663_4737 BA7	1.8
106661_4737 BA21	31.6	106676_3975 BA7	39.3

106664_4867 BA21	1.5	106673_3954 BA7	0.5
106674_3975 BA21	37.3	106679_4624 BA7	18.2
106671_3954 BA21	58.2	106683_4640 BA7	93.7

**Panel 1.3D Summary Ag2439** The NOV6 gene is widely expressed across the samples in this panel, with highest expression detected in the hippocampus (CT=28.1). This gene is also expressed at lower levels in the other tissues originating from the central nervous system, including the amygdala, cerebellum, cerebral cortex, substantia nigra, thalamus, and spinal cord. Thus, NOV6 gene expression may be used to distinguish hippocampus from other tissues. Please see CNS\_neurodegeneration\_panel\_v1.0 summary for a discussion of the potential utility of this gene in CNS disorders.

Among tissues involved in metabolic function, the NOV6 gene is also expressed in thyroid, adrenal gland, pituitary gland, pancreas, heart (adult and fetal), liver (adult and fetal), and adipose. Interestingly, this gene is more highly expressed in fetal skeletal muscle (CT = 29.4) than in adult skeletal muscle (CT = 33.4). This observation suggests that the NOV6 protein product may enhance muscular growth or development in the fetus and thus may also act in a regenerative capacity in the adult. Therefore, therapeutic modulation of the NOV6 gene could be useful in treatment of muscular related disease. More specifically, treatment of weak or dystrophic muscle with the protein encoded by this gene could restore muscle mass or function.

The NOV6 gene is also expressed at higher levels in gastric, colon, melanoma, lung, and colon cancer cell lines than in normal tissues. Therefore, this gene may be used as a marker for gastric cancer, colon cancer, melanoma, lung, and colon cancer cell lines. In addition, therapeutic modulation of the NOV6 gene product might be of use in the treatment of these cancers.

**Panel 2D Summary Ag2439** The NOV6 gene is most highly expressed in a sample derived from normal colon (CT=26). However, in general this gene appears to be more highly expressed in cancers than in normal tissues. Specifically, NOV6 gene expression is slightly higher in lung cancer (squamous cell type), gastric cancer, ovarian cancer, a kidney cancer sample and a sample of breast cancer relative to the normal controls. Thus, the expression of this gene could be used to distinguish malignant colon, lung, stomach, ovary and some breast and kidney tissue from normal tissue from these organs. In addition, therapeutic modulation of the NOV6 gene product might be of use in the treatment of these cancers.

**Panel 4D Summary Ag2439** The NOV6 gene is most highly expressed in ionomycin-treated Ramos B cells (CT = 26). This gene is also expressed at moderate levels in T cells, monocytes, dendritic cells, endothelial cells, smooth muscle cells, and airway epithelial cells both under resting and cytokine-stimulated conditions. Therefore, this gene may be useful as a marker for these resting and activated cells.

**Panel CNS\_neurodegeneration\_v1.0 Summary Ag2439** The NOV6 gene encodes a protein with homology to fatty acid binding protein and is expressed across the brain, although expression appears to be the highest in the hippocampus (Panel 1.3D). This gene does not appear to be differentially expressed in Alzheimer's disease based on the results from panel CNS\_Neurodegeneration\_V1.0, although the transcript is detected at low levels in many of the brain samples. Fatty acid binding protein expression is increased in development during axon growth and during the response to injury, probably for the transport of fatty acids for use as membrane components (ref. 1). Therefore, upregulation of the NOV6 gene or its protein product may be beneficial during neurite outgrowth and synaptogenesis in response to neuronal death or injury (Parkinson's disease, Alzheimer's disease, Huntington's disease, spinocerebellar ataxia, stroke, and head/spinal cord trauma).

#### References:

1. Liu Y, Molina CA, Welcher AA, Longo LD, De Leon M. Expression of DA11, a neuronal-injury-induced fatty acid binding protein, coincides with axon growth and neuronal differentiation during central nervous system development. *J Neurosci Res* 1997 Jun 15;48(6):551-62.

DA11 is the first fatty acid binding protein (FABP) for which gene expression has been shown to be upregulated following neuronal injury in the adult peripheral nervous system. To understand better the potential regulatory role(s) of this unique FABP in axonal growth and neuronal differentiation, we undertook a temporal and spatial study of DA11 gene expression in the developing rat central nervous system (CNS). Transient upregulation of DA11 mRNA and protein levels in CNS tissues were quantified by Northern blot hybridization and Western immunoblot analyses at different developmental ages. Homogenates of embryonic and neonatal cerebral cortex, cerebellum, brain stem, and hippocampal tissues contained 100-fold more DA11 mRNA and protein than corresponding adult tissues. Significant increase in DA11 mRNA was observed as early as embryonic day (E) 14 in cerebral cortex and cerebellum and

E19 in brain stem and hippocampus. Postnatal levels of DA11 remained elevated through postnatal day (P) 10 in cerebral cortex, P14 in brain stem and hippocampus, and P20 in cerebellum. Localization of DA11-like immunoreactivity to specific CNS tissues, cell types, and intracellular compartments at P9 revealed a spatial pattern of neuronal expression different than that reported for other FABPs. DA11 protein was detected in the nucleus, cytoplasm, axons, and dendrites of differentiating neurons in cerebral cortex, hippocampus, cerebellum, brain stem, spinal cord, and olfactory bulb. The strong association of DA11 gene expression with development throughout the CNS suggests that this unique FABP plays an important role in axonal growth and neuronal differentiation in many different neuronal populations.

PMID: 9210525

## NOV9

Expression of gene NOV9 was assessed using the primer-probe set Ag2771 described in Table 58. Results from RTQ-PCR runs are shown in Tables 59, 60, 61, and 62.

**Table 58. Probe Name Ag2771**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5'-TGAACAGAACTATGCGAAACAA-3'	58.5	22	223	153
Probe	FAM-5'-TCTGGTTAAGAAGTACTGCCCCAAACG-3'-TAMRA	68	27	253	154
Reverse	5'-GGCTCTTCATCTTTGGATGAA-3'	59.3	21	280	155

**Table 59. Panel 1.3D**

Tissue Name	Relative Expression(%) 1.3Dx4tm4869 f_ag2771_a1	Tissue Name	Relative Expression(%) 1.3Dx4tm4869 f_ag2771_a1
Liver adenocarcinoma	12.7	Kidney (fetal)	32.3
Pancreas	3.2	Renal ca. 786-0	22.1
Pancreatic ca. CAPAN 2	4.7	Renal ca. A498	13.0
Adrenal gland	2.1	Renal ca. RXF 393	13.1
Thyroid	14.5	Renal ca. ACHN	4.0
Salivary gland	10.5	Renal ca. UO-31	15.9
Pituitary gland	2.5	Renal ca. TK-10	22.2
Brain (fetal)	16.8	Liver	1.2

Brain (whole)	5.0	Liver (fetal)	3.7
Brain (amygdala)	5.0	Liver ca. (hepatoblast) HepG2	20.9
Brain (cerebellum)	8.1	Lung	9.9
Brain (hippocampus)	7.7	Lung (fetal)	23.0
Brain (substantia nigra)	2.1	Lung ca. (small cell) LX-1	20.5
Brain (thalamus)	5.5	Lung ca. (small cell) NCI-H69	14.0
Cerebral Cortex	28.3	Lung ca. (s.cell var.) SHP-77	38.9
Spinal cord	17.3	Lung ca. (large cell) NCI-H460	4.6
CNS ca. (glio/astro) U87-MG	2.2	Lung ca. (non-sm. cell) A549	9.5
CNS ca. (glio/astro) U-118-MG	5.1	Lung ca. (non-s.cell) NCI-H23	20.1
CNS ca. (astro) SW1783	21.6	Lung ca (non-s.cell) HOP-62	10.3
CNS ca.* (neuro; met ) SK-N-AS	26.8	Lung ca. (non-s.cl) NCI-H522	36.6
CNS ca. (astro) SF-539	12.7	Lung ca. (squam.) SW 900	21.3
CNS ca. (astro) SNB-75	12.3	Lung ca. (squam.) NCI-H596	11.8
CNS ca. (glio) SNB-19	34.0	Mammary gland	21.5
CNS ca. (glio) U251	20.2	Breast ca.* (pl. effusion) MCF-7	25.3
CNS ca. (glio) SF-295	12.4	Breast ca.* (pl.ef) MDA-MB-231	10.5
Heart (fetal)	6.2	Breast ca.* (pl. effusion) T47D	13.3
Heart	14.1	Breast ca. BT-549	8.0
Fetal Skeletal	23.1	Breast ca. MDA-N	13.2
Skeletal muscle	20.0	Ovary	15.2
Bone marrow	0.7	Ovarian ca. OVCAR-3	27.2
Thymus	10.1	Ovarian ca. OVCAR-4	3.2
Spleen	1.5	Ovarian ca. OVCAR-5	9.8
Lymph node	1.9	Ovarian ca. OVCAR-8	25.5
Colorectal	1.6	Ovarian ca. IGROV-1	12.1
Stomach	4.2	Ovarian ca.* (ascites) SK-OV-3	10.6
Small intestine	5.2	Uterus	2.6
Colon ca. SW480	10.7	Placenta	7.9
Colon ca.* (SW480 met)SW620	31.7	Prostate	6.2
Colon ca. HT29	41.9	Prostate ca.* (bone met)PC-3	13.2
Colon ca. HCT-116	13.9	Testis	11.8
Colon ca. CaCo-2	25.6	Melanoma Hs688(A).T	5.2
83219 CC Well to Mod Diff (ODO3866)	51.7	Melanoma* (met) Hs688(B).T	6.1
Colon ca. HCC-2998	11.9	Melanoma UACC-62	1.0
Gastric ca.* (liver met) NCI-N87	44.6	Melanoma M14	2.3
Bladder	100.0	Melanoma LOX IMVI	1.1
Trachea	23.9	Melanoma* (met) SK-MEL-5	13.3
Kidney	34.9	Adipose	10.5

Table 60. Panel 2D

Tissue Name	Relative Expression(%) 2dx4tm4680f_ ag2771_a2	Tissue Name	Relative Expression(%) 2dx4tm4680f_ ag2771_a2
Normal Colon GENPAK 061003	52.3	Kidney NAT Clontech 8120608	1.9
83219 CC Well to Mod Diff (ODO3866)	5.0	Kidney Cancer Clontech 8120613	5.4
83220 CC NAT (ODO3866)	5.0	Kidney NAT Clontech 8120614	2.4
83221 CC Gr.2 rectosigmoid (ODO3868)	2.8	Kidney Cancer Clontech 9010320	2.4
83222 CC NAT (ODO3868)	1.4	Kidney NAT Clontech 9010321	5.2
83235 CC Mod Diff (ODO3920)	17.6	Normal Uterus GENPAK 061018	2.5
83236 CC NAT (ODO3920)	5.7	Uterus Cancer GENPAK 064011	17.1
83237 CC Gr.2 ascend colon (ODO3921)	19.3	Normal Thyroid Clontech A+ 6570-1	12.9
83238 CC NAT (ODO3921)	7.3	Thyroid Cancer GENPAK 064010	16.9
83241 CC from Partial Hepatectomy (ODO4309)	18.0	Thyroid Cancer INVITROGEN A302152	11.4
83242 Liver NAT (ODO4309)	5.6	Thyroid NAT INVITROGEN A302153	17.6
87472 Colon mets to lung (OD04451-01)	4.7	Normal Breast GENPAK 061019	27.2
87473 Lung NAT (OD04451-02)	8.4	84877 Breast Cancer (OD04566)	26.0
Normal Prostate Clontech A+ 6546-1	39.8	85975 Breast Cancer (OD04590-01)	52.2
84140 Prostate Cancer (OD04410)	22.5	85976 Breast Cancer Mets (OD04590-03)	23.4
84141 Prostate NAT (OD04410)	32.5	87070 Breast Cancer Metastasis (OD04655-05)	22.5
87073 Prostate Cancer (OD04720-01)	18.4	GENPAK Breast Cancer 064006	19.3
87074 Prostate NAT (OD04720-02)	19.5	Breast Cancer Res. Gen. 1024	20.9
Normal Lung GENPAK 061010	23.5	Breast Cancer Clontech 9100266	11.1
83239 Lung Met to Muscle (ODO4286)	1.4	Breast NAT Clontech 9100265	6.9
83240 Muscle NAT (ODO4286)	5.2	Breast Cancer INVITROGEN A209073	23.8
84136 Lung Malignant Cancer (OD03126)	17.4	Breast NAT INVITROGEN A2090734	17.2
84137 Lung NAT (OD03126)	25.0	Normal Liver GENPAK 061009	4.1
84871 Lung Cancer (OD04404)	9.2	Liver Cancer GENPAK 064003	3.4
84872 Lung NAT (OD04404)	7.9	Liver Cancer Research Genetics RNA 1025	1.4



84875 Lung Cancer (OD04565)	3.2	Liver Cancer Research Genetics RNA 1026	2.1
84876 Lung NAT (OD04565)	5.2	Paired Liver Cancer Tissue Research Genetics RNA 6004-T	4.1
85950 Lung Cancer (OD04237-01)	15.9	Paired Liver Tissue Research Genetics RNA 6004-N	3.0
85970 Lung NAT (OD04237-02)	8.5	Paired Liver Cancer Tissue Research Genetics RNA 6005-T	2.2
83255 Ocular Mel Met to Liver (ODO4310)	5.2	Paired Liver Tissue Research Genetics RNA 6005-N	0.6
83256 Liver NAT (ODO4310)	3.8	Normal Bladder GENPAK 061001	37.1
84139 Melanoma Mets to Lung (OD04321)	7.7	Bladder Cancer Research Genetics RNA 1023	2.4
84138 Lung NAT (OD04321)	21.5	Bladder Cancer INVITROGEN A302173	14.4
Normal Kidney GENPAK 061008	42.4	87071 Bladder Cancer (OD04718-01)	11.7
83786 Kidney Ca, Nuclear grade 2 (OD04338)	26.6	87072 Bladder Normal Adjacent (OD04718-03)	4.8
83787 Kidney NAT (OD04338)	11.5	Normal Ovary Res. Gen.	1.5
83788 Kidney Ca Nuclear grade 1/2 (OD04339)	24.5	Ovarian Cancer GENPAK 064008	22.4
83789 Kidney NAT (OD04339)	29.5	87492 Ovary Cancer (OD04768-07)	48.4
83790 Kidney Ca, Clear cell type (OD04340)	43.4	87493 Ovary NAT (OD04768-08)	1.7
83791 Kidney NAT (OD04340)	26.5	Normal Stomach GENPAK 061017	12.0
83792 Kidney Ca, Nuclear grade 3 (OD04348)	1.4	Gastric Cancer Clontech 9060358	0.7
83793 Kidney NAT (OD04348)	18.8	NAT Stomach Clontech 9060359	5.3
87474 Kidney Cancer (OD04622-01)	2.6	Gastric Cancer Clontech 9060395	5.6
87475 Kidney NAT (OD04622-03)	3.1	NAT Stomach Clontech 9060394	6.2
85973 Kidney Cancer (OD04450-01)	100.0	Gastric Cancer Clontech 9060397	11.7
85974 Kidney NAT (OD04450-03)	34.1	NAT Stomach Clontech 9060396	0.7
Kidney Cancer Clontech 8120607	1.9	Gastric Cancer GENPAK 064005	22.9

Table 61. Panel 4D

Tissue Name	Relative Expression(%)	Tissue Name	Relative Expression(%)
-------------	------------------------	-------------	------------------------

	4dx4tm4548f_ ag2771 b2		4dx4tm4548f_ ag2771 b2
93768_Secondary Th1_anti- CD28/anti-CD3	0.1	93100_HUVEC (Endothelial) IL-1b	54.2
93769_Secondary Th2_anti- CD28/anti-CD3	0.3	93779_HUVEC (Endothelial) IFN gamma	50.4
93770_Secondary Tr1_anti- CD28/anti-CD3	0.0	93102_HUVEC (Endothelial) TNF alpha + IFN gamma	15.6
93573_Secondary Th1_resting day 4-6 in IL-2	0.0	93101_HUVEC (Endothelial) TNF alpha + IL4	25.7
93572_Secondary Th2_resting day 4-6 in IL-2	0.3	93781_HUVEC (Endothelial) IL-11	23.3
93571_Secondary Tr1_resting day 4-6 in IL-2	0.0	93583_Lung Microvascular Endothelial Cells_none	71.5
93568_primary Th1_anti- CD28/anti-CD3	0.0	93584_Lung Microvascular Endothelial Cells_TNFa (4 ng/ml) and IL1b (1 ng/ml)	15.2
93569_primary Th2_anti- CD28/anti-CD3	0.2	92662_Microvascular Dermal endothelium_none	75.3
93570_primary Tr1_anti- CD28/anti-CD3	0.6	92663_Microvascular Dermal endothelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	44.2
93565_primary Th1_resting dy 4-6 in IL-2	1.6	93773_Bronchial epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml) **	21.1
93566_primary Th2_resting dy 4-6 in IL-2	1.4	93347_Small Airway Epithelium_none	14.4
93567_primary Tr1_resting dy 4-6 in IL-2	0.5	93348_Small Airway Epithelium_TNFa (4 ng/ml) and IL1b (1 ng/ml)	5.3
93351_CD45RA CD4 lymphocyte_anti-CD28/anti- CD3	0.8	92668_Coronary Artery SMC_resting	47.2
93352_CD45RO CD4 lymphocyte_anti-CD28/anti- CD3	1.0	92669_Coronary Artery SMC_TNFa (4 ng/ml) and IL1b (1 ng/ml)	5.0
93251_CD8 Lymphocytes_anti- CD28/anti-CD3	0.5	93107_astrocytes_resting	46.9
93353_chronic CD8 Lymphocytes 2ry_resting dy 4- 6 in IL-2	2.1	93108_astrocytes_TNFa (4 ng/ml) and IL1b (1 ng/ml)	32.0
93574_chronic CD8 Lymphocytes 2ry_activated CD3/CD28	0.0	92666_KU-812 (Basophil) resting	3.2
93354_CD4 none	0.8	92667_KU-812 (Basophil) PMA/ionoycin	11.8
93252_Secondary Th1/Th2/Tr1_anti-CD95 CH11	0.2	93579_CCD1106 (Keratinocytes)_none	15.5
93103_LAK cells_resting	0.4	93580_CCD1106 (Keratinocytes)_TNFa and IFNg **	11.6
93788_LAK cells_IL-2	0.5	93791_Liver Cirrhosis	5.7

93787_LAK cells_IL-2+IL-12	0.8	93792_Lupus Kidney	13.5
93789_LAK cells_IL-2+IFN gamma	1.5	93577_NCI-H292	55.3
93790_LAK cells_IL-2+IL-18	1.0	93358_NCI-H292_IL-4	57.1
93104_LAK cells_PMA/ionomycin and IL-18	1.7	93360_NCI-H292_IL-9	92.5
93578_NK Cells_IL-2 resting	0.1	93359_NCI-H292_IL-13	37.0
93109_Mixed Lymphocyte Reaction_Two Way MLR	1.3	93357_NCI-H292_IFN gamma	84.6
93110_Mixed Lymphocyte Reaction_Two Way MLR	0.7	93777_HPAEC_-	31.5
93111_Mixed Lymphocyte Reaction_Two Way MLR	0.2	93778_HPAEC_IL-1 beta/TNA alpha	15.7
93112_Mononuclear Cells (PBMCs)_resting	1.1	93254_Normal Human Lung Fibroblast_none	3.4
93113_Mononuclear Cells (PBMCs)_PWM	14.1	93253_Normal Human Lung Fibroblast_TNFa (4 ng/ml) and IL-1b (1 ng/ml)	2.6
93114_Mononuclear Cells (PBMCs)_PHA-L	12.2	93257_Normal Human Lung Fibroblast_IL-4	4.9
93249_Ramos (B cell)_none	0.0	93256_Normal Human Lung Fibroblast_IL-9	3.8
93250_Ramos (B cell)_ionomycin	0.1	93255_Normal Human Lung Fibroblast_IL-13	3.2
93349_B lymphocytes_PWM	25.6	93258_Normal Human Lung Fibroblast_IFN gamma	6.2
93350_B lymphocytes_CD40L and IL-4	3.1	93106_Dermal Fibroblasts CCD1070_resting	6.8
92665_EOL-1 (Eosinophil)_dbcAMP differentiated	0.1	93361_Dermal Fibroblasts CCD1070_TNF alpha 4 ng/ml	5.3
93248_EOL-1 (Eosinophil)_dbcAMP/PMAionomycin	0.1	93105_Dermal Fibroblasts CCD1070_IL-1 beta 1 ng/ml	3.1
93356_Dendritic Cells_none	3.2	93772_dermal fibroblast_IFN gamma	2.2
93355_Dendritic Cells_LPS 100 ng/ml	2.2	93771_dermal fibroblast_IL-4	6.1
93775_Dendritic Cells_anti-CD40	1.6	93260_IBD Colitis 2	1.1
93774_Monocytes_resting	0.7	93261_IBD Crohns	6.2
93776_Monocytes_LPS 50 ng/ml	1.0	735010_Colon_normal	29.5
93581_Macrophages_resting	5.2	735019_Lung_none	20.0
93582_Macrophages_LPS 100 ng/ml	1.1	64028-1_Thymus_none	60.1
93098_HUVEC (Endothelial)_none	50.9	64030-1_Kidney_none	16.6
93099_HUVEC (Endothelial)_starved	100.0		

**Table 62. Panel CNS\_neurodegeneration\_v1.0**

Tissue Name	Relative Expression(%) tm7061f_ag27 71_b1s2	Tissue Name	Relative Expression(%) tm7061f_ag27 71_b1s2
106655_4951 Hippo	15.5	106677_4624 BA21	7.7
106657_4986 Hippo	35.1	106681_4640 BA21	50.4
106652_4933 Hippo	11.5	106654_4951 BA17	20.3
106649_4901 Hippo	16.4	cns_water	0.0
110138_3087 hippo	72.0	106651_4933 BA17	8.5
110121_3027 Hippo	<b>100.0</b>	106648_4901 BA17	27.5
106670_4971 Hippo	33.6	110123_3027 Occ Ctx	27.8
106666_4867 Hippo	22.9	110140_3087 occ ctx	38.3
106680_4624 Hippo	15.0	106659_4595 BA17	4.5
106653_4951 BA21	27.1	106668_4971 BA17	29.0
106656_4986 BA21	43.7	106662_4737 BA17	17.4
106650_4933 BA21	10.0	106665_4867 BA17	11.4
106647_4901 BA21	21.3	106675_3975 BA17	83.9
110136_3087 inf temp ctx	89.6	106672_3954 BA17	16.4
110137_3087 sup temp ctx	63.8	106678_4624 BA17	5.3
110118_3027 Inf Temp Ctx	77.7	106682_4640 BA17	17.5
110119_3027 Sup Temp Ctx	80.4	106660_4595 BA7	9.5
106658_4595 BA21	10.5	113670_106669 pool	62.9
106667_4971 BA21	31.8	106663_4737 BA7	11.5
106661_4737 BA21	20.8	106676_3975 BA7	53.6
106664_4867 BA21	13.2	106673_3954 BA7	28.4
106674_3975 BA21	72.4	106679_4624 BA7	7.7
106671_3954 BA21	48.4	106683_4640 BA7	35.3

5           **Panel 1.3D Summary Ag2771** Expression of the NOV9 gene is highest in normal bladder (CT=27.2). This gene is more highly expressed in colon cancer cell lines relative to normal colon as well as in some lung cancer cell lines relative to normal lung. Thus, expression of this gene could be used to distinguish between colon or lung cancer cell lines and other cell lines. Furthermore, therapeutic inhibition of the NOV9 gene or its protein

10           product, through the use of antibodies, small molecule or protein drugs, may be effective in the treatment of colon and lung cancers.

          The NOV9 gene is expressed at moderate levels throughout CNS, with expression detected in fetal brain, amygdala, cerebellum, hippocampus, substantia nigra, thalamus,

cerebral cortex and spinal cord (CTs=29-33). See CNS\_neurodegeneration\_panel\_v1.0 summary for potential utility of this gene in CNS disorders.

Among tissues with metabolic function, this gene shows low expression in pancreas, adrenal gland, pituitary gland, and liver (adult and fetal) with higher expression in thyroid, heart (adult and fetal), skeletal muscle (adult and fetal), and adipose. Therefore, the NOV9 gene product may play a role in the pathogenesis and/or treatment of metabolic diseases in any or all of these tissues, including obesity and diabetes.

**Panel 2D Summary Ag2771** Highest expression of the NOV9 gene is found in a kidney cancer sample (CT=23). However, this gene is rather ubiquitously expressed at moderate levels in all the tissue samples on this panel. Interestingly, the level of NOV9 gene expression appears to be lower in liver and lung tissues when compared to other organs. In addition, this gene appears to be overexpressed in ovarian cancers as well as in several colon cancers relative to the normal controls. The expression patterns suggest that this gene is required for the survival and proliferation of the majority of cell types.

**Panel 4D Summary Ag2771** The NOV9 gene encodes protein with homology to CDC-42-interacting protein 4 and is highly expressed in resting HUVEC endothelial cells (CT=24-25), lung microvascular endothelium, bronchial epithelium, small airway epithelium, coronary artery smooth muscle cells, as well as in mucoepidermoid cells (NCI-H292). Basal expression of this gene in these cells appears to be decreased by various treatments with proinflammatory cytokines, such as IL-1beta, IL-4, IL-11, IFN-gamma, and TNF-alpha. CDC-42-interacting protein 4 is a Cdc42 effector protein involved in cytoskeletal organization (ref. 1-2). Since cytokine-activated cells express lower levels of the NOV9 gene, increasing the activity of this gene product may reduce the pro-inflammatory effects of these cytokines. Therefore, the NOV9 gene product may be a useful target for agonistic small molecule therapeutics that increase activity of the protein, and such small molecule drugs may reduce the severity of symptoms of asthma and inflammatory bowel disease.

**Panel CNS\_neurodegeneration\_v1.0 Summary Ag2771** CDC42 has been implicated as a neuronal death effector in Alzheimer's disease and as playing an essential role in cerebellar granule neuron survival (refs. 4-6). Cdc42-interacting protein 4, which is homologous to the NOV9 gene, has been identified as a substrate of CDC42 (ref. 2). Therefore, drugs that inhibit NOV9 gene product activity may be effective in blocking processes downstream of CDC42, such as neuronal death in Alzheimer's disease. Since

CDC42 can mediate both desirable (cerebellar neuronal survival) and undesirable (Alzheimer's disease) processes, it is likely that specifically targeting distinct downstream substrates may enable the effective targeting of distinct processes without affecting other CDC42-mediated processes. The NOV9 gene is expressed in the human brain in all regions examined, including the cerebral cortex, hippocampus, amygdala, cerebellum, substantia nigra, spinal cord and thalamus (see Panel 1.3D). Additionally, this gene is expressed more highly in the hippocampus of some patients with Alzheimer's disease than in normal control brains, indicating a possible pathological role in neurodegenerative brain disease. Therefore, targeting the NOV9 gene product may have utility in selective targeting of undesirable CDC42-mediated processes, such as Alzheimer's disease.

### References:

1. Linder S, Hufner K, Wintergerst U, Aepfelbacher M. Microtubule-dependent formation of podosomal adhesion structures in primary human macrophages. *J Cell Sci* 2000 Dec;113 Pt 23:4165-76.

Podosomes are unique actin-rich adhesion structures of monocyte-derived cells such as macrophages and osteoclasts. They clearly differ from other substratum-contacting organelles like focal adhesions in morphological and functional regards. Formation of podosomes has been shown to be dependent on the small GTPase CDC42Hs and its effector Wiskott-Aldrich syndrome protein (WASp). In this study, we investigated the functional relation between podosomes and the microtubule system in primary human macrophages. We demonstrate that, in contrast to focal adhesions, assembly of podosomes in macrophages and their monocytic precursors is dependent on an intact microtubule system. In contrast, experiments using Wiskott-Aldrich syndrome (WAS) macrophages indicate that the microtubule system is not reciprocally dependent on podosomes. A potential linker between podosomes and microtubules may be WASp itself, considering that microinjection of the WASp polyproline domain prevents podosome reassembly. This polyproline domain is thought to link WASp to microtubules via CDC42 interacting protein 4 (CIP4). Consistently, macrophages microinjected with CIP4 constructs deficient in either the microtubule- or the WASp-binding domain also fail to reassemble podosomes. In sum, our findings show that microtubules are essential for podosome formation in primary human macrophages and that WASp and CIP4 may be involved in this phenomenon.

2. Tian L, Nelson DL, Stewart DM. Cdc42-interacting protein 4 mediates binding of the Wiskott-Aldrich syndrome protein to microtubules. *J Biol Chem* 2000 Mar 17;275(11):7854-61

The Wiskott-Aldrich syndrome is an inherited X-linked immunodeficiency characterized by thrombocytopenia, eczema, and a tendency toward lymphoid malignancy. Lymphocytes from affected individuals have cytoskeletal abnormalities, and monocytes show impaired motility.

The Wiskott-Aldrich syndrome protein (WASP) is a multi-domain protein involved in cytoskeletal organization. In a two-hybrid screen, we identified the protein Cdc42-interacting protein 4 (CIP4) as a WASP interactor. CIP4, like WASP, is a Cdc42 effector protein involved in cytoskeletal organization. We found that the WASP-CIP4 interaction is mediated by the binding of the Src homology 3 domain of CIP4 to the proline-rich segment of WASP. Cdc42 was not required for this interaction. Co-expression of CIP4 and green fluorescent protein-WASP in COS-7 cells led to the association of WASP with microtubules. In vitro experiments showed that CIP4 binds to microtubules via its NH(2) terminus. The region of CIP4 responsible for binding to active Cdc42 was localized to amino acids 383-417, and the mutation I398S abrogated binding. Deletion of the Cdc42-binding domain of CIP4 did not affect the colocalization of WASP with microtubules in vivo. We conclude that CIP4 can mediate the association of WASP with microtubules. This may facilitate transport of WASP to sites of substrate adhesion in hematopoietic cells.

3. Aspenstrom P. A Cdc42 target protein with homology to the non-kinase domain of FER has a potential role in regulating the actin cytoskeleton. *Curr Biol* 1997 Jul 1;7(7):479-87

BACKGROUND: Members of the Rho family of small GTPases have been shown to have a diverse role in cell signalling events. They were originally identified as proteins that, by regulating the assembly of the actin cytoskeleton, are important determinants of cell morphology, and have recently been shown to be involved in transcriptional activation by the JNK/SAPK signalling pathway. In order to understand the mechanisms underlying the effects of Rho GTPases on these processes, the yeast two-hybrid system has been used to identify

proteins that bind to an activated mutant of Cdc42, a Rho-family member. RESULTS: A cDNA encoding a previously unidentified Cdc42 target protein, CIP4, which is 545 amino-acids long and contains an SH3 domain at its carboxyl terminus, was cloned from a human B-cell library. The amino terminus of CIP4 bears resemblance to the non-kinase domain of the FER and Fes/Fps family of tyrosine kinases. In addition, similarities to a number of proteins with roles in regulating the actin cytoskeleton were noticed. CIP4 binds to activated Cdc42 in vitro and in vivo and overexpression of CIP4 in Swiss 3T3 fibroblasts reduces the amount of stress fibres in these cells. Moreover, coexpression of activated Cdc42 and CIP4 leads to clustering of CIP4 to a large number of foci at the dorsal side of the cells. CONCLUSIONS: CIP4 is a downstream target of activated GTP-bound Cdc42, and is similar in sequence to proteins involved in signalling and cytoskeletal control. Together, these findings suggest that CIP4 may act as a link between Cdc42 signalling and regulation of the actin cytoskeleton.

PMID: 9210375

15

4. Mota M, Reeder M, Chernoff J, Bazenet CE. Evidence for a role of mixed lineage kinases in neuronal apoptosis. *J Neurosci* 2001 Jul 15;21(14):4949-57

20

Superior cervical ganglion (SCG) sympathetic neurons die by apoptosis when deprived of nerve growth factor (NGF). It has been shown previously that the induction of apoptosis in these neurons at NGF withdrawal requires both the activity of the small GTP-binding protein Cdc42 and the activation of the c-Jun N-terminal kinase (JNK) pathway. The mixed lineage kinase 3 (MLK3) belongs to a family of mitogen-activated protein (MAP) kinase kinase kinases. MLK3 contains a Cdc42/Rac interactive-binding (CRIB) domain and activates both the JNK and the p38 MAP kinase pathways. In this study the role of MLK3 in the induction of apoptosis in sympathetic neurons has been investigated. Overexpression of an active MLK3 induces activation of the JNK pathway and apoptosis in SCG neurons. In addition, overexpression of kinase dead mutants of MLK3 blocks apoptosis as well as c-Jun phosphorylation induced by NGF deprivation. More importantly, MLK3 activity seems to increase by 5 hr after NGF withdrawal in both differentiated PC12 cells and SCG neurons. We also show that MLK3 lies downstream of Cdc42 in the neuronal death pathway. Regulation of MLK3 in neurons seems to be dependent on MLK3 activity and possibly on an additional cellular component, but not on its binding to Cdc42. These results suggest that MLK3, or a closely related kinase, is a physiological element of NGF withdrawal-induced activation of the

25

30



Cdc42-c-Jun pathway and neuronal death. MLK3 therefore could be an interesting therapeutic target in a number of neurodegenerative diseases involving neuronal apoptosis.

PMID: 11438570

5

5. Linseman DA, Laessig T, Meintzer MK, McClure M, Barth H, Aktories K, Heidenreich KA. An essential role for Rac/Cdc42 GTPases in cerebellar granule neuron survival. *J Biol Chem* 2001 Aug 16; [epub ahead of print]

10 Rho family GTPases are critical molecular switches that regulate the actin cytoskeleton and cell function. In the current study, we investigated the involvement of Rho GTPases in regulating neuronal survival using primary cerebellar granule neurons. *C. difficile* toxin B, a specific inhibitor of Rho, Rac and Cdc42, induced apoptosis of granule neurons characterized by c-Jun phosphorylation, caspase-3 activation and nuclear condensation. Serum and  
15 depolarization-dependent survival signals could not compensate for the loss of GTPase function. Unlike trophic factor withdrawal, toxin B did not affect the anti-apoptotic kinase Akt or its target glycogen synthase kinase-3beta. The pro-apoptotic effects of toxin B were mimicked by *C. sordellii* lethal toxin, a selective inhibitor of Rac/Cdc42. Although Rac/Cdc42 GTPase inhibition led to F-actin disruption, direct cytoskeletal disassembly with *C. botulinum*  
20 C2 toxin was insufficient to induce c-Jun phosphorylation or apoptosis. Granule neurons expressed high basal JNK and low p38 MAPK activities that were unaffected by toxin B. However, pyridyl imidazole inhibitors of JNK/p38 attenuated c-Jun phosphorylation. Moreover, both pyridyl imidazoles and adenoviral dominant-negative c-Jun attenuated apoptosis, suggesting that JNK/c-Jun signaling was required for cell death. The results indicate  
25 that Rac/Cdc42 GTPases, in addition to trophic factors, are critical for survival of cerebellar granule neurons.

PMID: 11509562

30 6. Zhu X, Raina AK, Boux H, Simmons ZL, Takeda A, Smith MA. Activation of oncogenic pathways in degenerating neurons in Alzheimer disease. *Int J Dev Neurosci* 2000 Jul-Aug;18(4-5):433-7

A number of recent findings have highlighted the similarities between neurogenesis during

development and neurodegeneration during Alzheimer disease. In fact, neuronal populations that are known to degenerate in Alzheimer disease exhibit phenotypic changes characteristic of cells re-entering the cell division cycle. In this study, we extended these findings by investigating components of the cell cycle, known to trigger progression through G1 through activation of signal transduction cascades. Specifically, we found that proteins implicated in G1 transition, namely Cdc42/Rac, are upregulated in select neuronal populations in cases of Alzheimer disease in comparison to age-matched controls. Importantly, Cdc42/Rac shows considerable overlap with early cytoskeletal abnormalities suggesting that these changes are an extremely proximal event in the pathogenesis of the disease. Given the functional role of Cdc42/Rac in various cellular processes known to be perturbed in Alzheimer disease, namely cytoskeletal organization, oxidative balance, and oncogenic signaling, it is likely that increased neuronal Cdc42/Rac is highly significant in relation to the pathogenic process and contributes to neuronal degeneration. In fact, these findings suggest that Alzheimer disease is an oncogenic process.

PMID: 10817927

#### NOV10a

Expression of gene NOV10a was assessed using the primer-probe set Ag1674 described in Table 63. Results from RTQ-PCR runs are shown in Tables 64, 65, and 66.

**Table 63. Probe Name Ag1674**

Primers	Sequences	TM	Length	Start Position	SEQ ID NO:
Forward	5' - CTCCTCACCACAAGGGAGTAA - 3'	59.3	22	519	156
Probe	FAM - 5' - TGACATCAAACCTAACAGTCCCCAGGA - 3' - TAMRA	68.8	27	548	157
Reverse	5' - GTCTAGGAGAGAGCTGAGCAAA - 3'	58.1	22	576	158

**Table 64. Panel CNS\_1**

Tissue Name	Relative Expression(%)	
	cns1x4tm6180f_ag1674_b2	cns1tm6571f_ag1674
102633_BA4 Control	1.3	3.6
102641_BA4 Control2	0.8	3.7

102625_BA4 Alzheimer's2	2.4	0.7
102649_BA4 Parkinson's	15.3	12.4
102656_BA4 Parkinson's2	30.0	23.0
102664_BA4 Huntington's	4.7	8.2
102671_BA4 Huntington's2	5.7	1.0
102603_BA4 PSP	3.3	3.9
102610_BA4 PSP2	0.9	4.2
102588_BA4 Depression	5.2	4.2
102596_BA4 Depression2	1.6	4.0
102634_BA7 Control	3.7	3.7
102642_BA7 Control2	0.0	3.7
102626_BA7 Alzheimer's2	0.0	0.8
102650_BA7 Parkinson's	1.5	15.2
102657_BA7 Parkinson's2	21.1	21.3
102665_BA7 Huntington's	7.9	3.3
102672_BA7 Huntington's2	14.7	10.3
102604_BA7 PSP	11.5	7.4
102611_BA7 PSP2	2.2	2.4
102589_BA7 Depression	3.0	2.5
102632_BA9 Control	0.0	2.5
102640_BA9 Control2	8.7	6.8
102617_BA9 Alzheimer's	2.4	0.8
102624_BA9 Alzheimer's2	3.6	5.4
102648_BA9 Parkinson's	3.6	8.4
102655_BA9 Parkinson's2	21.1	33.0
102663_BA9 Huntington's	15.7	12.5
102670_BA9 Huntington's2	6.7	8.4
102602_BA9 PSP	5.0	5.5
102609_BA9 PSP2	0.7	0.0
102587_BA9 Depression	1.0	2.9
102595_BA9 Depression2	3.8	4.8
102635_BA17 Control	2.8	7.7
102643_BA17 Control2	0.7	3.1
102627_BA17 Alzheimer's2	1.3	2.7
102651_BA17 Parkinson's	11.8	6.9
102658_BA17 Parkinson's2	33.4	27.5
102666_BA17 Huntington's	4.1	5.1
102673_BA17 Huntington's2	5.7	2.0
102590_BA17 Depression	5.1	5.1
102597_BA17 Depression2	2.4	9.3
102605_BA17 PSP	5.0	14.7
102612_BA17 PSP2	2.8	0.8
102637_Sub Nigra Control	12.3	15.9
102645_Sub Nigra Control2	3.1	9.7
102629_Sub Nigra Alzheimer's2	3.6	1.5

102660_Sub Nigra Parkinson's2	<b>100.0</b>	95.9
102667_Sub Nigra Huntington's	16.4	6.4
102674_Sub Nigra Huntington's2	15.3	16.7
102614_Sub Nigra PSP2	7.2	3.7
102592_Sub Nigra Depression	2.4	1.6
102599_Sub Nigra Depression2	5.9	3.7
102636_Glob Palladus Control	3.7	6.4
102644_Glob Palladus Control2	0.9	1.1
102620_Glob Palladus Alzheimer's	3.5	2.7
102628_Glob Palladus Alzheimer's2	0.0	2.8
102652_Glob Palladus Parkinson's	26.1	27.2
102659_Glob Palladus Parkinson's2	21.7	38.2
102606_Glob Palladus PSP	7.4	4.8
102613_Glob Palladus PSP2	0.3	0.9
102591_Glob Palladus Depression	3.9	3.7
102638_Temp Pole Control	0.7	1.9
102646_Temp Pole Control2	2.6	4.2
102622_Temp Pole Alzheimer's	0.5	3.9
102630_Temp Pole Alzheimer's2	1.8	2.1
102653_Temp Pole Parkinson's	6.2	11.3
102661_Temp Pole Parkinson's2	16.5	23.7
102668_Temp Pole Huntington's	1.0	4.0
102607_Temp Pole PSP	1.5	4.0
102615_Temp Pole PSP2	1.4	0.9
102600_Temp Pole Depression2	2.3	1.7
102639_Cing Gyr Control	9.0	23.5
102647_Cing Gyr Control2	6.2	3.8
102623_Cing Gyr Alzheimer's	4.0	6.5
102631_Cing Gyr Alzheimer's2	1.5	1.7
102654_Cing Gyr Parkinson's	0.0	15.8
102662_Cing Gyr Parkinson's2	46.1	<b>100.0</b>
102669_Cing Gyr Huntington's	14.0	10.9
102676_Cing Gyr Huntington's2	14.9	13.6
102608_Cing Gyr PSP	16.1	32.8
102616_Cing Gyr PSP2	1.8	0.8
102594_Cing Gyr Depression	3.2	4.9
102601_Cing Gyr Depression2	6.6	11.0

**Table 65. Panel CNS\_1.1**

Tissue Name	Relative Expression(%)	
	cns_1.1tm6733 f_ag1674_a2	cns_1.1tm6734 f_ag1674_a2
102601_Cing Gyr Depression2	3.0	2.4
102594_Cing Gyr Depression	1.6	0.4

102616_Cing Gyr PSP2	1.2	0.6
102608_Cing Gyr PSP	20.8	8.2
102676_Cing Gyr Huntington's2	19.7	5.5
102669_Cing Gyr Huntington's	9.6	6.6
102662_Cing Gyr Parkinson's2	45.3	<b>100.0</b>
102654_Cing Gyr Parkinson's	18.7	4.6
102631_Cing Gyr Alzheimer's2	0.0	0.4
102623_Cing Gyr Alzheimer's	4.3	0.8
102647_Cing Gyr Control2	2.8	1.2
102639_Cing Gyr Control	13.3	11.5
102600_Temp Pole Depression2	2.3	0.4
102615_Temp Pole PSP2	0.0	0.0
102607_Temp Pole PSP	1.4	0.4
102668_Temp Pole Huntington's	6.2	2.3
102661_Temp Pole Parkinson's2	10.0	10.0
102653_Temp Pole Parkinson's	6.0	0.3
102630_Temp Pole Alzheimer's2	1.3	0.0
102622_Temp Pole Alzheimer's	0.7	0.0
102646_Temp Pole Control2	3.3	2.3
102638_Temp Pole Control	4.1	0.0
102591_Glob Palladus Depression	1.4	0.8
102613_Glob Palladus PSP2	0.7	0.3
102606_Glob Palladus PSP	4.4	2.9
102659_Glob Palladus Parkinson's2	16.3	10.5
102652_Glob Palladus Parkinson's	16.6	7.0
102628_Glob Palladus Alzheimer's2	2.0	1.2
102620_Glob Palladus Alzheimer's	0.7	56.0
102644_Glob Palladus Control2	0.8	0.4
102636_Glob Palladus Control	6.3	2.3
102599_Sub Nigra Depression2	3.2	2.2
102592_Sub Nigra Depression	0.7	0.5
102614_Sub Nigra PSP2	4.5	1.4
102674_Sub Nigra Huntington's2	20.9	7.8
102667_Sub Nigra Huntington's	6.0	7.4
102660_Sub Nigra Parkinson's2	<b>100.0</b>	52.5
102629_Sub Nigra Alzheimer's2	0.7	0.7
102645_Sub Nigra Control2	2.1	1.0
102637_Sub Nigra Control	13.2	4.5
102597_BA17 Depression2	3.3	1.3
102590_BA17 Depression	1.4	1.0
102612_BA17 PSP2	0.2	0.3
102605_BA17 PSP	3.6	1.1
102673_BA17 Huntington's2	4.6	1.2
102666_BA17 Huntington's	4.8	2.8
102658_BA17 Parkinson's2	35.7	13.0

102651_BA17 Parkinson's	7.3	5.6
102627_BA17 Alzheimer's2	2.8	0.6
102643_BA17 Control2	1.9	1.1
102635_BA17 Control	0.0	3.6
102595_BA9 Depression2	0.7	0.2
102587_BA9 Depression	1.5	0.4
102609_BA9 PSP2	0.7	0.7
102602_BA9 PSP	2.7	0.6
102670_BA9 Huntington's2	7.6	3.1
102663_BA9 Huntington's	6.2	6.0
102655_BA9 Parkinson's2	26.7	10.8
102648_BA9 Parkinson's	6.3	1.3
102624_BA9 Alzheimer's2	1.3	0.4
102617_BA9 Alzheimer's	0.7	0.8
102640_BA9 Control2	4.7	2.8
102632_BA9 Control	0.0	1.9
102589_BA7 Depression	2.0	1.4
102611_BA7 PSP2	2.2	0.7
102604_BA7 PSP	7.4	5.3
102672_BA7 Huntington's2	8.4	4.3
102665_BA7 Huntington's	11.0	3.8
102657_BA7 Parkinson's2	19.8	6.9
102650_BA7 Parkinson's	3.1	2.9
102626_BA7 Alzheimer's2	0.7	1.7
102642_BA7 Control2	1.3	0.6
102634_BA7 Control	3.4	3.1
102596_BA4 Depression2	2.6	0.6
102588_BA4 Depression	0.8	2.1
102610_BA4 PSP2	1.7	0.0
102603_BA4 PSP	2.8	1.0
102671_BA4 Huntington's2	4.6	1.0
102664_BA4 Huntington's	5.5	1.7
102656_BA4 Parkinson's2	22.1	12.1
102649_BA4 Parkinson's	10.1	5.6
102625_BA4 Alzheimer's2	0.8	0.4
102641_BA4 Control2	2.6	1.3
102633_BA4 Control	3.3	1.4

Table 66. Panel CNS\_neurodegeneration\_v1.0

Tissue Name	Relative Expression(%) tm7056f_ ag1674_a2_s1	Tissue Name	Relative Expression(%) tm7056f_ ag1674_a2_s1
106655_4951 Hippo	20.8	106677_4624 BA21	2.9
106657_4986 Hippo	10.4	106681_4640 BA21	13.7

106652_4933 Hippo	3.6	106654_4951 BA17	15.2
106649_4901 Hippo	7.7	cns_water	0.0
110138_3087 hippo	45.2	106651_4933 BA17	2.3
110121_3027 Hippo	19.1	106648_4901 BA17	9.6
106670_4971 Hippo	12.9	110140_3087 occ ctx	7.7
106666_4867 Hippo	17.3	110123_3027 Occ Ctx	4.3
106680_4624 Hippo	2.2	106659_4595 BA17	2.3
106653_4951 BA21	21.6	106668_4971 BA17	16.3
106656_4986 BA21	9.1	106662_4737 BA17	8.5
106650_4933 BA21	3.4	106665_4867 BA17	10.9
106647_4901 BA21	19.7	106675_3975 BA17	22.0
110136_3087 inf temp ctx	<b>100.0</b>	106672_3954 BA17	2.4
110137_3087 sup temp ctx	51.9	106678_4624 BA17	0.4
110118_3027 Inf Temp Ctx	18.9	106682_4640 BA17	6.9
110119_3027 Sup Temp Ctx	18.1	106660_4595 BA7	4.6
106658_4595 BA21	3.7	113670_106669 pool	46.1
106667_4971 BA21	11.4	106663_4737 BA7	12.3
106661_4737 BA21	9.5	106676_3975 BA7	14.3
106664_4867 BA21	10.4	106673_3954 BA7	6.9
106674_3975 BA21	10.9	106679_4624 BA7	1.9
106671_3954 BA21	10.3	106683_4640 BA7	13.9

#### Panel CNS\_1/CNS\_1.1/Panel CNS\_neurodegeneration\_v1.0 Summary Ag1674

The NOV10a gene encodes a protein with homology to hepsin/plasma transmembrane serine proteases. This gene is more highly expressed in the substantia nigra, globus palladus, BA17, BA4, BA9 and cingulate gyrus region of a Parkinson's disease brain than in the control brains. In addition, expression of this gene is 5-fold higher in the substantia nigra and cingulate gyrus of a Parkinson's disease brain than in similar regions from the brains of patients with other neurodegenerative diseases, such as Alzheimer's or Huntington's disease; this observation suggests that overexpression of the NOV10a gene in these regions is particular to Parkinson's disease. The substantia nigra is particularly vulnerable in Parkinson's disease, indicating that the NOV10a gene product may be a component of the biological dysregulation that contributes to Parkinson's disease pathogenesis. In particular, the NOV10a gene may be involved in the neuronal death that occurs in the substantia nigra in Parkinson's disease.

The NOV10a gene is also more highly expressed in the superior/inferior temporal cortex, globus palladus, and hippocampus of an Alzheimer's brain when compared to the control brains. This observation suggests that this gene may play a role the neurodegeneration of other diseases in addition to Parkinson's disease, such as Alzheimer's disease. Based on the proposed role of transmembrane proteases such as beta secretase in neurodegenerative

disorders, agents that influence the activity of the NOV10a gene product may be useful in treating these disorders, especially Parkinson's disease in which this gene appears to be dramatically upregulated.

### 5           **Example 3. SNP analysis of NOVX clones**

**SeqCalling™ Technology:** cDNA was derived from various human samples representing multiple tissue types, normal and diseased states, physiological states, and developmental states from different donors. Samples were obtained as whole tissue, cell lines, primary cells or tissue cultured primary cells and cell lines. Cells and cell lines may have been

10   treated with biological or chemical agents that regulate gene expression for example, growth factors, chemokines, steroids. The cDNA thus derived was then sequenced using CuraGen's proprietary SeqCalling technology. Sequence traces were evaluated manually and edited for corrections if appropriate. cDNA sequences from all samples were assembled with themselves and with public ESTs using bioinformatics programs to generate CuraGen's human SeqCalling

15   database of SeqCalling assemblies. Each assembly contains one or more overlapping cDNA sequences derived from one or more human samples. Fragments and ESTs were included as components for an assembly when the extent of identity with another component of the assembly was at least 95% over 50 bp. Each assembly can represent a gene and/or its variants such as splice forms and/or single nucleotide polymorphisms (SNPs) and their combinations.

20       Variant sequences are included in this application. A variant sequence can include a single nucleotide polymorphism (SNP). A SNP can, in some instances, be referred to as a "cSNP" to denote that the nucleotide sequence containing the SNP originates as a cDNA. A SNP can arise in several ways. For example, a SNP may be due to a substitution of one nucleotide for another at the polymorphic site. Such a substitution can be either a transition or

25   a transversion. A SNP can also arise from a deletion of a nucleotide or an insertion of a nucleotide, relative to a reference allele. In this case, the polymorphic site is a site at which one allele bears a gap with respect to a particular nucleotide in another allele. SNPs occurring within genes may result in an alteration of the amino acid encoded by the gene at the position of the SNP. Intragenic SNPs may also be silent, however, in the case that a codon including a

30   SNP encodes the same amino acid as a result of the redundancy of the genetic code. SNPs occurring outside the region of a gene, or in an intron within a gene, do not result in changes in any amino acid sequence of a protein but may result in altered regulation of the expression



pattern for example, alteration in temporal expression, physiological response regulation, cell type expression regulation, intensity of expression, stability of transcribed message.

**Method of novel SNP Identification:** SNPs are identified by analyzing sequence assemblies using CuraGen's proprietary SNPTool algorithm. SNPTool identifies variation in assemblies with the following criteria: SNPs are not analyzed within 10 base pairs on both ends of an alignment; Window size (number of bases in a view) is 10; The allowed number of mismatches in a window is 2; Minimum SNP base quality (PHRED score) is 23; Minimum number of changes to score an SNP is 2/assembly position. SNPTool analyzes the assembly and displays SNP positions, associated individual variant sequences in the assembly, the depth of the assembly at that given position, the putative assembly allele frequency, and the SNP sequence variation. Sequence traces are then selected and brought into view for manual validation. The consensus assembly sequence is imported into CuraTools along with variant sequence changes to identify potential amino acid changes resulting from the SNP sequence variation. Comprehensive SNP data analysis is then exported into the SNPCalling database.

**Method of novel SNP Confirmation:** SNPs are confirmed employing a validated method known as Pyrosequencing (Pyrosequencing, Westborough, MA). Detailed protocols for Pyrosequencing can be found in: Alderborn et al. Determination of Single Nucleotide Polymorphisms by Real-time Pyrophosphate DNA Sequencing. (2000). *Genome Research*. 10, Issue 8, August. 1249-1265. In brief, Pyrosequencing is a real time primer extension process of genotyping. This protocol takes double-stranded, biotinylated PCR products from genomic DNA samples and binds them to streptavidin beads. These beads are then denatured producing single stranded bound DNA. SNPs are characterized utilizing a technique based on an indirect bioluminometric assay of pyrophosphate (PPi) that is released from each dNTP upon DNA chain elongation. Following Klenow polymerase-mediated base incorporation, PPi is released and used as a substrate, together with adenosine 5'-phosphosulfate (APS), for ATP sulfurylase, which results in the formation of ATP. Subsequently, the ATP accomplishes the conversion of luciferin to its oxi-derivative by the action of luciferase. The ensuing light output becomes proportional to the number of added bases, up to about four bases. To allow processivity of the method dNTP excess is degraded by apyrase, which is also present in the starting reaction mixture, so that only dNTPs are added to the template during the sequencing. The process has been fully automated and adapted to a 96-well format, which allows rapid screening of large SNP panels. The DNA and protein sequences for the novel single nucleotide polymorphic variants are reported. Variants are reported individually but any combination of all or a select

subset of variants are also included. In addition, the positions of the variant bases and the variant amino acid residues are underlined.

### Results

5 Variants are reported individually but any combination of all or a select subset of variants are also included as contemplated NOVX embodiments of the invention.

#### NOV1a SNP data:

NOV1a has two SNP variants, whose variant positions for its nucleotide and amino acid sequences is numbered according to SEQ ID NOs:1 and 2, respectively. The nucleotide  
10 sequence of the NOV1a variant differs as shown in Table 67.

Table 67. cSNP and Coding Variants for NOV1a				
NT Position of cSNP	Wild Type NT	Variant NT	Amino Acid position	Amino Acid Change
472	A	G	155	R->G
481	G	C	158	A->P
1121	T	C	373	V->A
1516	T	C		No change
1566	T	C	553	C->R

#### NOV4 SNP data:

15 NOV4 has four SNP variants, whose variant positions for its nucleotide and amino acid sequences is numbered according to SEQ ID NOs:13 and 14, respectively. The nucleotide sequence of the NOV4 variant differs as shown in Table 68.

Table 68. cSNP and Coding Variants for NOV7				
NT Position of cSNP	Wild Type NT	Variant NT	Amino Acid position	Amino Acid Change
347	T	A	114	H->Q

20

#### NOV7 SNP data:

NOV7 has one SNP variant, whose variant position for its nucleotide and amino acid sequences is numbered according to SEQ ID NOs:19 and 20, respectively. The nucleotide  
25 sequence of the NOV7 variant differs as shown in Table 69.

Table 69. cSNP and Coding Variants for NOV8				
NT Position of cSNP	Wild Type NT	Variant NT	Amino Acid position	Amino Acid Change
50	T	C	17	S->P
73	G	A	24	W->Stop
111	G	A	35	V->M
120	C	T	40	A->V
160	C	T		No change
177	G	A	59	G->D
238	A	G		No change
250	G	A		No change
278	A	G	93	M->V
285	G	A	96	A->T
297	A	G	122	K->E
746	A	-		Frameshift

The SNP at nucleotide 746 has a putative allele frequency of 0.250.

## OTHER EMBODIMENTS

Although particular embodiments have been disclosed herein in detail, this has been done by way of example for purposes of illustration only, and is not intended to be limiting with respect to the scope of the appended claims, which follow. In particular, it is  
5 contemplated by the inventors that various substitutions, alterations, and modifications may be made to the invention without departing from the spirit and scope of the invention as defined by the claims. The choice of nucleic acid starting material, clone of interest, or library type is believed to be a matter of routine for a person of ordinary skill in the art with knowledge of the embodiments described herein. Other aspects, advantages, and modifications considered to be  
10 within the scope of the following claims.